

Comment on cond-mat/0406502 by V. Yukalov and E. Yukalova on Coherent Nuclear Radiation

E. M. Chudnovsky¹ and D. A. Garanin²

¹*Department of Physics and Astronomy, Lehman College, City University of New York,
250 Bedford Park Boulevard West, Bronx, New York 10468-1589, U.S.A.*

²*Institut für Physik, Johannes-Gutenberg-Universität, D-55099 Mainz, Germany*

We show that the argument of Yukalov and Yukalova that dipole-dipole interaction prevents a system of magnetic dipoles from exhibiting superradiance unless assisted by a resonator is incorrect.

The e-print of Yukalov and Yukalova [1] is a review of theory and experiment on coherent electromagnetic radiation by nuclear spins. In addition, the authors suggested, in accordance with our earlier work [2], that crystals of molecular magnets can be sources of coherent microwave radiation that is essentially stronger than the one from nuclear spins. The purpose of this Comment is to correct an error that propagates through several sections of Ref. 1. It originates in Section 6.1 that states that in order for the spin superradiance (SR) to occur, the SR rate, Γ_{rad} , must exceed the rate of dephasing, Γ_2 , due to magnetic dipole interaction between the spins, electronic or nuclear. The ratio of the two rates has been computed by the authors of Ref. 1 as

$$\frac{\Gamma_{rad}}{\Gamma_2} = \frac{2}{3n_0}(k_0L)^3, \quad (1)$$

where n_0 is the number of nearest neighbors for each spin, k_0 is the wave vector of the radiation, and L is the size of the system. According to Yukalov and Yukalova, the condition $\Gamma_{rad} \gg \Gamma_2$ is in conflict with the other condition of the SR: $k_0L \ll 1$. This made them to conclude that the SR based upon magnetic dipoles was impossible unless assisted by a resonator. While we agree that the use of the resonator may greatly enhance SR, we disagree with the above argument.

We first notice that Eq. (1) does not distinguish between the SR based upon magnetic dipoles and the SR based upon electric dipoles. Thus, if the above argument was correct, it would invalidate not only the model of non-resonator SR induced by a field pulse in a crystal of molecular magnets [2] but also the conventional Dicke model [3] of SR based upon electric dipoles. The error stems from the use of

$$\Gamma_2 = n_0\rho\gamma\mu, \quad (2)$$

where ρ is the concentration of spins, γ is the gyromagnetic ratio, and μ is the magnetic moment associated with the spin. Eq. (2) is a frequency equivalent of the dipolar magnetic field exerted on any given spin by the neighboring spins. It was introduced by Abragam [4] as a measure

of the NMR linewidth when spins are randomly oriented. It is that distribution in the orientation of the magnetic dipoles that contributes to the linewidth. Random orientation is usually the case for nuclear spins – they are difficult to align even by a high field at low temperature due to the smallness of nuclear magnetic moments. The latter is not true for crystals of molecular nanomagnets, though.

In our model [2], the spins, initially aligned by strong magnetic field, preserve the phase coherence when the field is swept in the opposite direction. During the collective relaxation towards the direction of the field, the mutual orientation of the spins does not change. This is manifested by the conservation of the length of Dicke pseudospin. The same is true for the Dicke model [3]. If the electric dipoles considered by Dicke had random orientations, as Yukalov and Yukalova assumed for magnetic dipoles, there would be no Dicke superradiance.

One should also notice that the SR does not disappear when the size of the system, L , grows beyond the wavelength of the radiation, $\lambda_0 = 2\pi/k_0$. In this limit the intensity of the radiation simply oscillates on k_0L with amplitude that slowly goes to zero on increasing L [5]. Thus, the restriction on k_0L is significantly softer than $k_0L \ll 1$ assumed in Ref. 1.

The conclusion of Yukalov and Yukalova based upon equations (1) and (2) is an overkill. The dipole-dipole interaction does not prevent systems of electric or magnetic dipoles from exhibiting superradiance.

-
- [1] V. I. Yukalov and E. P. Yukalova, arXiv:cond-mat/0406502.
 - [2] E. M. Chudnovsky and D. A. Garanin, Phys. Rev. Lett. **89**, 157201 (2002).
 - [3] R. H. Dicke, Phys. Rev. **93** 99 (1954).
 - [4] A. Abragam, *The Principles of Nuclear Magnetism* (Clarendon, Oxford, 1961).
 - [5] E. M. Chudnovsky and D. A. Garanin, arXiv:cond-mat/0404239.