

The spherical $2+p$ spin glass model: an exactly solvable model for glass to spin-glass transition.

A. Crisanti and L. Leuzzi[†]

Dipartimento di Fisica, Università di Roma "La Sapienza" and
Istituto Nazionale Fisica della Materia, Unità di Roma,
and SMC, P.le Aldo Moro 2, I-00185 Roma, Italy

We present the full phase diagram of the spherical $2+p$ spin glass model with $p \geq 4$. The main outcome is the presence of a new phase with both properties of Full Replica Symmetry Breaking (FRSB) phases of discrete models, e.g., the Sherrington-Kirkpatrick model, and those of One Replica Symmetry Breaking (1RSB). The phase, which separates a 1RSB phase from FRSB phase, is described by an order parameter function $q(x)$ with a continuous part (FRSB) for $x < m$ and a discontinuous jump (1RSB) at $x = m$. This phase has a finite complexity which leads to different dynamic and static properties.

PACS numbers: 75.10.Nr, 11.30.Pb, 05.50.+q

In the last years many efforts have been devoted to the understanding of complex systems such as spin glasses, structural glasses and others. Common denominator of all these systems is a large number of stable and metastable states [1] whose complex structure determines their static or dynamic behaviors. In this framework mean-field models, and among them spherical models, represent a valuable tool of analytical and theoretical investigation since they can be largely solved. Up to now only spherical models with One Replica Symmetry Breaking (1RSB) phases were studied, mainly due to their relevance for the fragile glass transition [2, 3, 4].

To our knowledge the possibility in finite or Full Replica Symmetry Breaking (FRSB) phases in spherical models was first pointed out by Nieuwenhuizen [5] on the basis of the similarity between the replica free energy of some spherical models with multi-spin interactions and the relevant part of the free energy of the Sherrington-Kirkpatrick (SK) model [6, 7]. In this paper Nieuwenhuizen presented some results for the FRSB phase but a complete analysis was not provided. The problem was considered some years later [8] in connection with the possible different scenarios for the critical dynamics near the glass transition [9]. This paper, however, analyzed only the dynamic behavior in the 1RSB phase.

The Model. The model we shall consider is the spherical $2+p$ spin glass model without external field defined by the Hamiltonian

$$H = \sum_{i < j} J_{ij}^{(2)} s_i s_j + \sum_{i_1 < \dots < i_p} J_{i_1 \dots i_p}^{(p)} s_{i_1} \dots s_{i_p} \quad (1)$$

where $J_{i_1 i_2 \dots i_p}^{(p)}$ are uncorrelated zero mean Gaussian variables of variance

$$\overline{J_{i_1 i_2 \dots i_p}^{(p)2}} = \frac{J_p^2 p!}{2N^{p-1}}; \quad i_1 < \dots < i_p \quad (2)$$

and s_i are N continuous variables obeying the spherical constraint $\sum_{i=1}^N s_i^2 = N$. The properties of the model

strongly depend on the value of p : for $p = 3$ the model reduces to the usual spherical p -spin spin glass model in a field [3] with a low temperature 1RSB phase, while for $p \geq 4$ the model possesses an additional FRSB low-temperature phase [5]. A partial analysis of the phase space of the model was carried out in Ref. 8 where, however, only the dynamical stability of the 1RSB phase was considered leaving out a large part of the phase space and in particular the question of the transition between the 1RSB and the FRSB phases.

In this Letter we complete the study of the phase space focusing in particular on the transitions lines. We have studied the model by three complementary approaches. The first employs the replica method and analyzes the disorder-averaged logarithm of the partition function following Ref. 3. The second approach starts from the microscopic dynamics and extends the results of Ref. 8 while the latter uses the Thouless-Anderson-Palmer approach [10]. In this Letter we shall mainly follow the replica approach, discussing differences with other approaches when necessary. A complete analysis of the properties of the model is beyond the scope of the Letter and will be presented elsewhere.

Applying the standard replica method the free energy per spin f can be written as function of the symmetric $n \times n$ replica overlap matrix Q as [3]

$$f = f_0 + s(1) + \lim_{n \rightarrow 0} \frac{1}{n} \max_Q G[Q] \quad (3)$$

where f_0 is an irrelevant constant, $s(1) = (1 + \ln 2)/2$ the entropy per spin at infinite temperature $T = 1/\beta$,

$$G[Q] = \frac{1}{2} \sum_{\alpha} \tilde{x}^{\alpha} g(Q^{\alpha}) + \frac{1}{2} \ln \det Q \quad (4)$$

and

$$g(x) = \frac{2}{2} x^2 + \frac{p}{p} x^p; \quad (5)$$

where we have used the shorthand $p = (J^{(p)})^2 = 2p$. The spherical constraint is ensured by the condition $Q = \bar{Q} = 1$.

Following Parisi [11] the overlap matrix Q for a number R of steps in the replica symmetry breaking is divided into successive boxes of decreasing size p_r , with $p_0 = n$ and $p_{R+1} = 1$. The replica symmetric case and the FRSB case are obtained for $R = 0$ and $R \geq 1$, respectively. In the Parisi scheme the elements of Q are then given by

$$Q_{r+1} = Q_r = q_r; \quad r = 0; \quad ; R+1 \quad (6)$$

with $Q_{R+1} = \bar{Q}$, where the notation $\setminus = r$ means that \setminus and \setminus belong to the same box of size p_r but to two distinct boxes of size $p_{r+1} < p_r$. The matrix obtained is conveniently expressed using the function

$$x(q) = p_0 + \sum_{r=1}^R (p_{r+1} - p_r) (q - q_r) \quad (7)$$

which equals the fraction of pair of replicas with overlap less or equal to q . Inserting this structure into eqs. (3)-(5), neglecting terms of order $O(n^2)$, and replacing the sums by integrals, one gets after a little of algebra,

$$\begin{aligned} 2f = 2s(1) - 2f_0 + \int_0^1 dx x(q) \frac{d}{dq} g(q) \\ + \ln(1 - q(1)) + \int_0^1 \frac{dq}{dq^0 x(q^0)} \end{aligned} \quad (8)$$

where $q(1) = q_R$ and $q(x)$ is the inverse of $x(q)$. Maximization of f with respect to $q(x)$ leads to the self-consistent equation(s) for the order parameter function $q(x)$. Depending on the value of the coupling strengths $J^{(p)}$ and of the temperature T the function $q(x)$ displays different forms which characterize the different phases of the model. Figure 1 shows the phase diagram in the space of the "natural" parameters p and z for $p = 4$. In the following we shall limit ourselves to the case $p = 4$, however the results are qualitatively valid for any $p \geq 4$. The analysis of the figure reveals four different phases, which will be discussed in the forthcoming part of this Letter.

The Paramagnetic phase (PM). This phase exists for not too large values of coupling parameters strengths and/or high temperature and is characterized by a null order parameter function: $q(x) = 0$ in the whole range $x \in [0, 1]$. The phases become unstable above the line $z = 1$ (De Almeida-Thouless line) where the "replicon" $\lambda = 1 - z$ becomes negative. In this region for $p \geq 4$ and p not too large a 1RSB solution is also unstable and a more complex phase (FRSB) appears. Below $z = 1$ the PM phase remains stable for all values of p , similarly to what happens in the spherical p -spin model without a field [3], however as p increases a more thermodynamically favorable 1RSB phase with a non vanishing order parameter appears.




FIG. 1: Phase diagram of the spherical $2 + 4$ spin glass model. PM : Paramagnetic phase; 1RSB : one-replica symmetry breaking phase; FRSB : full replica symmetry breaking phase; 1-FRSB : one-full replica symmetry breaking. The dashed lines refers to dynamics. The m-line shown are obtained for $m = 0.5$. The continuous transition between the PM and the FRSB phases and between the FRSB and 1-FRSB phases are the same for statics and dynamics. Inset: the discontinuous transition between the FRSB and the 1-FRSB phases.

The One Replica Symmetry Breaking phase (1RSB). This phase is characterized by a step-like order parameter function $q(x) = q_1 (x - m)$ [12] and is stable as long as the replicon eigenvalue is positive:

$$\frac{1}{(1 - q_1 + m q_1)^2} \frac{d^2}{dq^2} g(q) \Big|_{q=q_1} > 0 \quad (9)$$

Maximization of f with respect to q_1 and m leads to the 1RSB equations whose solution can be conveniently expressed defining $q_1 = (1 - y)/(1 - y + m y)$ in term of the function

$$z(y) = 2y \frac{1 - y + \ln y}{1 - y} \quad (10)$$

introduced in Ref. 3 for the solution of the spherical p -spin spin glass model. For $p = 4$ the solution reads

$$\begin{aligned} z &= 2(1 - z(y)) \frac{(1 - y + m y)^4}{m^2 y (1 - y)^2} \\ z &= (2z(y) - 1) \frac{(1 - y + m y)^2}{m^2 y} \end{aligned} \quad (11)$$

By fixing the value of $m \in [0, 1]$ and varying y from $y_{min} : z(y_{min}) = 1/2 (z = 0)$ to $y_{max} : z(y_{max}) = 1/2 (1 + y_{max})$, where the replicon (9) vanishes, one obtains the so called m-lines. The transition between the PM and the 1RSB phases corresponds to the $m = 1$ -line. Along this line q_1 jumps discontinuously from zero (PM) to a finite value (1RSB) however, since $m = 1$, the thermodynamic quantities remain continuous. Inserting into (11) the value y_{max} , for which the replicon vanishes, and varying m from




FIG. 2: Schematic form of the order parameter function $q(x)$ in the 1-FRSB phase.

1 to 0 one obtains the critical line between the 1RSB and the 1-FRSB phase.

The static approach requires that f be maximal with respect to variations of m . The dynamics, on the other hand, leads to the different conditions (marginal condition)

$$\frac{1}{(1 - q_1)^2} \frac{d^2}{dq_1^2} g(q_1) = 0 \quad (12)$$

which can be stated by saying that the derivative of f with respect to m (the complexity) be maximal [13, 14, 15, 16]. As a consequence the transition lines for dynamics and statics do not coincide. Due to space limitations the equations for the dynamical transition lines will not be reported, but only drawn in Figure 1 for completeness.

The One-Full Replica Symmetry Breaking phase (1-FRSB). The analysis of the instability of the 1RSB solution reveals that in order to stabilize the phase above the line where the replicon (9) vanishes a non-zero q_0 would be needed. However in the absence of external fields the order parameter function must vanish as $x \rightarrow 0$, and hence a 1RSB solution is not possible. On the other hand the different location of the static and dynamical instability lines suggests that some sort of 1RSB-like form must survive in the solution. The way out is to look for a solution which below q_0 has a structure which vanishes as $x \rightarrow 0$. The most general form is an order parameter that has a discontinuity at $x = m$, is continuous below it and vanishes for $x = 0$:

$$q(x) = \begin{cases} q_1 & \text{for } x > m \\ q_0(x) & \text{for } x < m \end{cases} \quad (13)$$

with $q(0) = 0$ and $\lim_{x \rightarrow m^-} q(x) = q_0 \neq \lim_{x \rightarrow m^+} q(x) = q(1) = q_1$, see Figure 2. That this is the correct ansatz also follows from the numerical solution of the Parisi equations derived from stationarity of f with respect to $q(x)$ [17, 18].

The 1-FRSB equations are obtained by inserting the form (13) into the replica free energy (8) and imposing stationarity with respect to $q_0(x)$, q_1 and m . The resulting equations can be solved in terms of m -lines similarly to what done for the 1RSB case. For the $p = 4$ case the solution for the "discontinuous" part of $q(x)$ reads

$$\begin{aligned} 4 &= \frac{[1 - y + m y (1 - t)]^4}{m^2 y (1 - y)^2 (1 - t)^3 (1 + 2t)} \\ 2 &= \frac{[1 - y + m y (1 - t)]^2}{m^2 y (1 - t)^3 (1 + 2t)} [y(1 + t + t^2) - 3t^2] \end{aligned} \quad (14)$$

where

$$t = \frac{q_0}{q_1} = \frac{1 + y}{4z(y)} \frac{2z(y)}{3 - y} \quad (15)$$

and $q_1 = (1 - y)[1 - y + m y (1 - t)]$. The "continuous" part of $q(x)$ is given by

$$q = \int_0^{q_1} dq^0 [2 + 3 - 4q^0] (q^0)^2; \quad 0 \leq q \leq q_0 = tq_1 \quad (16)$$

where

$$(q) = 1 - q_1 + m(q_1 - q_0) + \int_{q_1}^{q_0} dq^0 x (q^0) \quad (17)$$

For any fixed value of $m \in [0; 1]$ these equations can be solved varying y from y_{min} : $x(y_{min}) = 0$ (transition line to 1RSB phase) up to $y = 1$ ($t(y = 1) = 1$) where the difference between q_1 and q_0 vanishes. These lines are the continuation into the 1-FRSB phase of the 1RSB m -lines. In particular the $m = 1$ line represents the transition between the 1-FRSB to the FRSB phase. For many aspects this transition is similar to the transition between the PM and the 1RSB phases, indeed q_1 jumps discontinuously from a null value (FRSB) to a finite value (1-FRSB) however the discontinuity appears at $m = 1$ so that the thermodynamic quantities are continuous across the transition. The critical $m = 1$ -line ends at the point where $q_1 = q_0$, which for $p = 4$ is

$$q_1 = q_0 = \frac{1}{4}; \quad 4 = \frac{4}{3}; \quad 2 = \frac{32}{27} \quad (18)$$

From this end-point on the transition between the FRSB and the 1-FRSB phases can only take place without a jump in the order parameter function. The value of t , the ratio between q_1 and q_0 , increases along the m -lines as one moves away from the transition line with the 1RSB phase, and the lines terminate when $t = 1$ ($q_0 = q_1$). The set of all end-points for $m \in [0; 1]$ defines the continuous critical line between the 1-FRSB and the FRSB phases, which for $p = 4$ reads

$$4 = \frac{1}{m^2} \frac{1 + 3m}{3}^4; \quad 2 = \frac{2}{3} \frac{1 + 3m}{3m}^2; \quad (19)$$

On this line $q_1 = q_0 = 1/(1 + 3m)$ and $x_c = x(q_0) = m$. It is worth to note that in going from the 1-FRSB to the FRSB phase the solution changes from stable to marginally stable, and remains marginally stable in the whole FRSB phase.

The presence of a discontinuity in the order parameter function leads to a finite complexity so that static and dynamics calculations leads to different solutions being the first associated with states of smallest (zero) complexity while the latter with states of largest complexity. As a consequence the m -lines in the two cases are different, as shown in Figure 1. We shall not report the expression for the dynamical m -line, this will be done elsewhere. The inset of the figure shows the different transition lines between the FRSB and the 1-FRSB phases. The discontinuity, and hence the complexity, vanishes on the continuous transition line and two solutions coincide on this line and in the whole FRSB phase.

The Full Replica Symmetry Breaking phase (FRSB). In this phase the order parameter function is continuous and given by eq. (16) with $q_1 = q_0$. By expressing (16) in terms of $q(x)$, instead of $x(q)$, and taking successive derivatives with respect to x a power expansion of $q(x)$ about $x = 0$ can be computed [17, 19]. For the $2 + 4m$ model it turns out that $q(x)$ contains only odd powers of x , the first of which are

$$q(x) = \frac{2^{3=2}}{3^{3/4}} x + \frac{2^{7=2}}{6^{4/4}} x^3 + \frac{13^{11=2}}{72^{3/4}} x^7 + \dots \quad (20)$$

Using this expression one can show that as the PM-FRSB transition line is approached from above $m \rightarrow 2 - 1/4^+$ then both $q_0 = q(x_c)$ and x_c vanishes linearly with m as

$$q_0 = \frac{2}{3} + O(m^2); \quad x_c = \frac{3^{3/4}}{2} + O(m^2); \quad m \rightarrow 2 - 1/4^+ \quad (21)$$

so that the transition between the PM and the FRSB phases takes place continuously without any jump in the order parameter function.

Conclusions. To summarize in this Letter we have provided the full phase diagram of the spherical $2 + p$ spin glass model with $p \leq 4$. Despite its simplicity the model has a rather rich phase. Indeed not only it presents a 1RSB phase similar to that of the spherical p -spin spin glass model and a FRSB phase similar to that of the SK model, but it also possesses a completely new phase with an order parameter made of a continuous part for $x < m - 1$ much alike the FRSB order parameter and a discontinuous jump at $x = m$ resembling the 1RSB case. To emphasize its mixed nature this phase, which separates the FRSB phase and the 1RSB phase, has been called 1-FRSB. For many aspects the 1-FRSB phase is similar to the 1RSB, in particular it is stable, at difference with discrete models as the SK model where a solution of this form is always unstable [20], and has a finite complexity counting metastable states that are strict minima of the free energy landscape.

The transition between the FRSB and the 1-FRSB phase can be either continuous (for m large enough) or discontinuous. In the first case the transition line is the continuation of the discontinuous transition between the PM and the 1RSB phases and as for the latter the discontinuity appears at $m = 1$. The presence of finite complexity in the 1-FRSB phase makes the static and dynamics transition different. The two lines join together at the end-point where the discontinuity at $m = 1$ in the order parameter function vanishes. From this point on the transition between the FRSB phase and the 1-FRSB can only take place continuously with the discontinuity of the 1-FRSB phase which vanishes at the transition. Along the continuous transition line the complexity vanishes and the static and dynamic approaches lead to the same results, in agreement with the conjecture made in Ref. 21 that the FRSB phase has vanishing complexity.

In conclusion we believe that this is a rather promising model since not only it can be fully solved, but it possesses different phases which can be fully analyzed. Moreover it possesses an interesting transition between two different glassy phases, similar to what found in some colloidal suspensions [22].

Electronic address: andrea.crisanti@phys.uniroma1.it

Electronic address: luca.leuzzi@roma1.infn.it

- [1] See for example Spin Glass Theory and Beyond, by M. Mezard, G. Parisi and M. Virasoro (World Scientific, Singapore 1987); Spin Glasses and Random Fields, edited by A. P. Young (World Scientific, Singapore 1998); Complex Behavior in Glassy Systems, edited by M. Rub and C. Perez-Vicente (Springer-Verlag, Berlin 1996); C. A. Angell, Science 267, 1924 (1995).
- [2] T. R. Kirkpatrick and D. Thirumalai, Phys. Rev. B, 36, 5388 (1987).
- [3] A. Crisanti and H.-J. Sommers, Z. Phys. B 87, 341 (1992).
- [4] For a review see: J. P. Bouchaud, L. F. Cugliandolo, J. Kurchan and M. Mezard in Spin glasses and random fields Ed. by A. P. Young (World Scientific, Singapore, 1997).
- [5] Th. M. Nieuwenhuizen, Phys. Rev. Lett. 74, 4289 (1995).
- [6] A. J. Bray and M. A. Moore, Phys. Rev. Lett. 41, 1086 (1978).
- [7] E. Pytte and J. Rudnick, Phys. Rev. B 19, 3603 (1979).
- [8] S. Cichini and A. Crisanti, Europhys. Lett. 49, 754 (2000).
- [9] W. Gotze and L. Sjorgren, J. Phys. Cond. Matt. 1, 4203 (1989).
- [10] D. J. Thouless, P. W. Anderson and R. G. Palmer, Phil. Mag. 35, 593 (1977).
- [11] G. Parisi, J. Phys. A 13, L115, 1101 and 1887 (1980).
- [12] The most general form of the order parameter function in the 1RSB phase is $q(x) = q_0(m - x) + q_1(x - m)$. However in the absence of external fields $q_0 = 0$.
- [13] A. Crisanti, H. Hømer and H.-J. Sommers Z. Phys. B 92, 257 (1993).
- [14] J. Kurchan, G. Parisi and M. A. Virasoro, J. Phys. I

- (France) 3, 1819 (1993).
- [15] A. C risanti and H -J. Som m ers J. Phys. I (France) 5, 805 (1995).
- [16] R. M onasson, Phys. Rev. Lett. 75, 2847 (1995).
- [17] H -J. Som m ers and W . D upont, J. Phys. C 17, 5785 (1984)
- [18] A. C risanti, T. R izzo, Phys. Rev. E 65, 46137 (2002).
- [19] H -J. Som m ers, J. Phys. (France) Lett. 46, L-779 (1985).
- [20] A. C risanti, L. Leuzzi, G. Parisi and T. R izzo, cond-m at/0309256, Phys. Rev B (2004) (in press).
- [21] A. C risanti, L. Leuzzi, G. Parisi and T. R izzo, Phys. Rev. Lett 92, 127203 (2004).
- [22] K. D aw son et. al, Phys. Rev. E 63, 011401 (2001).