Apparent fractal dim ensions in the HMF model

Luca Squanci¹, Dieter H.E.G $ross^2$ and Stefano Ru o^3

1. D ipartim ento di Energetica \S. Stecco", Universita di Firenze, Via SM arta, 3 I-50139, Firenze, Italy

2. Hahn Meitner Institute, Bereich SF5 Glienicker Str.100, D14109 Berlin, Germany

3. D ipartim ento di Energetica \S. Stecco" and CSDC, Universita di Firenze, INFM and INFN, V ia S.M arta, 3 I-50139, Firenze, Italy

Abstract

We show that recent observations of fractal dimensions in the -space of N-body Ham iltonian systems with long-range interactions are due to nite N and nite resolution e ects. We provide strong numerical evidence that, in the continuum (V lasov) limit, a set which initially is not a fractal (e.g. a line in 2D) remains such for all nite times. We perform this analysis for the Ham iltonian M ean Field (HMF) model, which describes the motion of a system of N fully coupled rotors. The analysis can be indirectly con med by studying the evolution of a large set of initial points for the Chirikov standard map.

Key words: Ham iltonian dynam ics, V lasov equation, Fractal dimensions.

PACS numbers: 05.45.-a Nonlinear dynamics and nonlinear dynamical systems, 05.45 D fFractals 52.65 F f Fokker-P lanck and V lasov equation.

It has been recently claimed [1,2] that structures characterized by a non integer fractaldimension $[\beta]$ form spontaneously in the -space of som eN-body Ham iltonian systems, even when starting from a set of initial conditions (points in -space) that does not show any fractality.

¹ E-m ail: luke@ dm a.uni .it

² E-m ail: gross@ hm i.de

³ E-mail: ru o@avanzi.de.uni .tt

The model analysed in Ref. [1] is the one-dimensional self-gravitating system [4] and the initial condition is a line in -space, corresponding to a vanishing virial ratio (zero velocity for all mass sheets). The model studied in Ref. [2] describes the motion of a fully coupled system of N rotors and goes under the name of H am iltonian M ean Field (HMF) model [5]; the initial condition is such that all rotors are xed at some given angle (conventionally zero) and angular velocities are uniform by spread in an interval, symmetric around zero.

There is a strong analogy between the two results since: i) both models possess a continuum mean-eld limit (N ! 1 at xed volume) for which the single particle distribution function (SPDF) f (q;p;t) evolves according to a V lasov-Poisson system [6], ii) both initial states are lines in -space when the continuum limit is taken.

In this short note we analyse the HMF model, considering the same initial state as the one studied in Ref. [2]. We give compelling numerical evidence that, as the number of rotors grows and as the resolution increases, the collection of initial points lies on a set of dimension one in the -space (which is two-dimensional for this model) at all nite times.

W e will explain that this is a som ewhat trivial consequence of the fact that the SPDF obeys a V lasov equation in the mean-eld lim it. M oreover, we will show that apparent fractal dim ensions can be indeed observed at low resolutions, justifying the observations of R ef. [2]. For com parison, we will show what happens for the standard C hirikov's map [7], which shares the sym plectic property with the HMF m odel, for which we can push the analysis to a larger num ber of initial points.

The Ham iltonian of the HMF model is

$$H_{HMF} = \frac{1}{2} \frac{x^{N}}{j=1} p_{j}^{2} + \frac{1}{2N} \frac{x^{N}}{jk=1} [1 \cos(q_{j} - q_{k})]; \qquad (1)$$

where $q_i 2$ [; [is the position (angle) of the i-th particle on a circle and p_i the corresponding conjugate variable. This system can be seen as representing particles moving on a unit circle interacting via an in nite range attractive cosine potential, or as classical XY rotors with in nite range ferror agnetic couplings (form ore details see R ef. [8]). The magnetization, de ned as

$$\overset{!}{M} (t) = (M_{x}; M_{y}) = \frac{1}{N} \sum_{j=1}^{N} (\cos q_{j}(t); \sin q_{j}(t));$$
(2)

is the main observable that characterizes the dynam ical and therm odynam ic state of the system .

In the continuum limit, that is keeping the volume (here the interval [;]) and the energy per particle xed as the number of particles N ! 1, the dynamics governed

by H am iltonian (1) is described by a V lasov equation. Indeed, the state of the nite N system can be described by a single particle time-dependent distribution function

$$f_{d}(q;p;t) = \frac{1}{N} \sum_{j=1}^{N} (q q_{j}(t);p p_{j}(t)); \qquad (3)$$

where is the D irac function. When N is large, it is natural to approximate the discrete density f_d by a continuous SPDF f (q;p;t). U sing this distribution, one can rewrite the two components of the magnetization M as

$$\frac{Z}{M_{x}}[f] = f(q;p;t) \cos q \, dq dp ;$$
(4)
$$\frac{Z}{M_{y}}[f] = f(q;p;t) \sin q \, dq dp :$$
(5)

W ithin this approximation the potential that a ects all the particles is

$$V(q)[f] = 1$$
 $\overline{M}_{x}[f] \cos q$ $\overline{M}_{y}[f] \sin q$: (6)

This potential enters the expression of the V lasov equation

$$\frac{\partial f}{\partial t} + p \frac{\partial f}{\partial q} \quad \frac{dV}{dq} [f] \frac{\partial f}{\partial p} = 0 \quad ; \tag{7}$$

which governs the spatio-tem poral evolution of the SPDF f.

The initial condition considered in [2] is $q_i = 0$; 8i and p_i random i.i.d. uniform by distributed in the interval [p;p]. As usual, we call this class of initial conditions water bags (W B). The value of p is chosen such that the energy per particle is $U = H_{HMF} = N = 0.69$ and N = 10000, but these data will be varied in this paper. For such an energy this initial state is V lasov unstable [9], and hence the SPDF evolves in time (see g.3 in Ref. [2]).

However, since initially the state is a line in the two-dimensional -space (q;p), it will remain a line for all nite times, although intricately stretched and folded. This is a trivial property related to the fact that V lasov equation de ness a ow $_t$ for the particles (points x = (q;p) in the -space continuum)

$$x(t) = t(x(0));$$
 (8)

which is a di com or sm . M oreover, since the Jacobian J = @x(t)=@x(0) is a sym – plectic m atrix, areas m odify their shape but are conserved.

The reason why the set does not appear to be a line is due to the nite particle number in the simulation. The line is stretched and folded by the two main phasespacem echanism s: the form ation of whorls around elliptic xed points and oftendrils in the hom oclinic tangle [10]. Moreover, local hyperbolicity in phase-space produces a dispersion of the initial points. As a consequence, the set of initial points, appears by visual inspection, in the course of time, to be partly distributed along lines and partly dispersed onto the two-dimensional -space (see again g.3 in Ref. [2] or g. 2 in Ref. [1]) This is why the authors of Refs. [1,2] have used fractal dimension to chacterize this set, more speci cally box dimension [1] and correlation dimension [2].

The number of boxes $N_b()$ of linear size which cover the set of $N = 10^{\circ}$ points at time t = 75 when starting the HMF model in a WB initial condition at U = 0.69 is shown as a function of resolution 1= in log-log scale in g. 1. These data are not qualitatively di erent from the ones in g. 3 of Ref. [1] and one might then t the central part of them with the Ansatz $N_b = D_0$ (since the maximal box size which contains all the set is normalized to one), obtaining D_0 1.930 0.002 and concluding that the set is a fractal, with dimension D_0 close to two, but de nitely distinct from two within num erical accuracy. This is a consequence of the set being a mixture of a set with one-dimensional features (the initial line) and two-dimensional ones caused by chaotic spread.



Fig.1.Num ber of occupied boxes of linear size versus resolution 1 = in log-log scale for theHMF model in a WB initial condition with U = 0:69 and N = 10^6 (). The short-dashed line superposed to the data is a power law twith apparent fractal dimension D₀ 1:93. The upper deviation from the power law is at $1 = 2^{10}$, i.e. roughly when the number of boxes $(1 =)^2$ is of the order of the number of points N. The dotted and long-dashed lines are the power law s with exponents 2 and 1, respectively.

However, as rst remarked in Ref. [11] the slope of the curve in g.1 is not constant and depends on the resolution. It is already evident in this gure that, as the resolution increases (! 0), the local slope tends to zero (on this scale most of the points are surrounded by very few neighbours). This is better shown in g.2a, where the discrete approximation of the local slope in logarithm is scale

$$s() = \frac{d \log_2 N_b()}{d \log_2 (1 =)};$$
(9)

is plotted as a function of for the same set of points. Going from low to high resolution, all dimensions from 2 to 0 are covered continuously and sm oothly. All are apparent fractal dimensions of a set which has indeed dimension 1. It might bok strange that one does not observe any signal of dimension 1 at any resolution. This is due to the inhom ogeneity, as shown in g.3, where one represents in black the occupied boxes at increasing resolutions. While at low resolutions all boxes are

occupied indicating a set of dimension 2, when the resolution increases, the set concentrates on a smaller set of boxes, but already at resolution $1 = 2^{\circ}$ one observes structures of dimension 1 (lines). However, their weight in the -space is negligible and, as the resolution is increased even more, the fraction of the set made by spread points dom inates and consequently dimension 0. A complete multifractal [3] analysis should be realized to highlight such inhom ogeneities.

For com parison, we have perform ed a sim ilar experim ent for the Chirikov standard map [7]

$$p_{t+1}^{i} = p_{t}^{i} + q_{t}^{i}$$
 (10)

$$q_{t+1}^{1} = q_{t}^{1} \quad K \sin(p_{t+1}^{1}) \mod 2$$
; (11)

where the index t = 0; ::: denotes time and the index i = 1; :::; N P the number of initial particles. Particles are put at t = 0, as above, at q = 0 and with momentum uniform by spread in the interval [p;p]. The phase space is closed by periodizing in 2 both the q and the p directions and norm alized in the unit square. The results of numerical experiments conducted above the chaos threshold, at K = 12, i.e. a parameter region with a mixed (chaotic and regular) phase space, are shown in g.2b (local slopes) and g.4 (occupied boxes in phase space at di erent resolutions). The behavior of all measured quantities is qualitatively the same as the one of the HM F m odel.



Fig. 2. Local slopes s() for resolutions ranging from $1 = \frac{2}{2}$ up to 2^{12} . In Figure a) we plot the results for the HMF m odel at t = 75 with W B initial condition at U = 0.69 and $N = 10^6$. Figure b) refers to the standard m ap at t = 160 for K = 1.2 and $N P = 10^6$.

Can one get from straight box-counting an indication of the presence of sets of dimension 1? To obtain this results it is enough to perform experiments at lower energies for the HMF model and at smaller values of parameter K in the standard m ap. By doing this, one decreases the fraction of phase space which is occupied by chaotic orbits and, hence, reduces particle dispersion. In g. 5 we plot the local slopes for the HMF model at U = 0:1 (g. 5a) and at K = 0.2 for the standard m ap (g. 5b). In these cases, the local slope, and hence fractal dimension, shows an evident plateau at the value 1 as the resolution increases, before dropping to zero when the individual points of the set are detected. Let us remark that the number



Fig. 3. HMF model. A set of N = 10^6 points is, initially, uniform by distributed along the p axis between p and p, such that U = $p^2=6 = 0.69$. From left to right and from top to bottom we show a portion S = [0.33;0.66[[0.36;0.50[of the -space, where a box of given linear dimension is colored black if at least one point of the initial condition is contained inside it at time t = 75. Resolution ranges from $1 = 2^7$ up to $1 = 2^{12}$.

of particles is the same as before $N = NP = 10^6$. This is a strong evidence that the line remains a line during time evolution, as expected.

Having understood that the limited number of particles is crucial in determining the results, we show in g. 6 and g. 7 the dependence of the local slopes on the number of particles for both the HMF m odel and the standard m ap. In all cases, s is an increasing function of the number of particles, and, at xed number of particles, as the resolution increases, the slope decreases. However, for a more chaotic phase space (g. 6), i.e. HMF at U = 0:69 and standard m ap at K = 1.2, the curves corresponding to the higher resolution show only a slight increase from zero slope and one cannot de nitely conclude that, as the number of particle increases, they will tend to slope 1 (while those which converge, clearly point to dimension 2). On the contrary, for a less chaotic phase space (HMF at U = 0:1 and standard m ap at K = 0.2), the curves corresponding to the higher resolutions converge to slope (dimension) 1 as the number of particles increases, approxim ately to $(1 =)^2$.⁴

⁴ For the standard m ap we have perform ed experim ents for up to 10^8 particles. We have obtained a more evident levelling of the slope as the number of particles increases and a convergence to 1 from above as the resolution subsequently increases. For the HMF model the convergence to 1 is followed by a further drop to smaller values at higher resolutions, as also shown in g. 5a: this is again a nite N e ect.



Fig. 4. Standard m ap at K = 1.2. A set of N P = 10^6 particles is, initially, uniform ly distributed along the p axis. From left to right and from top to bottom we show a portion S = [0;0:33[[0:86;1:00[of the phase space, where a box of given linear dimension is colored black if at least one particle of the initial condition is contained inside it at time t = 160. Resolution ranges from $1 = 2^7$ up to $1 = 2^{12}$.



Fig. 5. This gure show show the local slope changes for resolutions ranging from $1 = 2^8$ up to 2^{12} . In Figure a) we plot the results at t = 500 for the HMF m odel with W B initial condition corresponding to U = 0:1 and a num ber of particles N = 10^6 . Figure b) refers to the Standard M ap at t = 160 for K = 0.2 and N P = 10^6 . Again the \natural" slope s = 1 is roughly at $(1 =)^2$ N; N P.

W e think that we have solved the puzzle of the existence of a fractal set generated in a nite time when starting with a one-dimensional set (a line) in the phase space of H am iltonian system s. The fractal is not there and is a feature of the resolution with which the set is observed.

O f course this does not dim inish the role played by fractal concepts in system s with long-range interactions. Certainly, when the initial set is already a fractal [12], it's



Fig. 6. Local slope as a function of the number of particles (ranging from 10^3 up to 10^7) for di erent values of the resolution $1 = :2^7$ (4); 2^8 (); 2^2 (); 2^{10} (); 2^{11} (+); 2^{12} (). Figure a) shows the results for the HMF model at U = 0:69 and Figure b) the corresponding results for the standard map at K = 12.



Fig. 7. Same as g. 6 but for U = 0.1 (HMF) a) and K = 0.2 (standard map) b).

well possible that the set remains fractal during time evolution. Moreover, when studying evolving universe models, one electively introduces dissipative terms in the equations of motion [12,13] and one can indeed observe fractal sets generated from non fractals. A nother situation is the one of the advection of passive tracers [14] or vector elds [15], where one indeed can observe fractal sets in H am iltonian phase spaces. This is due to the presence of non escaping orbits inside the mixing region, which form a zero measure fractal repeller [14] or, in the kinematic dynam o problem [15], to the concentration of the magnetic ux onto a singular set ast ! 1.

A cknow ledgem ents

W e warm ly thank F.Bagnoli for useful discussions and help in num erical calculations. This work has been nantially supported by The University of Florence, INFN, MIUR-COFIN03 contract Order and chaos in nonlinear extended systems and the FIRB n.RBNE01CW 3M_01 project on synchronization.

- [1] H.Koyam a and T.Konishi, Phys. Lett. A 279, 226 (2001).
- [2] V.Latora, A.Rapisarda, C.Tsallis, Phys. Rev. E 64, 056134 (2001) and Physica A 305, 129 (2002).
- [3] K. Falconer, Fractal geometry: m athem atical foundations and applications, W iley (1990).
- [4] F.Hohland M.R.Feix, Astroph.J., 147, 1164 (1967).
- [5] M.Antoniand S.Ru o, Phys. Rev. E 52, 2361 (1995).
- [6] H. Spohn, Large Scale Dynam ics of Interacting Particles, Springer (1991).
- [7] B.V. Chirikov, Phys. Rep., 52, 263 (1979).
- [8] T. Dauxois, V. Latora, A. Rapisarda, S. Ru o, A. Torcini, The Ham iltonian Mean Field Model: from Dynamics to Statistical Mechanics and back, in Dynamics and thermodynamics of systems with long range interactions, T. Dauxois et al. Eds., Lecture Notes in Physics 602, Springer (2002).
- [9] Y.Y.Yam aguchi, J.Barre, F.Bouchet, T.Dauxois and S.Ru o, Physica A, 337, 36 (2004).
- [10] JM. Ottino, The kinematics of mixing: stretching, chaos and transport, Cambridge University Press (1989).
- [11] G. Benettin, D. Casati, L. Galgani and A. Giorgilli, Phys. Lett. A, 118, 325 (1986).
- [12] T. Tatekawa and K. Maeda, Astroph. J. 547, 531 (2001).
- [13] B N.M iller and J.L.Rouet, Phys.Rev.E 65, 056121 (2001).
- [14] T. Telet al, Chaos, 10, 89 (2000).
- [15] E.Ott and T.M. Antonsen Jr., Phys. Rev. A, 39, 3660 (1989).