

Josephson π -state in superconductor-Luttinger liquid hybrid systems

Nobuhiko Yokoshi and Susumu Kurihara

Department of Physics, Waseda University, Okubo, Shinjuku, Tokyo 169-8555, Japan

(Dated: April 14, 2024)

Josephson current through a Luttinger liquid (LL) under a magnetic field is theoretically studied. We derive an analytical expression of Josephson current for clean interfaces, by using quasiclassical Green's function and functional bosonization procedure. We show that critical currents can be renormalized by electron-electron interactions at perfect transparency when LL is adiabatically connected with superconductors. We also find that a generation of π -state, due to spin-dependent energy shift in Andreev bound states (ABS), is prohibited even at zero temperature when the strength of repulsive interactions reaches some critical value. The suppression of π -state is caused by the low energy fluctuations propagating in LL, and making the Zeeman splitting in ABS blurred.

PACS numbers: 71.10.Pm, 71.70.Ej, 74.50.+r

Transport phenomena in one-dimensional (1D) structures are strongly affected by electron-electron interactions. Fermi liquid description breaks down due to electron correlation, and systems are believed to behave as Luttinger liquids (LLs), where low-lying excitations are collective modes rather than single-particle excitations. This gives rise to characteristic phenomena such as spin-charge separation and charge fractionalization. These behaviors have been verified experimentally, e.g., in carbon nanotubes (CNTs) [1, 2] and quantum wires on GaAs/AlGaAs heterostructure [3, 4]. Inspired by advances in microfabrication techniques, hybrid structures connecting LL with other conductors have attracted attentions from theoretical and experimental sides as candidates of new mesoscopic electronic devices.

Recently, superconducting proximity effect on CNTs in contact with superconductors was reported in a couple of experiments [5, 6]. There are some previous works about Josephson currents in superconductor-Luttinger liquid (S-LL) hybrid systems [7, 8, 9, 10, 11, 12]. In such systems, electron-electron interactions modify the normal and the Andreev reflection (AR) process, and lead to unique power law behaviors of low energy properties such as proximity effect, conductance and local density of states (LDOS). On the other hand, a generation of metastable π -state has been studied theoretically [13] and experimentally [14, 15] in superconductor-ferromagnet (S-F) junctions. The π -state is a result of spin-dependent energy shift in Andreev bound states (ABSs) due to exchange energy in F. Here, we study Josephson current in the presence of the π -state in S-LL junctions. By applying a magnetic field to the LL region, we can consider that Zeeman energy in LL plays the same role as the exchange energy in F. In this paper, we derive the analytical expression of Josephson current and find that critical currents can be renormalized by the interactions at perfect transparency when LL is adiabatically connected with reservoirs. We also show that the Coulomb interactions affect the generation of the metastable π -state through the LDOS in 1D structures.

The system under consideration is a long S-LL-S junction where LL is adiabatically connected with s-wave su-




FIG. 1: A possible implementation of system setup: s-wave superconductors are deposited on quantum wire which is adiabatically connected with two dimensional Fermi liquids. We believe that superconducting proximity effect penetrates in Fermi liquid regions. The electrons in superconductors can tunnel into Luttinger liquid via Andreev reflection process. For sake of simplicity, barrier potentials at the junctions are assumed to be negligible.

perconductors (see Fig.1). Approximately in absence of the interactions in superconductors is neglected, and superconducting order parameters $\psi = \psi_0 e^{i\phi}$ is assumed to change abruptly. For LL region, we assume that Fermi velocity is spin-independent $v_F^\uparrow = v_F^\downarrow = v_F$. This assumption may be justified as far as the Zeeman energy is far smaller than the Fermi energy. In this case, we can consider that the spin-charge separation in LL does not break down [16]. Thus, Hamiltonian with usual g-ology is expressed as

$$H_{LL} = \sum_{a,s} \int dx \left[\frac{\hbar v_F}{2} \left(\frac{\partial}{\partial x} \psi_{a,s}^\dagger + \psi_{a,s} \right) \right] + \sum_{a,s} \int dx \left[g_{2,p} \psi_{a,s}^\dagger(x;t) \psi_{a,s}^\dagger(x;t) \right] + \sum_{a,s} \int dx \left[\frac{g_{4,p}}{2} \psi_{a,s}^\dagger(x;t) \psi_{a,s}^\dagger(x;t) \right] ; (1)$$

where $\psi_{a,s}$ and \hbar are particle density and Zeeman energy. Here, $a = \uparrow, \downarrow$ and $s = \uparrow, \downarrow$ denote direction of movement and spin, respectively. $p = \pm$ specifies interaction between electrons with parallel or anti-parallel spins. We consider the case where back-scattering and umklapp-scattering processes are irrelevant for simplic-




FIG. 2: Josephson current in unit of $J_c = ev_F L$ is plotted as a function of phase difference for different K 's (Here, we set $\alpha = 0.45$, $L=L_T = 0.4$, $K = 1$). We assume $g_{ik} = g_i$, thus $K_j^{-1} = u_j$. There are positive slopes at $\alpha = 0$ indicating the existence of the π -state. One can find critical currents are suppressed for repulsive interactions.

Here, K_j and $v_j = u_j v_F$ are usual Luttinger parameter and propagation velocity of j component.

Total Josephson current through the system can be obtained, summing up the contributions from all the ABSs. Because of the spin-charge separation, renormalization of the spin-up current and the spin-down one is exactly the same. Here we assume that excitation momentum unit of the charge and the spin bosons is infinitesimally small in LL connected with Fermi liquids. As a consequence, Josephson current for S-LL-S system has a following form

$$J = \frac{ev_F}{L} \sum_{m=1}^{\infty} \frac{2}{L_T} \frac{X^m}{\sin(m\alpha) \cos(m\alpha)} \frac{(1)^{m+1} m^2}{\sinh(\frac{mL}{L_T})}; \quad (16)$$

with $L_T = v_F/2\pi$ being thermal length. The influence of electron-electron interactions is included only in X , which has a following form

$$X = \left(\frac{T}{D}\right)^{K^{-1}+K^{-2}} \frac{u^{K^{-1}} u^K \sinh^2(\frac{L}{2L_T})}{\sinh^{K^{-1}}(\frac{L}{2uL_T}) \sinh^K(\frac{L}{2uL_T})}; \quad (17)$$

Here, D is a high energy cut-off; since we take into account the energy smaller than superconducting energy gap, we assume $D \gg \Delta$ [12]. Eq.(16) shows that each AR process is renormalized by the fluctuations with energies lower than $v_F L$ [23]. This can be interpreted that the "environment" at the interfaces directly affect the AR. Note that the exponent of renormalization differs from the one due to normal backscattering, e.g., at high barrier potential/ $(T=D)^{K^{-1}+K^{-1}+2}$ [24]. Therefore, we consider that it is Cooper pair tunneling process that the interactions renormalize. To see this explicitly, we utilize conventional tunnel Hamiltonian $H_T = \sum_{a;s} T_1 \psi_{a;s}^\dagger(\frac{L}{2}) \tilde{\psi}_{a;s}(\frac{L}{2}) + T_2 \psi_{a;s}^\dagger(\frac{L}{2}) \tilde{\psi}_{a;s}(\frac{L}{2}) + H.c.$




FIG. 3: Temperature dependence of J_π in unit of J_c is plotted at $\alpha = 0.35$ and $K = 1$. For repulsive interactions, one can see that π -state is strongly suppressed. Further, at $K = 0.4$, the generation of π -state is entirely suppressed even at zero temperature.

with $a_{a;s}(x)$ and $\tilde{a}_{a;s}(x) = e^{i(ak_F x + a_{a;s})}$ being annihilation operators in S and LL. At zero temperature, the second order perturbation of H_T at each interface, which corresponds to single AR process, is proportional to

$$\sum_Z \int dt < e^{i\alpha} \psi^\dagger(\frac{L}{2}, 0) \psi(\frac{L}{2}, 0) \psi(\frac{L}{2}, t) \psi^\dagger(\frac{L}{2}, t) > / \left(\frac{v_F}{D L}\right)^{K^{-1}+K^{-2}} = (T \neq 0): \quad (18)$$

This is in agreement with the result of Fazio et al. [7]. One can verify that higher order contributions are consistent with Eq.(16) as far as the power law dependence is concerned [18].

Current-phase relations for different values of K are shown in Fig 2 (Hereafter, we employ the parameters, such as α and L , modeling the experiment by Kasumov et al. [5]). Owing to the Zeeman phase shift α , the positive slopes are found to appear at $\alpha = 0$ (π -state). This is the same result as in S-F junction except the suppression (enhancement) by the repulsive (attractive) interactions [13]. However, the suppression by the repulsive interactions ($K < 1$) can play a key role as for the generation of π -state. We calculate the temperature dependence of the sub-critical current J_π which provides an indication of the existence of the π -state (see Fig. 3). In presence of the repulsive interactions, there is the regime for finite T where the π -state does not occur even at absolute zero, although finite Josephson current can flow. The condition for the existence of a π -state is given by

$$\sum_{m=1}^{\infty} \frac{X^m}{\cos(m\alpha)} \left(\frac{v_F}{D L}\right)^{(K^{-1}+K^{-2})m^2} > 0: \quad (19)$$

In non-interacting case ($K = K = 1$), $\alpha = 0.5$ for finite T . In contrast, if the Coulomb interactions are strong enough ($K \geq 0.2$) as in CNTs [2], the metastable π -state cannot be generated as such.




FIG. 4: The tunneling rate from the bottom of the $\bar{\sigma}$ -state is plotted for $\sigma = 0.35$ (dotted line) and $\sigma = 0.45$ (solid line). Declining line is the one plotted as a function of K at $K = 1$ (left axis), while increasing one as a function of K at $K = 0.3$ (right axis).

This behavior can be verified also from Josephson potential $E_J(\phi; \sigma)$. As a criterion for determining the existence of the metastable state, we estimate the transition probability T from the bottom of the $\bar{\sigma}$ -state. T is numerically calculated as a function of K and K within WKB method, and the result is shown in Fig. 4. One can see that there is a critical value of interaction strength, where T reaches unity and the generation of $\bar{\sigma}$ -state is prohibited completely. In the vicinity of the critical point, the tunneling probability becomes

$$T = 1 - \frac{3}{2} \frac{1}{2} \frac{r}{C} \frac{v_F}{e^2 L} ; \quad (20)$$

where C denotes junction capacitance. The disappearance of the metastable state is caused by the fact that the Zeeman splitting in ABSs becomes blurred due to the collective excitations arising from the interactions in LL. In practice, for repulsive interactions, the LDOS near the interfaces can be roughly estimated by

$$\begin{aligned} \frac{\langle \sigma \rangle}{(0)} &= \sum_{n,s} X_{n,s} 2e^{-j_{n,s}} \frac{\hbar}{(j_{n,s})^2 + \frac{1}{2}} + \frac{(v_F)}{D L} (1 - \cos \frac{f}{2}) + g \\ &+ \frac{j_{n,s}}{2D} \left(\frac{v_F}{D L} \right) + (1 - \cos \frac{f}{2}) + \frac{1}{2} g ; \end{aligned} \quad (21)$$

with the exponents $j_{\pm} = (K_{\pm}^{-1} + K_{\pm} - 2) = 4$ and $\gamma = (K_{\pm}^{-1} + K_{\pm} - 2) = 4$. Here γ is the finite spectral weight of each ABS and $\Gamma(x)$ is Gamma function. We further set the phase difference between the superconductors to zero for simplicity [12]. The first term in right hand side shows that the spectral weight is spread out through distribution function. In this light, the repulsive interactions can be interpreted to raise the effective temperature of the junction system. In the second one, which originates in $\chi_{a,s}$ of Eq.(10), the repulsive interactions suppress the Andreev resonance. For attractive case ($K > 1$), on the other hand, it becomes easier to generate the metastable state. An attractive back-scattering, in general, cannot be neglected. However, it does not change the situation qualitatively, because the attractive back-scattering tends to enhance the superconducting correlation [25]. Besides, in contrast to K , small K enhances the critical current, and makes the $\bar{\sigma}$ -state incident. This reveals a reflection of the symmetry of Cooper pairs in superconductors.

Finally, we mention the effect of breakdown of spin-charge separation due to the Zeeman effect. We have fo-

cused on the case where spin-charge separation is valid. This assumption may be acceptable for the system on electron doped GaAs which has small Lande's g -factor ($g \approx 0.4$) [4]. If, e.g., InSb quantum wires ($g \approx 50$) [26] are prepared, however, our approach needs some corrections. In such a case, the charge- and spin-density fluctuations exhibit coupled oscillation, whose coupling strength is proportional to $r = (v_F^{\uparrow} - v_F^{\downarrow}) / (v_F^{\uparrow} + v_F^{\downarrow})$. Therefore, \hat{M} in Eq.(12) has off-diagonal elements. This leads to different renormalization for spin-up and spin-down currents, and the ratio of renormalization factor has the following expression

$$\begin{aligned} \frac{r}{\#} &= \sinh\left(\frac{L}{2u_{\downarrow} L_T}\right) = \sinh\left(\frac{L}{2u_{\uparrow} L_T}\right) \frac{4r(u_{\downarrow} K^{-1} + u_{\uparrow} K)}{u_{\downarrow}^2 - u_{\uparrow}^2} \\ &\sinh\left(\frac{L}{2(1+r)L_T}\right) = \sinh\left(\frac{L}{2(1-r)L_T}\right)^2 ; \end{aligned} \quad (22)$$

where u denotes velocity renormalization of the normal mode [16]; the explicit form is given by

$$u^2 = \frac{1}{2} \frac{\hbar}{(u^2 + u^2 + 2r^2)} \frac{q}{(u^2 - u^2)^2 + 4r^2 (u K + u K^{-1}) (u K^{-1} + u K)} ; \quad (23)$$

Since Eq.(22) does not include high energy cut-off D , $\mu = \mu_0$ does not become so large. Thus the influence of the breakdown of the spin-charge separation is small, which may cause the circumstance that supercurrent is conveyed by singlet Cooper pairs.

In conclusion, current-phase relations of S-L-S junctions are analytically reexamined using functional bosonization. We have shown that the low energy excitations in LL can play crucial roles on a critical current and a generation of π -state if LL is adiabatically connected with superconducting reservoirs. We have also examined the tunneling from the metastable π -state. When we regard the junctions as macroscopic two level systems, the quantum leakage directly affects the stability of the metastable state. The decay rate is given by $\Gamma = 2B e^{S_0 E_J(\pi/2)^2}$, where $S_0 = \int_0^d dx CE_J(\pi/2) = 4e^2$ is the action of symmetric potential (B is the actual

tunneling determinant without zero mode [27, 28]). This indicates that the macroscopic nature is inevitably related with many body correlations in 1D conduction. Our method can be easily applied to hybrid systems of superconductors with other symmetries or interacting conductor with higher dimensions [21, 29]. Experimentally in semiconducting heterostructure, one can vary the carrier density and the effective interactions to some extent by gate voltage [30]. Further tunnel junctions between 1D-2DEG have been also examined [31]. It may be feasible in the future to construct the system under consideration, using S-2DEG-S junction with a gate [32].

This work is partly supported by a Grant for The 21st Century COE Program (Physics of Self-organization Systems) at Waseda University from the Ministry of Education, Sports, Culture, Science and Technology.

-
- [1] M. Bockrath, D. H. Cobden, P. L. McEuen, N. G. Chopra, A. Zettl, A. Thess and R. E. Smalley, *Science* 275, 1922 (1997); Z. Yao, H. W. C. Postma, L. Balents and C. Dekker, *Nature* 402, 273 (1999).
- [2] H. Ishii, H. Kataura, H. Shiozawa, H. Yoshioka, H. Otsubo, Y. Takayama, T. Miyahara, S. Suzuki, Y. Achiba, M. Nakatake, T. Narimura, M. Higashiguchi, K. Shimada, H. Namatame and M. Taniguchi, *Nature* 426, 540 (2003).
- [3] A. R. Goni, A. Pinczuk, J. S. Weiner, J. M. Calleja, B. S. Dennis, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* 67, 3298 (1991).
- [4] Y. Tserkovnyak, B. I. Halperin, O. M. Auslaender and A. Yacoby, *cond-mat/0312159* (2003).
- [5] A. Yu. Kuznetsov, R. Deblock, M. Kociak, B. Reulet, H. Bouchiat, I. L. Khodos, Yu. B. Gorbatov, V. T. Volkov, C. Joumet and M. Burghard, *Science* 284, 1508 (1999); A. Kuznetsov, M. Kociak, M. Ferrier, R. Deblock, S. Gueron, B. Reulet, I. Khodos, O. Stephan, and H. Bouchiat, *Phys. Rev. B* 68, 214521 (2003).
- [6] A. F. Morpurgo, J. Kong, C. M. Marcus and H. Dai, *Science* 286, 263 (1999).
- [7] R. Fazio, F. W. J. Hekking and A. A. Odintsov, *Phys. Rev. Lett.* 74, 1843 (1995).
- [8] D. L. Maslov, M. Stone, P. M. Goldbart and D. Loss, *Phys. Rev. B* 53, 1548 (1996).
- [9] Y. Takane, *J. Phys. Soc. Jpn.* 66, 537 (1997).
- [10] I. Alek, J.-S. Caux and A. M. Zagoskin, *Phys. Rev. B* 62, 1433 (2000); J.-S. Caux, H. Saleur and F. Siano, *Phys. Rev. Lett.* 88, 106402 (2002).
- [11] M. P. A. Fisher, *Phys. Rev. B* 49, 14550 (1994); H. W. Lee, H. C. Lee, H. Yi and H.-Y. Choi, *Phys. Rev. Lett.* 90, 247001 (2003).
- [12] C. Winkelholz, R. Fazio, F. W. J. Hekking and G. Schon, *Phys. Rev. Lett.* 77, 3200 (2003).
- [13] L. N. Buluevskii, V. V. Kuzii and A. A. Sobyanin, *Sov. Phys. JETP Lett.* 25, 290 (1977); A. I. Buzdin, L. N. Buluevskii and S. V. Panyukov, *JETP Lett.* 35, 179 (1982).
- [14] V. V. Ryzanov et al., *Phys. Rev. Lett.* 86, 2427 (2001).
- [15] T. Kontos, M. Aprili, J. Lesueur and X. G. G. rison, *Phys. Rev. Lett.* 86, 304 (2001); W. Guichard, M. Aprili, O. Bourgeois, T. Kontos, J. Lesueur and P. Gandit, *Phys. Rev. Lett.* 90, 167001 (2003).
- [16] T. Kimura, K. Kuroki and H. Aoki, *Phys. Rev. B* 53, 9572 (1996).
- [17] R. L. Stratonovich, *Sov. Phys. Dokl.* 2, 416 (1958); J. Hubbard, *Phys. Rev. Lett.* 3, 77 (1959).
- [18] V. M. Yakovenko and H.-S. Goan, preprint.
- [19] G. Eilenberger, *Z. Phys.* 214, 195 (1968).
- [20] U. Günsenheim and A. D. Zaikin, *Phys. Rev. B* 50, 6317 (1994).
- [21] A. V. Zaitsev, *Sov. Phys. JETP* 59, 1015 (1984) [*Zh. Eksp. Teor. Fiz.* 86, 1742 (1984)].
- [22] P. Kopietz, *Bosonization of interacting fermions in arbitrary dimensions* (Springer Verlag, Berlin, 1997).
- [23] In Ref. 8, such a renormalization does not appear for perfectly transmitting regime. They consider a case that LL has strictly finite length L ; then momentum of fluctuations takes the discrete value $n\pi/L$. Our method reproduces the result of Ref. 8 in such situation.
- [24] C. L. Kane and M. P. A. Fisher, *Phys. Rev. B* 46, R7268 (1992); 46, 15233 (1992).
- [25] J. Solym, *Adv. Phys.* 28, 201 (1979).
- [26] *Crystal and Solid State Physics*, edited by O. Madelung et al., Landolt-Bornstein, New Series, Group III, vol. 16, Pt. a (Springer, Berlin, 1982).
- [27] H. Aoyama, T. Harano, H. Kikuchi, M. Sato and S. Wada, *Phys. Rev. Lett.* 79, 4052 (1997).
- [28] Here, the interference between decay paths to $\pi = 0/2$ is neglected, which can be important in some restricted case; N. Hatakenaka, *Phys. Rev. Lett.* 81, 3753 (1998).
- [29] J. Rollbühler and H. Grabert, *Phys. Rev. Lett.* 91, 166402 (2003).
- [30] Y. Y. Proskuryakov, A. K. Savchenko, S. S. Safonov, M. Pepper, M. Y. Simmons and D. A. Ritchie, *Phys. Rev. Lett.* 89, 076406 (2002).
- [31] R. de Picciotto, H. L. Stormer, A. Yacoby, L. N. Pfeiffer, K. W. Baldwin and K. W. West, *Phys. Rev. Lett.* 85, 1730 (2000).
- [32] T. A. Kazaki, H. Takayanagi, J. Nitta and T. Enoki, *Appl. Phys. Lett.* 68, 418 (1996).