

## Scaling approach to glassy stationary states of spin-glasses under chaos effects

Hajime YOSHINO<sup>1</sup>, and Petra E. JÖNSSON<sup>2</sup>,

<sup>1</sup>*Department of Earth and Space Science, Faculty of Science, Osaka University,  
Toyonaka, 560-0043 Osaka, Japan*

<sup>2</sup>*ISSP, University of Tokyo, Kashiwa-no-ha 5-1-5, Kashiwa, Chiba 277-8581, Japan*

Dynamics of spin-glasses subjected to slow continuous changes of working environment such as slow changes of temperature or interaction bonds are studied based on scaling arguments and numerical simulations of continuous bond changes. Such perturbations lead to a glassy stationary state where the *age* or domain size of the system is pinned to macroscopic but finite values due to competition between relaxation and chaos effects (rejuvenation). Fluctuation-dissipation relation is also pinned to that of a finite age. The scenario explains the anomalously weak cooling rate dependence of spin glasses.

**Introduction-** Dynamics of glassy systems have been actively studied over the past several decades. One general interest has been the aging effect,<sup>1)</sup> i.e. a glassy system ages extremely slowly in a fixed working environment specified for example by the temperature of the heatbath. Recently responses of such glassy systems to various kinds of perturbations have attracted much attention including restart of aging or rejuvenation of spin-glasses after sudden changes of the temperature of the heatbath,<sup>2),3)</sup> responses of soft glassy systems to shear.<sup>4)</sup>

In the present work we study relaxation of spin-glasses subjected to slow changes of the working environment, namely slow changes of temperature or interaction bonds. This study is motivated by the recent experimental observations that spin-glasses exhibit surprisingly weak dependence on the cooling rates.<sup>2),3)</sup> We propose a scaling ansatz for the effective age of the system under continuous changes of the working environment assuming that the competition between aging and rejuvenation due to the chaos effects<sup>5)</sup> leads to a stationary state. We performed Monte Carlo (MC) simulations of a 4-dimensional Edwards-Anderson (EA) model subjected to continuous changes of bonds and found good agreement with the scenario.

**Model-** Specifically we consider the Edwards-Anderson (EA) spin-glass model with Ising spins  $S_i$  ( $i = 1 \dots N$ ) on a  $d$ -dimensional (hyper)-cubic lattice with  $N = L^d$  lattice points. The Hamiltonian is given by  $H = - \sum_{\langle i,j \rangle} J_{ij} S_i S_j - h \sum_i S_i$ . We use the  $\pm J$  model where the interaction bonds  $J_{ij}$  between nearest-neighbours  $\langle i, j \rangle$  take random  $\pm J$  values. The temperature is measured as  $k_B T/J$  with the Boltzmann's constant set as  $k_B = 1$ . The magnetic field  $h$  is measured as  $h/J$ . The relaxational dynamics is modeled by the usual heatbath single spin flip MC algorithm.

A continuous bond changes is realized by flipping the sign of a fraction  $p \ll 1$  of the bonds chosen randomly in 1 Monte Carlo step (MCS). This amounts to flip a fraction  $p_{\text{eff}}(t) = [1 - (1 - 2p)^t]/2$  of the original set of bonds  $J_{ij}(0)$  to create the set of bonds  $J_{ij}(t)$  at  $t$  (MCS). For  $pt \ll 1$   $p_{\text{eff}}(t) = pt$  and  $p_{\text{eff}}(t) \rightarrow 1/2$  for  $pt \gg 1$ .

A continuous temperature changes is specified by a cooling or heating rate  $v_T =$

$\delta T/\delta t$  where  $\delta T$  is the change of the temperature within an interval of time  $\delta t$ . Thus the resultant change of the temperature after a time  $t$  is  $\Delta T(t) = v_T t$ . Unfortunately the temperature-chaos effect is much weaker than the bond chaos effect<sup>6)</sup> in the present model and we limit ourselves to the bond perturbations in the simulations.

**Scaling arguments-** Let us consider a spin-glass whose spins are evolving in a fixed working environment  $W = (\mathcal{J}, T)$  specified by a set of bonds  $\mathcal{J} = \{J_{ij}\}$  and the heatbath temperature  $T$ . If the initial spin configuration is far from the equilibrium state associated with  $W$ , the system is equilibrated only up to a certain length scale  $L_T(t)$  after a time  $t$ . The domain size  $L_T(t)$  grows slowly with time  $t$  by thermally activated droplets excitations as  $L_T(t) = L_0[(T/J) \log(t/\tau_c(T))]^{1/\psi}$  where  $L_0$  is a unit of length scale,  $\tau_c(T)$  is the attempt time of the activated dynamics and  $\psi > 0$  is the exponent of the free-energy barriers (See<sup>7)</sup> for the details).

Now we perturb the system by slowly changing  $W$  either by continuous temperature changes characterized by  $\Delta T(t) = v_T t$  or continuous bond changes characterized by  $\Delta J(t) = J\sqrt{p_{\text{eff}}(t)}$ .<sup>6)</sup> Within an interval of time  $\tau$ , the temporal working environment  $W(t)$  changes from  $W(0)$  to  $W(\tau)$ . Naturally the overlap length<sup>5)</sup> between the target equilibrium states associated with the two environments can be introduced as  $L_{\Delta J(\tau)} \sim L_0(\Delta J(\tau)/J)^{-1/\zeta}$  and  $L_{\Delta T(\tau)} \sim L_0(\Delta T(\tau)/T_g)^{-1/\zeta}$ . The target equilibrium states are completely different at length scales much larger than the overlap lengths by the chaos effects.<sup>5)</sup> During the continuous perturbations, the relaxational process which tends to grow the domains of  $W(t)$  and rejuvenation due to the chaos effects must be competing with each other. Naturally we expect that the system attains a certain stationary state characterized by a certain fixed size of the domains  $L_T(\tau^*)$  with respect to the temporal working environment  $W(t)$ . In other words the *age* of the system is pinned to a certain value  $\tau^*$ . As proposed in Ref,<sup>3),8)</sup> such a length scale should play the role of a sort of renormalized overlap length which parameterize the heating/cooling rate effects in temperature-cycling experiments. Simplest possible scaling form for the stationary domain size is

$$L_T(\tau^*) \sim L_0(\Delta J(\tau^*)/J)^{-1/\zeta} \quad L_T(\tau^*) \sim L_0(\Delta T(\tau^*)/T_g)^{-1/\zeta} \quad (1)$$

It can be seen that by choosing slower perturbations  $p \rightarrow 0$  or  $v_T \rightarrow 0$ , the pinned age increases  $\tau^* \rightarrow \infty$  and the stationary domain size increases  $L_T(\tau^*) \rightarrow \infty$ .

It can be also seen that in the limit of slow perturbations  $p \rightarrow 0$  or  $v_T \rightarrow 0$  actual change of the working environment becomes infinitesimal  $\Delta J(\tau^*) \rightarrow 0$  or  $\Delta T(\tau^*) \rightarrow 0$ . The later feature is quite unusual compared with analogues of more conventional cases such as systems under shear with shear rate  $\dot{\gamma}$ . There one typically finds that relaxation times scales as  $\tau^* \propto \dot{\gamma}^{-\nu}$  with  $\nu \leq 1$  so that actual deformation of the system  $\gamma \propto \dot{\gamma}^{1-\nu}$  within the relaxation time  $\tau^*$  can *diverge* as  $\dot{\gamma} \rightarrow 0$ .

During a continuous change of temperature the growth law  $L_T(t)$  would also change. However it is asymptotically irrelevant in the above scenario since  $L_{T+\Delta T}(t)/L_T(t) = (T + \Delta T/T)^{1/\psi} = O(1)$ . Thus for a given cooling/heating rate  $v_T$  the relaxation time  $\tau^*$  is defined at every local temperature  $T$ . The effective age  $\tau^*$  of a system cooled down to a certain target temperature  $T_m$  will be determined only by local cooling rate in the vicinity of  $T_m$  and the cooling rates used at higher temperatures are irrelevant. This feature observed experimentally in spin-glasses<sup>2)</sup> would hardly

be present in systems with activated dynamics but without the chaos effect.




Fig. 1. Spin autocorrelation function  $C(\tau + t_w, t_w) = (1/N) \sum_{i=1}^N \overline{S_i(\tau + t_w) S_i(t_w)}$  under continuous bond perturbation of strength  $p = 7.8125 \times 10^{-6}$  at  $T/J = 1.2$ . Here the averages over different realizations of thermal noises and sequences of bonds are denoted as  $\langle \dots \rangle$  and  $\overline{\dots}$  respectively. The lines in the left figure are reference data of standard isothermal aging.<sup>7)</sup> The system exhibits interrupted aging: waiting time  $t_w$  dependence saturates. The tail parts of the autocorrelation functions are fitted to a simple exponential form  $A \exp(-t/\tau^*(p))$  by which the characteristic relaxation time  $\tau^*(p)$  is extracted. In this example  $\tau^* = 6013$  (MCS) is obtained.




Fig. 2. The correlation  $C_{\text{bond}}(\tau^*)$  between the set of bonds at two times separated by the relaxation time  $\tau^*(p)$  is shown in the left figure. It is defined as  $C_{\text{bond}}(t) \equiv \overline{J_{ij}(0) J_{ij}(t)} / J^2 = \exp(-t/\tau_{\text{bond}}(p))$  with the decorrelation time of the bonds given as  $\tau_{\text{bond}}(p) = -1/\log(1-2p) \simeq 1/p$ . As shown in the inset the relaxation time of the spins  $\tau^*(p)$  grows with decreasing  $p$  but less rapidly than the decorrelation time of the bonds  $\tau_{\text{bond}}(p)$ . In the right figure the scaling ansatz Eq. 1 is tested. The open symbols are the reference data of  $L_T(\tau)$  (from Ref<sup>7)</sup>). The filled symbols are the overlap length  $a(\Delta J(\tau)/J)^{-1/\zeta}$  with  $\Delta J(t) = J\sqrt{p_{\text{eff}}(t)}$  at the relaxation time  $t = \tau^*(p)$  with  $p = 0.002 \times 2^{-n}$  with  $n = 0, \dots, 7$  (from left to right) obtained at  $T/J = 0.8, 1.2$  and  $1.6$ . Here  $\zeta = 0.9$  as obtained in Ref.<sup>3)</sup> (see also<sup>6)</sup>) and chosen the prefactor  $a = 0.22$ .

**Numerical analysis-** We performed MC simulations of the dynamics of the 4-dimensional EA model ( $T_g \simeq 2.0J$ ) under continuous bond changes starting from random initial configurations. In Fig. 1 we show the data of the spin autocorrelation function  $C(\tau + t_w, t_w)$ . Apparently the system becomes stationary within a finite

time scale  $\tau^*$  due to the continuous bond changes. The relaxation time  $\tau^*$  extracted from the data are examined as shown in Fig. 2. It appears that the actual change of the bonds within the relaxation time  $\tau^*$  of the bonds becomes negligible as the intensity of the driving  $p$  becomes smaller. As shown in the right figure of Fig. 2, the relaxation time  $\tau^*$  follows the expected scaling Eq. (1).

Finally let us examine the fluctuation dissipation relation (FDR) as proposed by the dynamical mean field theory (DMFT).<sup>9)</sup> As shown in Fig. 3 the fluctuation dissipation theorem is satisfied at short time scales and violated at larger time scales in a non-trivial manner as predicted by the DMFT. Thus the stationary state can be considered as a glassy state. The result is very similar to that of the same model during isothermal aging<sup>7)</sup> and driven dynamics induced by asymmetric coupling.<sup>10)</sup> It must be noted however that the FDR systematically depends on  $p$  as those of isothermal aging systematically depends on  $t_w$  so that the FDR in the asymptotic limit  $p \rightarrow 0$  may be different from the prediction of the DMFT.<sup>7)</sup>

**Discussion-** To conclude we studied glassy stationary dynamics of spin-glasses under continuous temperature or bond changes. Let us consider a typical cooling rate  $v_T = O(10^{-1})$ (K/sec) used in “quench” experiments, which is actually as slow as  $v_T = 10^{-15}$  J/MCS since  $T_g = O(10)$  K and the microscopic time scale is  $\tau_0 = 10^{-13}$  (sec). The scaling ansatz Eq. (1) implies it yields  $\tau^*$  of only order 1 – 10 (sec) at a target temperature  $T_m = 0.7T_g$  in the case of a canonical spin-glass AgMn sample,<sup>3)</sup> which allows one to observe “clean” aging at  $T_m$ . Indeed the effective age of the system after such slow cooling is found to be only  $O(10)$  (sec) experimentally.<sup>3)</sup>




Fig. 3. The ZFC susceptibility  $\chi_{\text{ZFC}}(\tau + t_w, t_w)$  vs. the spin autocorrelation function  $C(\tau + t_w, t_w)$  with  $t_w \gg \tau^*$ . In this example  $p = 2.5 \times 10^{-4}$ .  $\chi_{\text{ZFC}}$  is measured as usual<sup>7)</sup> with  $h/J = 0.1$  but under continuous bond changes. The lines represent FDT  $\chi_{\text{ZFC}} = (1-C)/T$  associated with the heatbath temperatures  $T$ .

## References

- 1) J.-P. Bouchaud, L. F. Cugliandolo, J. Kurchan, and M. Mézard, in *Spin Glasses and Random Fields*, edited by A. P. Young, pages 161–223, World Scientific, Singapore, 1997.
- 2) K. Jonason, E. Vincent, J. Hammann, J.-P. Bouchaud, and P. Nordblad, Phys. Rev. Lett. **81**, 3243 (1998).
- 3) P. E. Jönsson, R. Mathieu, H. Yoshino, P. Nordblad, H. A. Katori, and A. Ito, cond-mat/0307640 (2003).
- 4) P. Sollich, F. Lequeux, P. Hebraud, and M. E. Cates, Phys. Rev. Lett. **78**, 2020 (1997).
- 5) A. J. Bray and M. A. Moore, Phys. Rev. Lett. **58**, 57 (1987); D. S. Fisher and D. A. Huse, Phys. Rev. B **38**, 386 (1988);
- 6) M. Ney-Nifle, Phys. Rev. B **57**, 492 (1998); M. Sasaki et al, in this volume.
- 7) H. Yoshino, K. Hukushima, and H. Takayama, Phys. Rev. B **66**, 064431 (2002).
- 8) H. Yoshino, J. Phys. A **36**, 10819 (2003).
- 9) L. F. Cugliandolo and J. Kurchan, Phys. Rev. Lett. **71**, 173 (1993).
- 10) L. Berthier, J. Phys. A **36**, 10667 (2003).