

Do quantum dots allow one access to pseudogap Kondo physics?

John Hopkinson¹, Karyn Le Hur, and Emilie Dupont

Département de Physique, Université de Sherbrooke, Sherbrooke, Québec, Canada

Abstract

For the last decade[1], tunable quantum dot systems have allowed the investigation of Kondo physics wherein the quenching of a single spin on an artificial atom affects the conductance. The pseudogap Kondo model (pKM) featuring a density of states $\rho(\epsilon) = C|\epsilon|^\alpha$, introduced by Wirtho and Fradkin [2] in 1990 was predicted to exhibit Kondo-like physics above a critical value of the Kondo coupling, J_c , which several groups have shown by numerical renormalization group (RG) is finite for $\alpha < \frac{1}{2}$. Gonzalez-Buxton et al [3] showed that the strong coupling limit of the particle-hole symmetric model leads to a non-trivial $\frac{(1-\alpha)}{2}$ phase shift at low temperatures indicating incomplete screening of the local moment, while away from particle-hole symmetry (p-hs) one generically flows towards a ground state with $\frac{\pi}{2}$. We examine the implications of this model for quantum dots whose leads are Fermi-liquid-like, yet possess a tunneling density of states (TDOS) which is suppressed at the Fermi energy as a power law.

Key words:

pseudogap Kondo model, quantum dots, quantum critical point

Many of the unusual features arising in strongly correlated electron systems have been attributed to the proximity to quantum critical points. In particular, heavy fermion systems are thought to feature a competition between RKKY and Kondo interactions, within the context of a Kondo lattice model, leading to (an increasingly heavy) metallic antiferromagnetic phase which is suppressed as a function of pressure, chemical substitution or magnetic field to produce a cone of non-Fermi liquid physics preceding a (decreasingly heavy) metallic state. Unfortunately, despite decades of intensive research on the Kondo lattice model, the detailed nature of its solution remains a subject of some debate. A simple model exhibiting a quantum phase transition between a state with local moments and spin-screened state, the pKM, was introduced in Ref. 2 and studied extensively by subsequent groups [3-5]. This paper explores the possibility that this model could be experimentally realizable in a quantum dot system.

Resonant level limit We consider the small dot, weak tunneling limit in which the dot is a collection of discrete levels of average spacing E_d , an order of magnitude smaller than the charging energy, E_c ($\propto \frac{e^2}{2C}$), of the dot (e : electron charge; C : capacitance of dot-lead system). The energy to add or subtract an electron from the dot can be estimated as $E_c = E_c(Q_G, n+1) - E_c(Q_G, n) = (n + \frac{1}{2} - C_G V_G) \frac{e^2}{C}$, where n is the initial (final) number of electrons on the dot for + (-), V_G is an external gate voltage coupled capacitively (C_G) to the dot. A resonant condition periodic in the number of electrons on the dot corresponds to $E_c^{(+)} = 0$, $E_c^{(+)} = \frac{e^2}{C}$. For temperatures satisfying $E_d < T < E_c$, the physics of the dot is dominated by Coulomb blockade. For temperatures below E_d , one can write the current through the dot in the single level Landauer-Buttiker (elastic scattering) form,

$$I = \frac{2e}{h} \sum_D \frac{4(f_L(\epsilon_D) - f_R(\epsilon_D))}{(\epsilon_D)^2 + (\Gamma_L + \Gamma_R)^2} \frac{1}{2} \int_{-\infty}^{\infty} d\epsilon \quad (1)$$

$f_{L(R)}(\epsilon)$ is the Fermi function of the left (right) lead,

Corresponding Author: Université de Sherbrooke, QC, CA
Phone: (819) 821-8000 ext. 3043 Fax: (819) 821-8046

¹ This work was supported by NSERC, FQRNT and CIAR.

Following Glazman and Raikh [7], the effects of correlation in the leads ($r=0$ for free electrons) is to decrease the effective conductance relative to unitarity at the Kondo limit as $G = G_0 \sin^2(\frac{r}{2})$, a result one might have anticipated from Kane and Fisher's [8] repulsive interactions in Luttinger liquids (there $G_0 \propto g G_0$, g : Luttinger parameter). In the Kondo limit, Ref. 3 found that at p-hs, the problem shows to a strong coupling fixed point corresponding to the $U=0$ limit of the pAM provided r exceeds a critical value which increases monotonically with r to diverge at $\frac{1}{2}$. The phase shift of this model at the Fermi energy is $\delta = \frac{\pi}{2} (1 - \text{sgn}(r))$ [3,4]. However, this strong coupling fixed point is unstable with respect to the potential scattering term V . At the symmetric point, $V=0$, one should [5] enter the particle-hole symmetric fixed point below the Kondo scale $T_K \propto E_F (\frac{J}{J_0})^{\frac{1}{1-r}}$. If $r \neq 0$, for $0 < r < 0.375$, the initial flow is towards the p-hs fixed point. Below a temperature scale $T_0 = \frac{J_0}{J} T_K$ the problem flows to an asymmetric strong coupling fixed point [5] whose phase shift [3] about the Fermi energy is $\delta = \frac{\pi}{2}$. While such a phase shift would be opaque to conductance measurements,

