

# Comment on "Kondo resonances and Fano antiresonances in transport through quantum dots"

J. Bonča, A. Ramšak, and T. Rejec

*J. Stefan Institute, Jamova 39, 1000 Ljubljana, Slovenia*

*Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, 1000 Ljubljana, Slovenia*

(Dated: 19 July 2004)

We show that the numerical method [M. E. Torio, K. Hallberg, A. H. Ceccatto, and C. R. Proetto, Phys. Rev. B **65**, 085302 (2002)] does not reproduce correctly the Kondo physics in quantum dot systems with interaction.

In a recent paper, M. E. Torio, K. Hallberg, A. H. Ceccatto, and C. R. Proetto (THCP) presented a numerical calculation of the conductance of an interacting system consisting of a quantum dot coupled to the leads.<sup>1</sup> Their results exhibit a fair qualitative agreement with the corresponding experiments by J. Göres *et al.*<sup>2</sup> However, a closer quantitative inspection of the results concerning the Kondo regime clearly shows that their method fails to reproduce adequately the main features of the conductance in this regime.

Numerical method of THCP consists of exact diagonalization of a small cluster containing the region with interaction, coupling the cluster to external reservoirs (leads), and finally self-consistent calculation of the corresponding Dyson equation. As pointed out in THCP, the method should be reliable as long as the corresponding Kondo cloud does not exceed the size of the cluster. In order to meet these criteria, THCP chose  $t'/t = 1/\sqrt{2}$  and estimated (without giving arguments) the corresponding Kondo correlation length of about 10 lattice sites for  $U/\Delta = 12$  with  $\Delta = 2t'^2/t$ . Using this set of parameters for the Anderson model<sup>3</sup> they calculated the conductance of substitutional quantum dot configuration. For the symmetric situation,  $\epsilon_0 = -U/2$ , their conductance reaches the unitary limit,  $G = G_0 = 2e^2/h$ . They further claimed that this is a nontrivial check of the method which proves that their finite-system approach produces a Kondo peak with the exact spectral weight at the Fermi energy.<sup>1</sup>

In this Comment we show that although the conductance of THCP reaches the unitary limit, the method does not reproduce correctly the Kondo peak. In Fig. 1(a) the conductance of THCP is shown compared to the result obtained with the two-point sine conductance formula from Ref. 4 using the variational wave function of Gunnarsson and Schönhammer<sup>5</sup> together with the constrained path quantum Monte Carlo method (CPMC).<sup>6,7</sup> In the empty orbital regime,  $|\epsilon_0 + U/2| \gtrsim U/2$ , all results match perfectly. However, for  $|\epsilon_0 + U/2| \lesssim U/2$ , except in the symmetric point, the result of THCP drastically fails to reproduce the main fingerprint of Kondo physics, i.e., the plateau in the conductance in the Kondo regime. The reason is the low energy resolution which in turn is a fingerprint of finite-size effects of a small clusters used by the method of THCP. It should be pointed out that

even relatively large  $t'$  cannot remedy this deficiency of the method, because the problem falls into the unusually *narrow* bandwidth regime  $2t \ll U$ , and  $\epsilon_0 \ll -2t$ ,  $\epsilon_0 + U \gg 2t$ , where the scattering processes which involve electrons or holes in the band edges cannot cause real charge fluctuations of the impurity,<sup>8</sup> resulting in the reduction of the Kondo temperature. On the one hand, narrow bandwidth should improve the energy resolution of the method,<sup>1</sup> on the other hand, however, the Kondo temperature  $T_K$  in this limit is reduced,<sup>9</sup>

$$T_K \sim 2t \left( \frac{\Delta U}{|\epsilon_0| |\epsilon_0 + U|} \right)^{1/2} e^{-\pi|\epsilon_0||\epsilon_0 + U|/(2\Delta U)}. \quad (1)$$

Smaller  $t$  (or larger  $t'$ ) thus cannot bring the method of THCP into the regime of reliability unless the number of sites in the cluster is significantly increased or the strength of the interaction,  $U/\Delta$ , is reduced.

In a more common limit of a wide bandwidth,  $2t \gg U$ ,<sup>10</sup> the conductance can be calculated exactly using the Friedel sum rule and the Bethe ansatz result.<sup>11</sup> In Fig. 1(b) the exact conductance for wide bandwidth is presented, together with the result of the two-point sine formula, using variational wavefunction and the CPMC method. The results agree for all regimes. Small discrepancies can be attributed to the finite bandwidth of both numerical methods ( $t = 1.04U$ ) and to the variational nature of the wave functions used. We would like to stress that due to the lower Kondo temperature in the narrow bandwidth regime, Fig. 1(a), the Kondo plateau is obviously more pronounced compared to the wide bandwidth limit, Fig. 1(b).

In Fig. 1(b) the conductance using an elementary Hartree-Fock calculation is also presented. It is well known that this approximation cannot adequately reproduce the Kondo phenomenon. Nevertheless, the unitary limit in the symmetric point,  $\epsilon_0 + U/2 = 0$ , is in this approach as well reproduced. It is a consequence of the particle-hole symmetry of the Anderson model combined with the Friedel sum rule. The existence of the unitary limit can in general serve only as a test of internal numerical consistency of a particular numerical method and is not a direct evidence of a correct description of the Kondo physics as argued by THCP.




Figure 1: (a) In  $2t \ll U$  regime the conductance of THCP (diamonds) compared to results obtained from the two-point sine conductance formula,<sup>4</sup> using variational wave function of Gunnarsson and Schönhammer (full line)<sup>5</sup> and CPMC method (bullets).<sup>7</sup> (b) Conductance for  $2t \gg U$  obtained with exact Bethe ansatz (dashed-dotted)<sup>11</sup> and Hartree-Fock (dashed). In numerical calculation  $t = 1.04U$  was used for the case of variational wave function (full line)<sup>4,5</sup> and CPMC (bullets).<sup>7</sup>

In summary, the method used by THCP seems to be promising for the calculation of conductance of systems where the interaction is very weak. However, in the limit of strong electron-electron interaction it fails to reproduce many-body effects as is, e.g., the Kondo physics in transport through a quantum dot.

Authors acknowledge the support of the Ministry of Education, Science and Sport of Slovenia under grant P1-0044.

- 
- <sup>1</sup> M. E. Torio, K. Hallberg, A. H. Ceccatto, and C. R. Proetto, Phys. Rev. B **65**, 085302 (2002).  
<sup>2</sup> J. Göres, D. Goldhaber-Gordon, S. Heemeyer, M. A. Kastner, Hadas Shtrikman, D. Mahalu, and U. Meirav, Phys. Rev. B **62**, 2188 (2000).  
<sup>3</sup> All model parameters throughout this Comment are defined as in Ref. 1.  
<sup>4</sup> T. Rejec and A. Ramšak, Phys. Rev. B **68**, 033306 (2003).  
<sup>5</sup> O. Gunnarsson and K. Schönhammer, Phys. Rev. B **31**, 4815 (1985).  
<sup>6</sup> S. Zhang, J. Carlson, and J. E. Gubernatis, Phys. Rev. Lett. **74**, 3652 (1995); Phys. Rev. B **55**, 7464 (1997).  
<sup>7</sup> J. Bonča, A. Ramšak, and T. Rejec, unpublished.  
<sup>8</sup> J. H. Jefferson, J. Phys. C **10**, 3589 (1977).  
<sup>9</sup> A. C. Hewson, *The Kondo Problem to Heavy Fermions* (Cambridge University Press, Cambridge, 1993).  
<sup>10</sup> F. D. M. Haldane, Phys. Rev. Lett. **40**, 416 (1978).  
<sup>11</sup> P. B. Wiegman, Sov. Phys. JETP Lett. **31**, 392 (1980); N. Andrei, Phys. Rev. Lett. **45**, 379 (1980).