Site-Bond Representation and Self-Duality for Totalistic Probabilistic Cellular Automata in One Dimension

N. Konno[†], X. Ma

Department of Applied Mathematics, Yokohama National University, Hodogaya-ku, Yokohama, 240-8501, Japan

Abstract.

We study the one-dimensional two-state totalistic probabilistic cellular automata (TPCA) having an absorbing state with long-range interactions, which can be considered as a natural extension of the Domany-Kinzel model. We establish the conditions for existence of a site-bond representation and self-dual property. Moreover we present an expression of a set-to-set connectedness between two sets, a matrix expression for a condition of the self-duality, and a convergence theorem for the TPCA.

1. Introduction

Probabilistic versions of cellular automata can be considered as discrete-time Markov processes with parallel updating, which are useful in a large number of scientific areas [1]. As a special case, the Domany-Kinzel (DK) model introduced and studied by [2] is one of the simplest examples of the probabilistic cellular automata. The model is defined on a lattice, and the two different states for a site can be said as empty or occupied. The state of a given site depends on the number of the occupied states in its two nearest neighbors at the previous time step, where this property can be expressed with the totalistic rule [1, 3, 4]. In the DK model, the transition probability from an empty neighborhood to an occupied state is zero, so the empty configuration is an absorbing state. The totalistic probabilistic cellular automaton (TPCA) is a natural extension of the DK model, where we extend the number of neighbors to a finite integer $N \geq 2$. Remark that N = 2 case becomes the DK model. Concerning the DK model, various results are known (see our recent papers [5, 6, 7, 8] and references therein). As for N = 3case with two absorbing states, a rich phase diagram is reported by using mean-field approximations and numerical simulations (see Refs. [9, 10], for examples). However there are few rigorous results on the TPCA for a general $N \geq 3$. In this situation, the purpose of this paper is to give some rigorous results for a two-state N-neighbor TPCA with an absorbing state. More precisely, we reveal a relation between the site-bond representation and the self-duality for the TPCA. Furthermore we present an expression of a set-to-set connectedness between two sets, a matrix expression for a condition of the self-duality, and a convergence theorem for the TPCA. It is known that the self-duality is a very useful technique in the study of stochastic interacting models. Because problems in uncountable state space (typically configurations of $\{0,1\}^{\mathbb{Z}}$) can be reformulated as problems in countable state space (typically finite subsets of \mathbf{Z}). For some applications of the self-duality on discrete-time case (e.g., oriented bond percolation model) and continuous-time case (e.g., contact process), see Refs. [11, 12, 13, 14]. So our results here might be useful in obtaining rigorous results on the phase diagram and critical properties of such models.

We have organized the present paper as follows: Section 2 is devoted to the definition of our TPCA. In Section 3, we explain a site-bond representation and a self-duality for the TPCA. Moreover an expression of a set-to-set connectedness for the TPCA is given. Section 4 treats a matrix expression for the self-duality of the model. Section 5 contains a convergence theorem. Finally, concluding remarks and discussions are presented in Section 6.

2. Definition of the TPCA

First we give the definition of the DK model (N = 2 case). Let $\xi_n^A \subset \mathbf{Z}$ be the state of the process with parameters $(p_1, p_2) \in [0, 1]^2$ at time *n* starting from a set $A \subset 2\mathbf{Z}$. Its

evolution is described by

$$P(x \in \xi_{n+1}^{A} | \xi_{n}^{A}) = f(\xi_{n}^{A}(x-1) + \xi_{n}^{A}(x+1)),$$

and given ξ_n^A , the events $\{x \in \xi_{n+1}^A\}$ are independent, where P(B|C) is the conditional probability that B occurs given that C occurs, and

$$f(0) = 0, \quad f(1) = p_1, \quad f(2) = p_2.$$

If we write $\xi(x, n) = 1$ for $x \in \xi_n^A$ and $\xi(x, n) = 0$ otherwise, each realization of the process is identified with a configuration $\xi \in \{0, 1\}^S$ with $S = \{s = (x, n) \in \mathbb{Z} \times \mathbb{Z}_+ : x + n = \text{even }\}$, where $\mathbb{Z}_+ = \{0, 1, 2, \ldots\}$. The model with $(p_1, p_2) = (1, 0)$ becomes Wolfram's rule 90. For more detailed information, see Section 5 in [12].

From now on, we introduce a long-range TPCA. In order to clarify the definition, we consider two cases of N = even and N = odd respectively, where N is the number of the neighborhood.

(i) N = 2L (L = 1, 2, ...) case. The space of sites is denoted by $S_0 = \{s = (x, n) \in \mathbb{Z} \times \mathbb{Z}_+ : x + n = \text{even} \}$ and the space of bonds is

$$B_0 = \{((x, n+1), (x-2L+1, n)), \dots, ((x, n+1), (x-1, n)), ((x, n+1), (x+1, n)), \dots, ((x, n+1), (x+2L-1, n)) : (x, n+1) \in S_0\}.$$

For any initial set $A \subset 2\mathbf{Z}$, the process ξ_n^A satisfies

$$P(x \in \xi_{n+1}^{A} | \xi_{n}^{A}) = f(\sum_{k=-(L-1)}^{L} \xi_{n}^{A} (x - 2k + 1)),$$

and given ξ_n^A , the events $\{x \in \xi_{n+1}^A\}$ are independent, where

$$f(0) = p_0 = 0, f(1) = p_1, \dots, f(2L) = p_{2L},$$

with $p_1, p_2, \ldots, p_{2L} \in [0, 1]$. This process is considered on the space S_0 . The case L = 1 is equivalent to the DK model.

(ii) N = 2L + 1 (L = 1, 2, ...) case. The space of sites is denoted by

$$S_1 = \{ s = (x, n) \in \mathbf{Z} \times \mathbf{Z}_+ \},\$$

and the space of bonds is

$$B_1 = \{((x, n+1), (x-L, n)), \dots, ((x, n+1), (x-1, n)), ((x, n+1), (x, n)), ((x, n+1), (x+1, n)), \dots, ((x, n+1), (x+L, n)) : (x, n+1) \in S_1\}.$$

For any initial set $A \subset \mathbf{Z}$, the process ξ_n^A satisfies

$$P(x \in \xi_{n+1}^{A} | \xi_{n}^{A}) = f(\sum_{k=-L}^{L} \xi_{n}^{A} (x+k)),$$

and given ξ_n^A , the events $\{x \in \xi_{n+1}^A\}$ are independent, where

$$f(0) = p_0 = 0, f(1) = p_1, \dots, f(2L+1) = p_{2L+1},$$

with $p_1, p_2, \ldots, p_{2L+1} \in [0, 1]$. This process is considered on the space S_1 .

If $p_0 = 0 \le p_1 \le p_2 \le \ldots \le p_{N-1} \le p_N$, then the N-neighbor TPCA is said to be *attractive*. If not, it is said to be *non-attractive*. The attractiveness means that having more particles at one time implies there will be more particles at the next time(s).

In general, few rigorous results on the non-attractive model are known compared with the attractive model. Much more informations on the results for the non-attractive N = 2 case are shown for instance in [7]. So studying a general model including the non-attractive case, such as is the one in this paper, is very important.

3. Site-bond representation and self-duality

From now on we consider only $(p_1, p_2) \in D_*$ case, where the subset of parameter space D_* is defined in the following:

$$D_* = \{(p_1, p_2) : 0 < p_1 \le 1, \ 0 < p_2 \le 1 \text{ and } p_2 < 2p_1\}.$$

A reason for introducing the set is that for any point outside of D_* , it will be shown that there exists no site-bond representation for the TPCA (see Proposition 1 below). Suppose that $p_1, p_2 \in D_*$ are given. We put $\alpha = p_1^2/(2p_1 - p_2)$, $\beta = 2 - p_2/p_1$. Let $(S, B) = (S_0, B_0)$ or (S_1, B_1) and **R** denote the set of real numbers. If there exist $\alpha, \beta \in \mathbf{R}$ such that $p_n \in [0, 1]$ is given by

$$p_n = \alpha [1 - (1 - \beta)^n], \tag{1}$$

for n = 3, 4, ..., N, then we call "an *N*-neighbor TPCA with $\{p_n : 0 \le n \le N\}$ has a site-bond representation with α and β ". In fact, when $\alpha, \beta \in (0, 1]$, the TPCA can be considered as an oriented mixed site-bond percolation with a long range interaction in the following way. On the space (S, B) we define

$$X(S) = \{0,1\}^S, \quad X(B) = \{0,1\}^B, \quad X = X(S) \times X(B).$$

For given $\zeta = (\zeta_1, \zeta_2) \in X$, we say that $s = (y, n + k) \in S$ can be reached from $s' = (x, n) \in S$ and write $s' \to s$, if there exists a sequence $s_0, s_1, s_2, \ldots, s_k$ of members of S such that $s' = s_0, s = s_k$, and

$$\zeta_1(s_i) = 1, i = 0, 1, \dots, k; \ \zeta_2((s_i, s_{i+1})) = 1, i = 0, 1, \dots, k-1.$$

we also say that $G \subset S$ can be reached from $G' \subset S$ and write $G' \to G$, if there exist $s \in G$ and $s' \in G'$ such that $s' \to s$.

We introduce the signed measure m on X defined by

$$m(\Lambda) = \alpha^{k_1} (1 - \alpha)^{j_1} \beta^{k_2} (1 - \beta)^{j_2},$$

for any cylinder set

$$\Lambda = \{ (\zeta_1, \zeta_2) \in X \colon \zeta_1(s_i) = 1, i = 1, 2, \dots, k_1; \, \zeta_1(s'_i) = 0, i = 1, 2, \dots, j_1; \\ \zeta_2(b_i) = 1, i = 1, 2, \dots, k_2; \, \zeta_2(b'_i) = 0, i = 1, 2, \dots, j_2 \},$$

where $s_1, s_2, \ldots, s_k, s'_1, s'_2, \ldots, s'_j$ are distinct elements of S, and $b_1, b_2, \ldots, b_k, b'_1, b'_2, \ldots, b'_j$ are distinct elements of B and $\alpha = p_1^2/(2p_1 - p_2), \ \beta = 2 - p_2/p_1$. We should remark the next three facts. If $p_2 < 2p_1$ and $p_2 > 2p_1 - p_1^2$, then $\alpha > 1$ and $\beta \in (0, 1)$. If $p_2 \leq 2p_1 - p_1^2$ and $p_2 \geq p_1$, then $\alpha, \beta \in (0, 1]$. Moreover, if $p_2 < p_1$, then $\alpha \in (0, 1)$ and $\beta \in (1, 2]$. From the above observation, we see that the measure is not probability measure in the first and third cases, since $\alpha > 1$ and $1 - \beta < 0$ respectively. See Refs. [5] and [7] for more details on the site-bond representation.

It is noted that if an N-neighbor TPCA has a site-bond representation with α and β , then $\alpha = p_1^2/(2p_1 - p_2)$, $\beta = 2 - p_2/p_1$, since Eq. (1) gives

$$p_1 = \alpha \beta, \qquad p_2 = \alpha [1 - (1 - \beta)^2].$$

As we will see in the proof of Proposition 1, the following result can be obtained: an *N*-neighbor TPCA having a site-bond representation with α and β is attractive if and only if $p_1 \leq p_2$. In other words, an *N*-neighbor TPCA having a site-bond representation with α and β is non-attractive if and only if $p_1 > p_2$. Then we have

Proposition 1 We consider N-neighbor TPCA with $\{p_n : 0 \leq n \leq N\}$. Assume $(p_1, p_2) \in D_*$. Let $\alpha = p_1^2/(2p_1 - p_2)$ and $\beta = (2p_1 - p_2)/p_1$. (i) $p_1 > p_2$ (non-attractive case) and $p_n = \alpha [1 - (1 - \beta)^n] (0 \leq n \leq N)$, then $p_n \in [0, 1] (0 \leq n \leq N)$. (ii) $p_1 \leq p_2$ (attractive case), $p_n = \alpha [1 - (1 - \beta)^n] (0 \leq n \leq N)$, and $p_N \leq 1$, then $p_n \in [0, 1] (0 \leq n \leq N)$.

Proof. First we consider a relation among $p_0 = 0, p_1, \ldots, p_N$. By using $p_n = \alpha [1 - (1 - \beta)^n]$ for $n = 0, 1, \ldots, N$, we see

$$p_n - p_{n-1} = \alpha \beta (1 - \beta)^{n-1},$$
 (2)

$$p_n - p_{n-2} = \alpha (1 - \beta)^{n-2} \{ 1 - (1 - \beta)^2 \}.$$
(3)

Recall that

$$1 - \beta = (p_2 - p_1)/p_1, \qquad \alpha \beta = p_1.$$
 (4)

Now we consider $p_1 > p_2$ (non-attractive) case. In this case, Eq. (4) gives

 $-1 < 1 - \beta < 0, \qquad 0 < \alpha \beta \le 1, \qquad 0 < \alpha < 1.$ (5)

From Eqs. (2), (3), and (5), we see

$$p_0 = 0 < p_2 < p_4 < p_6 < \ldots < p_{2n} < p_{2n-1} < \ldots < p_5 < p_3 < p_1 \le 1.$$

Thus we obtain $p_n \in [0, 1]$ for any n = 0, 1, ..., N. Next we consider $p_1 \leq p_2$ (attractive) case. It is easily checked that $p_n \leq p_{n+1}$ for any n = 0, 1, ..., N - 1 by using Eq. (2) and $0 \leq 1 - \beta < 1$. So we have

$$p_0 = 0 < p_1 \le p_2 \le p_3 \le \ldots \le p_{N-1} \le p_N.$$

Therefore a necessary and sufficient condition for the site-bond representation in this case is $p_N = \alpha [1 - (1 - \beta)^N] \leq 1$. The proof of Proposition 1 is complete.

Remark that $p_n = \alpha [1 - (1 - \beta)^n] \to \alpha$ as $n \to \infty$, since $|1 - \beta| < 1$. Noting that $\alpha \le 1$ if and only if $p_2 \le 2p_1 - p_1^2$, we conclude that a sufficient (and not necessary) condition for a site-bond representation of an N-neighbor TPCA is $p_2 \le 2p_1 - p_1^2$. Furthermore

Eq. (2) implies

Corollary 2 If an *N*-neighbor TPCA with $\{p_n : 0 \le n \le N\}$ and $N \ge 3$ has the site-bond representation, then p_n can be expanded with p_1 and p_2 as follows:

$$p_n = \frac{(p_2 - p_1)^{n-1}}{p_1^{n-2}} + \frac{(p_2 - p_1)^{n-2}}{p_1^{n-3}} + \dots + \frac{(p_2 - p_1)^2}{p_1} + p_2,$$

for any n = 3, 4, ..., N.

Next we define a set-to-set connectedness for the TPCA from a set A to a set B by

$$\sigma(A,B) = \lim_{n \to \infty} P(\xi_n^A \cap B \neq \emptyset),$$

if the right-hand side exists. As in a similar argument of N = 2 case (the DK model) [5], we can easily extend the case to a general N-neighbor TPCA:

Proposition 3 Let $(p_1, p_2) \in D_*$. We assume that an *N*-neighbor TPCA with $\{p_n : 0 \le n \le N\}$ has a site-bond representation with α and β , where $\alpha = p_1^2/(2p_1 - p_2)$ and $\beta = (2p_1 - p_2)/p_1$. Then for any *A* with $|A| < \infty$, we have

$$\sigma(2\mathbf{Z}, A) = \sum_{D \subset A, D \neq \emptyset} \alpha^{|D|} (1 - \alpha)^{|A \setminus D|} \sigma(D, 2\mathbf{Z}),$$

when N = 2L, and

$$\sigma(\mathbf{Z}, A) = \sum_{D \subset A, D \neq \emptyset} \alpha^{|D|} (1 - \alpha)^{|A \setminus D|} \sigma(D, \mathbf{Z}),$$

when N = 2L + 1.

From now on, we consider a self-duality of the TPCA. An N-neighbor TPCA ξ_n is said to be *self-dual* with a *self-duality parameter* x if

$$E\left(x^{|\xi_n^A \cap B|}\right) = E\left(x^{|\xi_n^B \cap A|}\right) \qquad (n = 0, 1, \ldots)$$

holds for any $A, B \subset \mathbb{Z}$ with $|A| < \infty$ or $|B| < \infty$. The above equation is called a *self-duality equation*. Then we have

Theorem 4 Let $(p_1, p_2) \in D_*$ with $\alpha [1 - (1 - \beta)^N] \leq 1$. An *N*-neighbor TPCA with $\{p_n : 0 \leq n \leq N\}$ is self-dual with a self-duality parameter $(\alpha - 1)/\alpha$ if and only if this model has a site-bond representation with α and β , where $\alpha = p_1^2/(2p_1 - p_2)$ and $\beta = (2p_1 - p_2)/p_1$.

Proof. Theorem 1 in [15] implies that an N-neighbor TPCA with transition probabilities $\{p_n : 0 \le n \le N\}$ is self-dual with a self-duality parameter x is equivalent to

$$(p_i x + q_i)^j = (p_j x + q_j)^i, (6)$$

for $0 \le i, j \le N$, where $q_i = 1 - p_i$. It can be confirmed that $x = (\alpha - 1)/\alpha$ with $\alpha = p_1^2/(2p_1 - p_2)$ satisfies Eq. (6). Moreover we see that Eq. (6) if and only if

$$(p_1 x + q_1)^n = p_n x + q_n, (7)$$

for $0 \le n \le N$. In fact if we take i = 1 and j = n, Eq. (6) becomes Eq. (7). Conversely we have

$$(p_i x + q_i)^j = (p_1 x + q_1)^{ij} = (p_j x + q_j)^i$$

Therefore we put y = x - 1. Then Eq. (7) becomes $p_n y + 1 = (p_1 y + 1)^n$. So we see that

$$p_{n} = \sum_{r=1}^{n} {n \choose r} y^{r-1} p_{1}^{r}$$

$$= \sum_{r=1}^{n} {n \choose r} \left(\frac{p_{2} - 2p_{1}}{p_{1}^{2}}\right)^{r-1} p_{1}^{r}$$

$$= \left(\frac{p_{2} - 2p_{1}}{p_{1}^{2}}\right)^{-1} \sum_{r=1}^{n} {n \choose r} \left(\frac{p_{2} - 2p_{1}}{p_{1}}\right)^{r}$$

$$= \alpha \sum_{r=0}^{n} {n \choose r} \left(1 - \frac{p_{2} - 2p_{1}}{p_{1}}\right)^{r}$$

$$= \alpha [1 - (1 + (p_{2} - 2p_{1})/p_{1})^{n}]$$

$$= \alpha [1 - (1 - \beta)^{n}].$$

So the proof of Theorem 4 is complete.

From this theorem it can be seen that two conditions for the self-duality and for the site-bond representation are equivalent under an assumption that $(p_1, p_2) \in D_*$ with $p_N = \alpha [1 - (1 - \beta)^N] \le 1.$

4. Matrix expression

In this section we consider a criterion for the self-duality based on a matrix expression. Let

$$X(x) = \left[\begin{array}{cc} 1 & 1 \\ 1 & x \end{array} \right],$$

and

$$X_N(x) = X(x)^{\otimes N},$$

where \otimes indicates the tensor product. For example, in the case of N = 2, we have

$$X_2(x) = X(x) \otimes X(x) = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & x & 1 & x \\ 1 & 1 & x & x \\ 1 & x & x & x^2 \end{bmatrix}$$

For fixed $i, j \in \{1, 2, ..., 2^N\}$, the values of $i_1, i_2, ..., i_N$ and $j_1, j_2, ..., j_N \in \{0, 1\}$ are defined by the binary expansion of i and j in the following:

$$i - 1 = i_1 2^{N-1} + i_2 2^{N-2} + \dots + i_{N-1} 2^1 + i_N 2^0,$$

$$j - 1 = j_1 2^{N-1} + j_2 2^{N-2} + \dots + j_{N-1} 2^1 + j_N 2^0,$$

and $I = i_1 + i_2 + \cdots + i_N$. Moreover we introduce $2^N \times 2^N$ matrix P_N whose (i, j) element is defined by

$$p_I^{(j_1)} p_I^{(j_2)} \cdots p_I^{(j_N)},$$

where $p_I^{(1)} = p_I$, $p_I^{(0)} = 1 - p_I = q_I$. For N = 2 case, the above definition gives

$$P_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ q_1^2 & p_1 q_1 & p_1 q_1 & p_1^2 \\ q_1^2 & p_1 q_1 & p_1 q_1 & p_1^2 \\ q_2^2 & p_2 q_2 & p_2 q_2 & p_2^2 \end{bmatrix},$$

since $p_0 = 0$, $q_0 = 1$. Then we have the next result:

Theorem 5 A necessary and sufficient condition of the self-duality for an N-neighbor TPCA with $\{p_n : 0 \le n \le N\}$ with a self-duality parameter x is that x satisfies

$$P_N X_N(x) = {}^t (P_N X_N(x)), \tag{8}$$

where t means the transposed operator.

Proof. Let $c_{ik}^{(N)}(x)$ denote (i, k) element of $P_N X_N(x)$. First i_1, i_2, \ldots, i_N and k_1, k_2, \ldots, k_N are defined by $i - 1 = i_1 2^{N-1} + i_2 2^{N-2} + \cdots + i_{N-1} 2^1 + i_N 2^0$ and $k - 1 = k_1 2^{N-1} + k_2 2^{N-2} + \cdots + k_{N-1} 2^1 + k_N 2^0$, respectively. We put $I = i_1 + i_2 + \cdots + i_N$ and $K = k_1 + k_2 + \cdots + k_N$. Moreover $\{k_1, k_2, \ldots, k_N\}$ is divided into two sets such as $\{k_{m_1}, k_{m_2}, \ldots, k_{m_K} : k_u = 1 \text{ for any } u \in \{m_1, m_2, \ldots, m_K\}\}$ and $\{k_{l_1}, k_{l_2}, \ldots, k_{l_{N-K}} : k_v = 0 \text{ for any } v \in \{l_1, l_2, \ldots, l_{N-K}\}\}$. Then we see that

where $x_{(1)} = x$ and $x_{(0)} = 1$. In a similar way, we have

$$c_{ki}^{(N)}(x) = (p_K x + q_K)^I.$$

So we have

$$P_N X_N(x) = {}^t (P_N X_N(x)),$$

if and only if

$$c_{ik}^{(N)}(x) = c_{ki}^{(N)}(x),$$

for $1 \leq i, k \leq 2^N$. For thermore it is shown that the last equation is equivalent to

$$(p_I x + q_I)^K = (p_K x + q_K)^I$$

for $1 \leq I, K \leq N$. Combaining the above result with Theorem 1 in [15], we obtain the desired conclusion.

For N = 2 case (the DK model), Theorem 5 gives

$$(p_1x + q_1)^2 = p_2x + q_2.$$

This result was shown in [8, 15, 16] by using different methods.

5. Convergence theorem

In order to obtain a convergence theorem for the TPCA with $\{p_n : 0 \le n \le N\}$ having a site-bond representation with $\alpha = p_1^2/(2p_1-p_2)$ and $\beta = (2p_1-p_2)/p_1$, which corresponds to the result given by [8], a new process η_n is introduced as follows. For simplicity, we first consider the DK model (N = 2 case) with $(p_1, p_2) \in D_*$. Put $p_* = \max\{p_1, p_2\}$. A new process defined below is called p_* -DK-dual. We can see the thinning-relationship by coupling the DK model and p_* -DK-dual. We split both the DK model and the p_* -DK-dual into two phases, and we will allow the first phase to occur at times n + (1/2) for $n \in \mathbb{Z}_+$.

- (i) Let μ be the distribution of the p_* -DK-dual at time 0.
- (ii) At time n = 1/2, it undergoes a p_* -thinning. In general, for $p \in [0, 1]$, the *p*-thinning of a set $A \subset \mathbb{Z}$ is the random subset of A obtained by independently removing each element of A with probability 1 p.
- (iii) Start the DK model at time n = 0 with the same configuration as the p_* -DK-model (which is defined by f(0) = 0, $f(1) = p_1/p_*$, $f(2) = p_2/p_*$) at time n = 1/2.
- (iv) Couple the processes together until time $n_0 (1/2)$ for the DK model, n_0 for the p_* -DK-model. This can be done because the transitions for the DK model are the same as those for the p_* -DK-model lagged by time unit 1/2.
- (v) Perform a p_* -thinning for the DK model at time n_0 .

The distribution of the DK model started and ended as a p_* -thinning of the p_* -DKdual. As in the DK model, we can define a new process η_n as p_* -*TPCA-dual* for the *N*-neighbor TPCA, where $p_* = \max\{p_1, p_2, \ldots, p_N\}$. Recall that the following duality holds for the TPCA ξ_n^A and the p_* -TPCA-dual η_n^A starting from A, (see Theorem 2 (2) in [15]). **Theorem 6** ([15]) We assume an *N*-neighbor TPCA with $\{p_n : 0 \le n \le N\}$ having a site-bond representation with $\alpha = p_1^2/(2p_1 - p_2)$ and $\beta = (2p_1 - p_2)/p_1$. Suppose $(p_1, p_2) \in D_*$. For any A, B with $|A| < \infty$ or $|B| < \infty$, we have

$$E\left(\tilde{x}^{|\xi_n^A \cap B|}\right) = E\left(\tilde{x}^{|\eta_n^B \cap A|}\right),$$

for any $n \ge 0$, if $\tilde{x} = 1 - (2p_1 - p_2)p_*/p_1^2$.

Note that if $p_1 > p_2$ (non-attractive case for the TPCA with a site-bond representation), then $p_* = p_1$. By using the same argument for the DK model in Section 7 of [8], we can get the following convergence theorem for the TPCA:

Theorem 7 Assume $p_1 > p_2$ with $(p_1, p_2) \in D_*$. If the initial measure ν of the TPCA is a.s. (almost sure) infinite, then we have

$$\xi_n^{\nu} \to \mu_{\eta},$$

as $n \to \infty$, where the limit measure is uniquely determined by $E((-(p_1-p_2)/p_1)^{|\mu_\eta \cap A|}) = P(|\eta_{\infty}^A| = 0)$ for any A with $|A| < \infty$.

6. Conclusions and discussions

In this work we have presented rigorous results on the site-bond representation, the set-to-set connectedness, the self-duality, the matrix expression, and the convergence theorem for an N-neighbor TPCA with $\{p_n : 0 \le n \le N\}$, where $p_0 = 0$ and $N \ge 2$. An interesting feature of our model is that the dominant parameters on some properties are only p_1 and p_2 among $\{p_n : 0 \le n \le N\}$, see Proposition 1 and Corollary 2, for examples.

Arrowsmith and Essam [17] gave an expansion formula for a point-to-point connectedness for the oriented mixed site-bond percolation, in which each term is characterized by a graph. Konno and Katori [6] extended this formula to N = 2 case (the DK model). So it is shown that site-bond representation, self-duality, and the above graphical expansion formula hold in the DK model. Thus one of the interesting future problems is to extend the relation to a general N-neighbor TPCA considered here.

Finally we mention a relation between our discrete-time model and a continoustime one corresponding to it. In the continuous-time case, an infinitesimal generator for the model corresponds to P_N in Theorem 5. In fact, Eq. (8) in Theorem 5 corresponds to Eq. (17) in [18] for N = 2 case (an extension of their result to an N-neighbor case can be easily obtained). Recent works on the duality for continuous-time interacting particle systems are presented in [19, 20, 21, 22].

Acknowledgments. We would like to thank Akinobu Shimizu for useful discussions. This work was partially financed by the Grant-in-Aid for Scientific Research (B) (No.12440024) of Japan Society of the Promotion of Science.

- [1] Wolfram S, 2002 A New Kind of Science (Champaign, IL: Wolfram Media)
- [2] Domany E and Kinzel W, 1984 Phys. Rev. Lett. 53 311
- [3] Bagnoli F, 1992 Int. J. Mod. Phys. C 3 307
- [4] Bagnoli F and Rechtman R, 1999 Phys. Rev. E 59 R1307
- [5] Katori M, Konno N and Tanemura H, 2000 J. Stat. Phys. 99 603
- [6] Konno N and Katori M, 2000 J. Stat. Phys. 101 747
- [7] Katori M, Konno N and Tanemura H, 2002 Ann. Probab. 30 933
- [8] Katori M, Konno N, Sudbury A and Tanemura H, 2004 J. Theoret. Probab. 17 131
- [9] Bagnoli F, Boccara N and Rechtman R, 2001 Phys. Rev. E 63 046116
- [10] Atman A P F, Dickman R and Moreira J G, 2003 Phys. Rev. E 67 016107
- [11] Liggett T M, 1985 Interacting Particle Systems (New York: Springer)
- [12] Durrett R, 1988 Lecture Notes on Particle Systems and Percolation (Belmont, CA: Wadsworth)
- [13] Konno N, 1994 Phase Transitions of Interacting Particle Systems (Singapore: World Scientific)
- [14] Liggett T M, 1999 Stochastic Interacting Systems: Contact, Voter and Exclusion Processes (New York: Springer)
- [15] Konno N, 2002 J. Stat. Phys. 106 915
- [16] Konno N, 2002 J. Stat. Phys. 106 923
- [17] Arrowsmith D K and Essam J W, 1977 J. Math. Phys. 18 235
- [18] Sudbury A W and Lloyd P, 1995 Ann. Probab. 23 1816
- [19] Bandt C, 1999 J. Stat. Phys. 96 883
- [20] Sudbury A W, 2000 J. Theoret. Probab. 13 695
- [21] López F J and Sanz G, 2000 Markov Processes Relat. Fields 6 305
- [22] Hooyberghs J and Vanderzande C, 2001 Phys. Rev. E 63 041109