

Persistent superfluid phase in a three-dimensional quantum XY model with ring exchange

R. G. Melko and D. J. Scalapino

Department of Physics, University of California Santa Barbara, California 93106

(Dated: March 23, 2022)

We present quantum Monte Carlo simulation results on a quantum $S = 1/2$ XY model with ring exchange (the J - K model) on a three-dimensional simple cubic lattice. We first characterize the ground state properties of the pure XY model, obtaining estimations for the energy, spin stiffness and spin susceptibility at $T = 0$ in the superfluid phase. With the ring exchange, we then present simulation data on small lattices which suggests that the superfluid phase persists to very large values of the ring exchange K , without signatures of a phase transition. We comment on the consequences of this result for the search for various exotic phases in three dimensions.

The search for microscopic spin $1/2$ Hamiltonians that exhibit a spin liquid phase continues with the hope that these studies will provide insight into the type of materials that can support such a phase. Focus has recently turned to multi-particle ring exchange models,^{1,2,3} which have been identified as promising candidates for the presence of spin liquid as well as other exotic phases and critical points. One such Hamiltonian is an easy-plane $S = 1/2$ J - K model on the square lattice, with an XY exchange coupling J and a four-site ring exchange coupling K . Quantum Monte Carlo studies¹ of this model have shown that, at a critical value $(K/J)_c$, there appears to be a continuous quantum critical point separating an XY-superfluid phase from a valence bond solid (VBS) phase. Recently, Senthil *et al.*⁴ have proposed that at this type of critical point – separating conventional phases described in terms of different broken symmetries – one can have an isolated spin-liquid point characterized by an emergent global $U(1)$ symmetry and deconfined spinon excitations. In two dimensions, the theory finds that the $U(1)$ spin liquid is unstable away from the critical point because of the proliferation of monopole excitations, which are only absent precisely at $(K/J)_c$. They suggest⁴ that it is this instability that is responsible for the formation of the VBS phase seen in the simulations¹ for $K/J > (K/J)_c$. However, in three dimensions, monopole excitations can be gapped, so within the theoretical framework⁴ an extended region of stable $U(1)$ spin liquid phase could exist.⁵ Motivated by this possibility, we have carried out a quantum Monte Carlo study of the simple cubic J - K model to look for signatures of a zero-temperature phase transition out of the superfluid phase (at large J/K) to a quantum disordered or insulating phase (at large K/J).

The Hamiltonian under study is

$$H = -J \sum_{\langle ij \rangle} B_{ij} - K \sum_{\langle ijkl \rangle} P_{ijkl}, \quad (1)$$

where bond and plaquette exchange operators are defined as

$$B_{ij} = S_i^+ S_j^- + S_i^- S_j^+ = 2(S_i^x S_j^x + S_i^y S_j^y), \quad (2)$$

$$P_{ijkl} = S_i^+ S_j^- S_k^+ S_l^- + S_i^- S_j^+ S_k^- S_l^+. \quad (3)$$

Here, $\langle ij \rangle$ denotes a pair of nearest-neighbor sites and $\langle ijkl \rangle$ are sites on the corners of a plaquette (see Fig. 1). In three dimensions, each spin is shared by six bonds and twelve plaquettes, which form the faces of the simple cubic lattice. For $K = 0$ this is the standard quantum spin- $1/2$ XY-model, or, equivalently, a hard-core boson model at half-filling, studied previously using quantum Monte Carlo techniques by Pedersen and Schneider.⁶ The model is analogous to a quantum lattice gas model of He^4 , and undergoes a lambda-transition at $T/J \approx 2$ to a bulk superfluid state.^{6,7} The K -term corresponds to retaining only the x - and y -spin operators of the full four-spin cyclic exchange studied in relation to some high- T_c parent compounds.⁸




FIG. 1: The labeling of the plaquette indices in Eq. (1) are illustrated on left. At right, three intersecting plane segments of the cubic lattice show that each spin site shares six bonds and twelve plaquettes with other sites.

Our simulations use the stochastic series expansion (SSE) quantum Monte Carlo method,^{9,10} which is based on importance sampling of elements of the power series expansion of the partition function. The method is numerically exact within statistical error bars, and has broad applicability to a wide range of spin and boson models where the sign problem can be avoided. In order to efficiently sample off-diagonal J terms in the Hamiltonian, the recently developed “directed-loop”^{11,12} algorithm is used. However, for large values of the parameter K , the loop moves become ineffective, and a new sampling procedure was developed.¹³ This new “multi-branch cluster” update is a Swendsen-Wang or Wolff-type sampling scheme generalized to flip clusters of plaquette operators in the SSE operator-list. It is observed to significantly decrease autocorrelation times of the simulations for large K , even for $K/J \rightarrow \infty$ in two dimen-

sions. Details of the SSE algorithm, simulation procedure and calculation of physical observables can be found in Ref. 13.

The energy per spin is simply calculated in terms of the SSE estimator for the order of the power series expansion, n , and the temperature:

$$E = -\frac{\langle n \rangle T}{N}, \quad (4)$$

where $N = L^3$ is the system size. The main quantity of interest in this paper is the spin stiffness,

$$\rho_s = \frac{\partial^2 E(\phi)}{\partial \phi^2} \quad (5)$$

$$= \frac{T (\langle W_x^2 \rangle + \langle W_y^2 \rangle + \langle W_z^2 \rangle)}{3N}, \quad (6)$$

where ϕ is a twist imposed on bonds in either the x , y or z lattice directions. In Eq. (6) we calculate the derivative at $\phi = 0$ using the winding number estimators $W_{x,y,z}$ which measure the net spin current across the periodic boundaries in each direction.^{10,13} We have also calculated in some cases the spin susceptibility χ , easily obtainable in the S^z basis using the squared magnetization estimator,

$$\chi = \frac{1}{TN} \left\langle \left(\sum_{i=1}^N S_i^z \right)^2 \right\rangle. \quad (7)$$

In three dimensions, even the smallest useful lattice sizes have Hilbert spaces far too large to allow an exact diagonalization study to be performed. In order to test the Monte Carlo and to characterize the $K = 0$ ground state, we study the asymptotics of the pure XY model, setting $J = 0.5$ in the Hamiltonian Eq. (1). Second-order spin-wave calculations carried out by Weihong *et al.*¹⁴ give numerical values for the ground state energy per spin and spin susceptibility as $E_0 \approx -0.7914$ and $\chi \approx 0.1517$, without error bars. Although we are not aware of any detailed numerical results for the zero-temperature spin stiffness, mean field theory predicts a value of $\rho_s = S^2 = 0.25$ using Eq. (5). One might expect that quantum fluctuations slightly increase the value of ρ_s in the full $S = 1/2$ XY model, as observed in two dimensions.¹⁵

We simulated the pure XY model ($K = 0$, $J = 0.5$) for several lattice sizes, $L = 6, 8, 10, 12$, and 14 , using the full SSE code, for several values of T/J down to 0.04 . The relevant estimators either showed absolute temperature convergence to within error bars, or were extrapolated to $T = 0$. Data obtained from the SSE simulation for the ground-state energy per spin, E_0 , is most accurate. Chiral perturbation theory¹⁶ predicts that the energy depends on system size and spatial dimension (d) as

$$E_0(L) - E_0 \sim \frac{C}{L^{d+1}}, \quad (8)$$

where C is some constant. This finite-size dependence has been shown to be in good agreement with similar




FIG. 2: Ground-state energy versus $1/L^4$ for system sizes $L = 6, 8, 10, 12$ and 14 . The curve is a fit to Eq. (8).

quantum Monte Carlo studies of the two-dimensional (2D) XY model.¹⁵ Setting $d = 3$ in Eq. (8), we use our data to obtain an extrapolation to $L \rightarrow \infty$, giving $E_0 = -0.79182(2)$. The fit is illustrated in Fig. 2, which has a minimum in the chi-squared value per degree of freedom for $C = -1.14$. Data for the spin susceptibility and spin stiffness have considerably larger error bars, within which the larger system size data is converged, making a finite-size scaling extrapolation irrelevant. Our best estimates for the ground state values in these cases are $\chi = 0.1500(2)$ and $\rho_s = 0.2623(2)$




FIG. 3: The finite-temperature behavior of the spin stiffness. Data for both $L = 6$ and $L = 24$ is included for $K/J = 0$ to illustrate finite-size effects at the lambda-transition.

Turning to the full model, we performed simulations including the K term in Eq. (1), keeping J fixed at $1/2$. First, the $T \rightarrow 0$ temperature convergence was established for each value of K/J simulated. Fig. 3 compares the finite-temperature behavior of the spin stiffness for an $L = 6$ system at $K/J = 20$ versus the XY-point at $K/J = 0$. Results for the ground-state spin stiffness ver-

sus K/J are illustrated in Fig. 4. Data for $L = 4$ and $L = 6$ up to $K/J = 48$ was taken at temperatures below $T/J = 0.5$, depending on the specific K/J where good $T \rightarrow 0$ convergence was established. However, the three largest K/J values for $L = 6$ began to show sticking, evidence that the Monte Carlo time-scales are getting very large or ergodicity is being lost. To combat this, data for $L = 8$ was taken at slightly higher temperature, $T/J < 2.0$ (for large K/J), which was nonetheless within the range of the zero-temperature convergence.

For the 2D J - K model, the zero-temperature critical point separating the superfluid phase from the VBS occurs at $(K/J)_c \approx 8$.¹ However, as illustrated in Fig. 4, for the three-dimensional (3D) case the superfluid phase is clearly much more robust. In particular, extensive simulations for the $L = 6$ system, out to $K/J = 144$ fail to show an abrupt transition out of the superfluid phase. Similarly, data for $L = 8$ remains strongly superfluid for K/J up to at least 60. As evident from Fig. 4, finite-size effects for the spin stiffness become much more pronounced as K/J gets large. Further, observation of the staggered magnetization showed no indications of the onset of any Néel (or boson charge-density wave) ordered state at any K/J which was simulated. Unfortunately, algorithm performance becomes less efficient for this model for large values of K/J . This, coupled with the fact that the lattice-plaquette coordination contributes an extra factor of three in the CPU time size scaling of the algorithm in three dimensions as compared to two dimension, makes a more detailed study of larger lattice sizes impractical at this time.




FIG. 4: Ground-state spin stiffness for the 3D J - K model. The XY exchange coupling was set to a constant, $J = 0.5$. Open symbols indicate data which showed signs of ergodicity loss, and therefore error bars may be under-estimated.

The consistency of data for the two largest system sizes points toward robust superfluidity in the system well above the value for $(K/J)_c$ where the onset of insulating behavior occurs in the 2D model.¹ Although impossible to prove without a detailed finite-size scaling study, we believe that the data supports the conjecture that the superfluid ground state persists in the model, becoming critical only at $K/J \rightarrow \infty$. This of course leaves the interesting question regarding the nature of the $K/J \rightarrow \infty$ phase open for future study. One suggestion is that the $J = 0$ model actually has long-range in-plane ferromagnetic order, in accord with a simple spin-wave analysis.¹⁷ The critical value for the spin stiffness in this case would be $\rho_s = 0$, however for finite J as in Fig. 4, ρ_s may be non-zero in the thermodynamic limit.

In summary, we have presented data from SSE quantum Monte Carlo simulations of the 3D simple cubic J - K model. Simulations of the XY point reveal a ground state energy, spin susceptibility, and spin stiffness of

$$\begin{aligned} E_0 &= -0.79182(2), \\ \chi &= 0.1500(2), \\ \rho_s &= 0.2623(2), \end{aligned}$$

consistent with predictions. In particular, the ground state energy agrees with second-order spin-wave theory¹⁴ to within 0.05%. The full J - K model appears to remain superfluid for all finite K/J , with no phase transition signature up to $K/J = 144$. The large K insulating phase is still unknown, however no indications of either a charge-density wave or any quantum disordered state are present. The robustness of the superfluid phase is consistent with the notion that the relative strength of quantum fluctuations introduced by the ring-exchange (K) term is reduced as one increases the dimensionality of the system. Finally, the failure of this model to support an emergent $U(1)$ spin liquid illustrates the difficulty in realizing exotic phases in microscopic quantum models suggested by arguments arising from generalized actions.

Acknowledgments

The authors would like to thank T. Senthil for motivating discussions. We appreciate extensive and helpful collaboration with A. Sandvik. Thanks also to L. Balents, A. Burkov, M. P. A. Fisher, M. Hermele, and O. Motrunich for useful conversations. Supercomputer time was provided by NCSA under grant number DMR020029N. This work was also supported by the Department of Energy, Grant No. DOE85-45197.

¹ A. Sandvik, S. Daul, R.R.P. Singh, and D.J. Scalapino, Phys. Rev. Lett. **89**, 247201 (2002).

² L. Balents, M.P.A. Fisher, and S.M. Girvin, Phys. Rev.

B **65**, 224412 (2002); V. Rousseau, G. G. Batrouni, and R. T. Scalettar, cond-mat/0312018; M. Hermele, M. P. A. Fisher, and Leon Balents, Phys. Rev. B **69**,

- 064404 (2004); V. Gritsev, B. Normand, and D. Baeriswyl, Phys. Rev. B **69**, 094431 (2004).
- ³ A. Paramekanti, L. Balents and M. P. A. Fisher, Phys. Rev. B **66**, 054526 (2002).
- ⁴ T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, Science **303**, 1490 (2004); T. Senthil, L. Balents, S. Sachdev, A. Vishwanath, and M. P. A. Fisher; cond-mat/0312617.
- ⁵ D. A. Huse, W. Krauth, R. Moessner, and S. L. Sondhi, Phys. Rev. Lett. **91**, 167004 (2003); R. Moessner and S. L. Sondhi, Phys. Rev. B **68**, 184512 (2003); O. I. Motrunich and A. Vishwanath, cond-mat/0311222; O. I. Motrunich and T. Senthil, cond-mat/0407368.
- ⁶ M. H. Pedersen and T. Schneider, Phys. Rev. B **53**, 5826 (1996).
- ⁷ D. D. Betts, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and M. S. Green (Academic, London, 1974), Vol. 3, pp. 569-652.
- ⁸ M. Roger and J. M. Delrieu, Phys. Rev. B **39**, 2299 (1989); Y. Honda, Y. Kuramoto, and T. Watanabe, Phys. Rev. B. **47**, 11329 (1993). J. Lorenzana, J. Eroles, and S. Sorella, Phys. Rev. Lett. **83**, 5122 (1999); R. Coldea, *et al.* Phys. Rev. Lett. **86** 5377 (2001); B. Normand, and A. M. Oles, cond-mat/0403481.
- ⁹ A.W. Sandvik and J. Kurkijärvi, Phys. Rev. B **43**, 5950 (1991); A.W. Sandvik, J. Phys. A **25**, 3667 (1992).
- ¹⁰ A.W. Sandvik, Phys. Rev. B **56**, 11678 (1997).
- ¹¹ A.W. Sandvik, Phys. Rev. B **59**, R14157 (1999).
- ¹² O.F. Syljuåsen and A.W. Sandvik, Phys. Rev. E **66**, 046701 (2002).
- ¹³ R. G. Melko and A. W. Sandvik, in preparation.
- ¹⁴ Z. Weihong, J. Oitmaa, and C. J. Hamer, Phys. Rev. B **44**, 11869 (1991).
- ¹⁵ A. W. Sandvik and C. J. Hamer, Phys. Rev. B **60**, 6588 (1999).
- ¹⁶ P. Hasenfratz and F. Niedermayer, Z. Phys. B **92**, 91 (1993).
- ¹⁷ L. Balents, private communication.