

Quantum criticality and the metal-insulator transition in 2D: a critical test.

B. Hosseinkhani and J. Zaanen

Instituut Lorentz for Theoretical Physics, Leiden University, POB 9506, 2300 RA Leiden, The Netherlands
(March 23, 2022)

Using recent insights obtained in heavy fermion physics on the thermodynamic singularity structure associated with quantum phase transitions, we present here an experimental strategy to establish if the zero-temperature transition in the disordered two dimensional gas is a real quantum phase transition. We derive a overcomplete set of scaling laws relating the density and temperature dependence of the chemical potential and the effective mass, which are in principle verifiable by experiment.

The observation of a metal-insulator transition (MIT) in the two dimensional electron/hole gasses with quenched disorder (2DE/HG's) [1] revived the interest in the nature of disordered, interacting fermion systems [2]. A consensus has emerged that the transition is between a disordered Wigner crystal (Wigner glass) at low electron density and a strongly interacting Fermi-liquid state at higher densities. However, the nature of the phase transition itself is subject of much debate. One viewpoint is that the interacting electron system near the transition is characterized by strong density inhomogeneities with puddles of high density electron gas immersed in a low density insulating background [3,4], suggesting a transition of the percolation kind [5]. On the other hand, it has been suggested that the transition is a genuine continuous quantum phase transition (QPT) [6], finding support in for instance the scaling collapse of the resistivity [1,2]. In the past only transport quantities were accessible by experiment, but recently it appeared possible to also measure thermodynamic quantities: the inverse compressibility [7–10] and the effective mass [11,12], both of which appear to diverge near the transition.

Thermodynamical properties are often easier to interpret than transport quantities. Very recently it was realized by Si, Rosch and coworkers in the context of heavy fermion physics that thermodynamics becomes extremely revealing in the vicinity of QPT's [13,14]. Based on almost trivial scaling considerations, they demonstrated that an unambiguous diagnostic can be constructed for a QPT, in the form of a plethora of scaling laws involving thermodynamic properties. The fact that thermodynamics is more informative at a strongly interacting QPT than at a thermal phase transition is caused by the peculiar role of temperature as an effective finite size in the former. The other important quantity is the zero-temperature control parameter, tuning the system through the QPT. This coupling constant determines which thermodynamic quantities reveal the quantum singularity. As we first realized in the context of cuprate superconductivity [15], the electron density is a particularly interesting control parameter, because the singu-

larity shows itself through the temperature- and density derivatives of the chemical potential, in addition to independent information contained in the specific heat. It are precisely these quantities which have become accessible by experiment in the MOSFET's and we will present here a recipe for how to use this information to uncover the QPT, if present in these systems.

The thermodynamic singularity structure of a QPT is of a different kind than the one encountered at thermal phase transitions. The role of temperature is taken by the coupling constant y , a quantity tuning the system through its zero temperature transition at y_c . We take as a working definition for a 'genuine' QPT that (a) at zero temperature the quantum dynamics in $D = d + z$ dimensional space time (z is the dynamical exponent) becomes scale invariant at y_c , and (b) this critical state is universal in the sense that it obeys hyperscaling. In Euclidean path-integral representation, temperature sets the inverse compactification radius of the imaginary time direction and acts therefore as a finite size. These postulates can be compactly represented in the form of a scaling relation for the singular part of the free energy,

$$F_s(r, T) = b^{-(d+z)} F_s(b^{y_r} r, b^z T) \quad (1)$$

where $r = (y - y_c)/y_c$, the reduced coupling constant. Following Zhu *et al.* [13], Eq. (1) is equivalent to the following scaling forms for the free energy density,

$$\begin{aligned} F_s(r, T) &= -\rho_0 r^{(d+z)/y_r} \tilde{f}\left(\frac{T}{T_0 r^{z/y_r}}\right) \\ &= -\rho_0 \left(\frac{T}{T_0}\right)^{(d+z)/z} f\left(\frac{r}{(T/T_0)^{y_r/z}}\right) \end{aligned} \quad (2)$$

where ρ_0 and T_0 are non-universal constants, while $f(x)$ and $\tilde{f}(x)$ are universal scaling functions. The first form is useful in the low temperature regime of the (dis)ordered phase in the proximity of the QPT, while the second form describes the quantum critical regime itself. Accordingly, since there is no singularity at $r = 0, T > 0$, the 'quantum critical' function f can be expanded as $f(x \rightarrow 0) \simeq f(0) + x f'(0) + (1/2) x^2 f''(0) + \dots$. On the other hand, in the stable phase the scaling function

is expanded as $\tilde{f}(x) = \tilde{f}(0) + g(x)$ where $g(x)$ describes the low temperature thermodynamics of the phases to the left- or right side of the QPT. By taking derivatives to the coupling constant and/or temperature one easily derives the thermodynamic quantities. The fact that temperature enters through finite size scaling is a blessing; the outcome is a highly overcomplete set of scaling laws [13,15] relating the zero- and finite temperature exponents of the thermodynamic quantities via y_r and z .

Let us now apply this general wisdom to the 2D MIT. The coupling constant clearly corresponds with the carrier density n while the transition occurs at a critical n_c such that the reduced coupling constant is $r = (n - n_c)/n_c$. The thermodynamic quantities associated with the density are the chemical potential $\mu = \partial F/\partial n$ and the inverse compressibility (or incompressibility) $(1/\kappa) = (1/n)(\partial^2 F/\partial n^2)$. In addition one has the specific heat $C = -T(\partial^2 F/\partial T^2)$ and the temperature derivative of μ , $\Upsilon_n = (\partial\mu/\partial T) = (\partial^2 F/\partial T\partial n)$. In principle, one can derive a total of 9 scaling forms for the singular parts of $1/\kappa$, η and C , applicable to the quantum critical regime ($r = 0$, finite T) and the low temperature regimes ($T \rightarrow 0$, $r \neq 0$) of the two phases in the proximity of the QPT, respectively. Although the physics is in any other regard very different, the 2d MIT shares the density as coupling constant with the (suspected) QPT in optimally doped cuprates [16] and one can find a full set of scaling forms in a paper [15] dealing with the specifics of the cuprates.

Another difference with the cuprates is that the quantities of crucial importance (compressibility, chemical potential) are now routinely measured in the MOSFET's [7–10] while this information is not available yet in the cuprates [15]. In the remainder we will be primarily interested in the high density side of the MIT. All we have to know is the function $g(x)$ in this regime, and this can be deduced from the low temperature behavior of the specific heat C . Because of their ‘lack of volume’ C cannot be measured directly and we will assume that this state is ‘like a Fermi-liquid’ in the sense that the following Fermi-liquid relation is still valid: $(C/C_0) = 1 + F_1^s/3 = (m^*/m)$, where $C_0 = \nu T$, the specific heat in the non-interacting limit (ν is the density of states) while F_1^s is the Landau parameter which also governs the transport effective mass m^*/m . This effective mass is directly measured in the magneto-transport experiments [12], and found to be finite at zero temperature, suggesting a linear specific heat $C = \gamma T$. It is easily checked that this implies $g(x) = cx^{y_0+1}$ (massless) spectrum with $y_0 = 1$.

The scaling relations for the various quantities are easily obtained and let us directly specialize to the case of the ‘Fermi-liquid like’ regime in the proximity of the 2D MIT ($d = 2$ and $y_0 = 1$). The unknown exponents are the dynamical exponent z and the coupling constant (electron density) exponent y_r . It is convenient

to reparametrize the latter in terms of an exponent having a similar status as the specific heat exponent α in the case of a thermal phase transition [15],

$$\alpha_r = 2 - (2 + z)/y_r \quad (3)$$

The singular, low temperature contributions to the various thermodynamics quantities in the liquid have the following form,

$$\begin{aligned} C_{cr}(T \rightarrow 0, r) &= \gamma_{cr} T \\ \Upsilon_{r,cr}(T \rightarrow 0, r) &= v_{cr} T \\ \frac{1}{\kappa_{cr}(T \rightarrow 0, r)} &= K_{cr,0} + K_{cr,T} T^2 \end{aligned} \quad (4)$$

where the various coefficients which depend critically on $r = (n - n_c)/n_c$ are given by,

$$\begin{aligned} \gamma_{cr} &= \frac{2\rho_0 c}{T_0^2} r^{(2-\alpha_r)(2-z)/(2+z)} \\ v_{cr} &= -\frac{2\rho_0 c}{T_0^2} \frac{(2-z)}{(2+z)} (2 - \alpha_r) r^{(2-\alpha_r)(2-z)/(2+z)-1} \\ K_{cr,0} &= -\rho_0 \tilde{f}(0)(1+z)(2 - \alpha_r) r^{-\alpha_r} \\ K_{cr,T} &= -\frac{c\rho_0}{T_0^2} (2-z)((2 - \alpha_r) \left(\frac{2-z}{2+z}\right) - 1) \\ &\quad \times r^{(2-\alpha_r)(2-z)/(2+z)-2} \end{aligned} \quad (5)$$

One infers that the divergence of the prefactors γ, v, K_T of the temperature dependent parts of C, Υ and $1/\kappa$ are all governed by the fundamental mass exponent defined through $\gamma \sim m^*/m = r^{\alpha_m}$ with

$$\alpha_m = (2 - \alpha_r)(2 - z)/(2 + z), \quad (6)$$

such that $v \sim r^{\alpha_m-1}$ and $K_T \sim r^{\alpha_m-2}$. These relations between the exponents are just a consequence of the assumption that r enters the free energy in the form of a power law and the Fermi-liquid form for \tilde{f} . Hence, by measuring with high precision the temperature dependence of the chemical potential and incompressibility one can critically test if the system is like a Fermi-liquid with a mass diverging at the MIT. According to the transport measurements [11,12] $m^* \sim r^{\alpha_m}$ with $\alpha_m = -0.5 \pm 0.1$ and when the interpretations are correct it has to be that $v \sim r^{-1.5}$ and $K_T \sim r^{-2.5}$.

In fact, the first genuine scaling laws in this liquid regime can be deduced from universal *amplitude ratio's*. Usually amplitudes are non-universal but Zhu et al. [13] realized that in the case of pressure driven QPT's the Grueneisen ratio becomes universal. The analogon of the Grueneisen ratio in the case of the density driven transition is the ratio of Υ and C which is expressed entirely in the exponents while the non-universal factors (ρ_0, T_0, c) drop out,

$$\Gamma_r = \frac{\Upsilon_{r,cr}}{C_{cr}} = \left(\frac{(2-z)(2-\alpha_r)}{2+z} \right) r^{-1} \quad (7)$$

In addition, Eq. (5) implies a second universal ratio which has not been identified before,

$$\Gamma'_r = \frac{T\nu}{K_T} = \left(\frac{2(2 - \alpha_r)}{(2 - \alpha_r)(2 - z) - 2 - z} \right) r^{-1} \quad (8)$$

It follows from the scaling analysis that the divergence of the zero-temperature incompressibility K_0 is governed directly by the elementary ‘energy exponent’ α_r – this is no wonder since the incompressibility has the same status as a specific heat when the roles of temperature and density are exchanged. Hence, the zero temperature incompressibility is the quantity which reveals the quantum singularity most directly in case of density-driven transitions. As an interesting aside, the screening length $s \sim 1/\kappa$ in a 2D charged liquid and this length is therefore governed at zero temperature by the exponent α_r . One could be tempted to identify the screening length with the correlation length but this is not quite right: the correlation length is governed instead by the exponent $\nu = 1/y_r$. In fact, the correlation length is the length scale associated with order parameter correlations and the screening length has a completely different, ‘truly thermodynamic’ status. There might well be an order parameter at work at the MIT but we have left it completely implicit in the present analysis which is equivalent to the finite size scaling of the specific heat singularity at a thermal phase transition.

Up to this point we have indentified three scaling laws governing the liquid: one relating α_r to the mass exponent α_m via Eq. (6), and the universal ratio’s Eq.’s (7,8). This is not all, because an other set of scaling relations can be deduced governing the quantum critical regime itself. At the quantum-critical point the scales associated with the stable fixed points should collapse. Approaching the MIT from the metallic side the situation is in this regard analogous to what is found in the heavy fermion systems: the Fermi-energy (the scale of the Fermi-liquid) has to vanish [17], or equivalently, the effective mass has to diverge. In the quantum critical regime the temperature dependence of the thermodynamic properties are governed by a different set of scaling laws which are also governed by the fundamental exponents (z, α_r) . The quantum critical regime can be accessed right at the QPT, but also away from the critical point, as long as temperature exceeds the Fermi-energy [6]. Hence, if the MIT is governed by a genuine QPT it has to be that upon approaching the phase transition a temperature window opens up of universal behaviors bounded from the below by the Fermi-energy and from the above by a cross-over to a non-universal ultraviolet.

The temperature dependences of C , Υ and $1/\kappa$ in this quantum critical regime are easily deduced from the second form for the singular part of the free-energy in Eq. (2),

$$\begin{aligned} C_{cr}(T, r=0) &= 2\rho_0 f(0) \frac{(2+z)}{z^2} \left(\frac{T}{T_0} \right)^{2/z} \\ \Upsilon_{cr}(T, r=0) &= -\frac{\rho_0 f'(0)}{T_0} \frac{1 - \alpha_r}{2 - \alpha_r} \frac{(2+z)}{z} \\ &\quad \times \left(\frac{T}{T_0} \right)^{(2(1-\alpha_r)-z)/(z(2-\alpha_r))} \\ \frac{1}{\kappa_{cr}(T, r=0)} &= -\rho_0 f''(0) \left(\frac{T}{T_0} \right)^{((2+z)\alpha_r)/(z(2-\alpha_r))} \end{aligned} \quad (9)$$

It follows that the temperature exponent of the specific heat directly reveals the dynamical exponent z – a direct consequence of the role of temperature as a finite size. Since there is no a-priori relation between transport and specific heat in the quantum critical regime it appears that this quantity is beyond the reach of experiment. However, again the chemical potential is a valuable source of information. Its temperature and density derivatives should obey two other independent scaling laws.

In summary, we have identified five scaling relations depending on two unknown exponents (α_r, z) , governing the behaviors of quantities which can be measured in principle by present day experimental techniques, which should be obeyed if a genuine QPT is at work in the 2DEG’s. How does the above relate to the thermodynamic experiments, feasible in the MOSFET’s? High precision measurements of the effective mass and the chemical potential in the same samples are required, to establish if these quantities reveal divergences at the same critical density. The dynamical range should be improved to densities much closer to the transition and a main message of the present work is that the temperature dependence of especially the chemical potential contains valuable information.

What can be said on basis of the data which are at present available? The divergence of the effective mass seems to be reasonably well established, suggesting a mass exponent $\alpha_m = -0.5 \pm 0.1$ [11,12]. The incompressibility data suggest that K_0 diverges at some critical density [7–10]. However, the dynamical range covering this divergence is in the present data too small. Pending the choice of n_c we find that the incompressibility upturn can equally well be fitted with an α_r in the range 0.2 – 2 [18]. As we already emphasized, α_r is analogous to the specific heat exponent of thermal phase transitions, and these are generically small compared to unity. Even within the uncertainties, the present data seem to suggest that the incompressibility divergence is much stronger than the specific heat divergence at a typical classical phase transition. It is easy to derive some bounds on the allowed values of the exponents – if these are violated one is surely not dealing with a QPT. The fundamental exponent is the coupling constant dimension $y_r = (d+z)/(2-\alpha_r)$. To ensure the relevancy of the coupling constant $y_r > 0$ and this implies an upper

bound $\alpha_r < 2$.

Another bound follows from the condition that both the effective mass and the incompressibility are diverging at the transition. The divergence of the mass implies $\alpha_m = -|\alpha_m|$, while a diverging incompressibility needs $\alpha_r > 0$. Using the scaling law Eq. (6) to express α_r in terms of α_m as $\alpha_r = 2 + |\alpha_m|(2+z)/(2-z)$, we find a lower bound on $z > (2 + |\alpha_m|)/(1 - |\alpha_m|/2)$. For instance, taking the experimental value $|\alpha_m| \simeq 1/2$, it follows that $z > 10/3$. This is a surprisingly large lower bound, once again demonstrating that quantum phase transitions involving fermions have little in common with their bosonic- or classical siblings [17].

Summarizing, we have presented in the above a powerful diagnostic to establish whether the metal-insulator transition in 2DEG's is a genuine quantum phase transition. We are not advocating the view that *it has to be* a quantum phase transition. The only way to find it out is by more and better experiments. Alternatively, one could take the viewpoint that the MIT cannot possibly be a QPT given the intrinsic tendencies of the system to micro-phase separate. The various arguments [3,4] in support of this viewpoint are persuasive, but only so to the extent that the starting assumption is satisfied: a good metal separated from an insulating crystal by a first order transition which is in turn 'masked' by electrostatic effects and/or the disorder potential. In this context, the 'quality' of the metal refers to its screening length being small. In 2D, the screening length $s \sim 1/\kappa$ and according to a Fermi-liquid relation $\nu/\kappa = (m/m^*)(1 + F_0^s)$. Even when F_0^s can be ignored, the incompressibility and thereby the screening length has to diverge in the approach to the transition, because the effective mass is divergent. It is hard to imagine how a mass divergence can arise from inhomogeneities – it is most likely associated with the physics of the uniform liquid [19]. Stronger, as is obvious from the scaling analysis, the screening length and the mass are in principle governed by different exponents, which implies that also F_0^s might behave singularly. Microscopically this might have to do with the effect discovered by Das Sarma in the 1980's that quenched disorder enhances the screening length [20], a physics which was recently discussed in the context of the MIT by Si and Varma [21]. The bottomline is that the homogeneous liquid has an intrinsic tendency to become increasingly incompressible approaching the MIT, and thereby increasingly similar to the Wigner glass in its electrostatic responses. This in turn will tend to diminish the tendency towards the formation of inhomogeneities, potentially paving the road for a continuous quantum phase transition. In an upcoming publication we will ad-

dress these matters on a more quantitative level.

Acknowledgements We acknowledge fruitful discussions with T. Klapwijk, S. Das Sarma, C.M. Varma, Q. Si and J.M.J. van Leeuwen.

-
- [1] S.V. Kravchenko *et al.*, Phys. Rev. B **51**, 7038 (1995); Phys. Rev. Lett. **77**, 4938 (1996).
 - [2] E. Abrahams, S.V. Kravchenko and M.P. Sarachik, Rev. Mod. Phys. **73**, 251 (2001).
 - [3] S. He and X.C. Xie, Phys. Rev. Lett. **80**, 3324 (2001); J. Shi and X.C. Xie, Phys. Rev. Lett. **88**, 4951 (2002).
 - [4] B. Spivak, Phys. Rev. B **67**, 125205 (2003); B. Spivak and S. Kivelson, cond-mat/0310712; cond-mat/xxxxxx.
 - [5] S. Das Sarma *et al.*, cond-mat/0406655
 - [6] S. Sachdev, *Quantum Phase transitions* (Cambridge Univ. Press, 1999).
 - [7] S. Ilani, A. Yacoby, D. Mahalu and H. Shtrikman, Phys. Rev. Lett. **84**, 3133 (2000).
 - [8] S.H. Dultz and H.W. Jiang, Phys. Rev. Lett. **84**, 4689 (2000).
 - [9] S. Ilani, A. Yacoby, D. Mahalu and H. Shtrikman, Science **292**, 1354 (2001).
 - [10] S.H. Dultz, B. Alavi and H.W. Jiang, cond-mat/0210584.
 - [11] A.A. Shashkin, S.V. Kravchenko, V.T. Dolgoplov, and T.M. Klapwijk, Phys. Rev. B **66**, 073303 (2002).
 - [12] A.A. Shashkin, M. Rahimi, S. Anissimova, S.V. Kravchenko, V.T. Dolgoplov, and T.M. Klapwijk, Phys. Rev. Lett. **91**, 046403 (2003); A.A. Shashkin, S.V. Kravchenko, V.T. Dolgoplov, and T.M. Klapwijk, J. Phys. A: Math. Gen. **36**, 9237 (2003).
 - [13] L. Zhu, M. Garst, A. Rosch and Q. Si, Phys. Rev. Lett. **91**, 066404 (2003).
 - [14] R. Kuchler *et al.*, Phys. Rev. Lett. **91**, 066405 (2003).
 - [15] J. Zaanen and B. Hosseinkhani, Phys. Rev. B, in press (cond-mat/0403345).
 - [16] A.V. Chubukov and S. Sachdev, Phys. Rev. Lett. **71**, 169 (1993); C. Castellani, C. di Castro and m. Grilli, Phys. Rev. Lett. **75**, 4650 (1995); R.B. Laughlin, Adv. Phys. **47**, 943 (1998); C.M. Varma, Phys. Rev. Lett. **83**, 3538 (1999).
 - [17] J. Custers *et al.*, Nature **424**, 524 (2003); P. Coleman *et al.*, J. Phys.: cond. mat. **13**, R723 (2001).
 - [18] B. Hosseinkhani, *On the thermodynamics at the quantum phase transitions in two dimensional electron systems*, PhD thesis, Leiden University (2004).
 - [19] It was recently argued that this mass divergence might even reside in the clean limit: Y. Zhang and S. Das Sarma, cond-mat/0312565.
 - [20] S. Das Sarma, Phys. Rev. Lett. **50**, 211 (1983).
 - [21] Q. Si and C.M. Varma, Phys. Rev. Lett. **81**, 4951 (1998).