

Competition Between Charge-Density Waves and Superconductivity in Striped Systems

Enrico Arrigoni ^{a,*} Eduardo Fradkin ^b Steven A. Kivelson ^c

^a *Institute for Theoretical Physics, Graz University of Technology, Petersgasse 16, A-8010 Graz, Austria*

^b *Department of Physics, University of Illinois at Urbana-Champaign, 1110 W. Green St., Urbana, IL 61801-3080*

^c *Department of Physics, University of California at Los Angeles, 405 Hilgard Ave., Los Angeles, CA 90095*

Abstract

Switching on interchain coupling in a system of one-dimensional strongly interacting chains often leads to an ordered state. Quite generally, there is a competition between an insulating charge-density-wave and a superconducting state. In the case of repulsive interactions, charge-density wave usually wins over superconductivity.

Here, we show that a suitable modulation in the form of a period 4 bond-centered stripe can reverse this balance *even in the repulsive case* and produce a superconducting state with relatively high temperature.

Key words:

Superconductivity, Quasi-one-dimensional systems, competing phases

There is strong *theoretical* evidence that quasi-one dimensional ladder systems with purely repulsive electronic interaction, such as two-leg Hubbard or $t - J$ ladders, show a strong tendency toward the formation of a large spin gap associated with substantial superconducting (SC) “d-wave-like” pair-field correlations [1,2,3,4], i. e. the pairing susceptibility diverges as

$$\chi_{SC} \sim \Delta_s / T^{2-1/K} . \quad (1)$$

Here, Δ_s is the spin gap, T the temperature and K the charge Luttinger parameter of the ladder. One could thus imagine to obtain a “true” superconductor with long-ranged correlations by coupling an infinite number of ladders into a two-dimensional array. However, besides SC, also charge-density-wave (CDW) susceptibility with wave vector $4K_F$ diverges with a power law in the temperature:

$$\chi_{CDW} \sim \Delta_s / T^{2-K} . \quad (2)$$

Numerical calculations [5,6,7,8] have shown that, except in a very small region in parameter space, K is smaller than 1 for strong repulsive interactions, so that CDW susceptibility diverges stronger than SC. For this reason, upon switching on a weak coupling between the ladders, one expects CDW correlations to condense before the SC ones, so that in the end one obtains a CDW insulator instead of a superconductor. This can be seen from the RG equations for the interchain CDW and Josephson couplings, V and J respectively:

$$dV/d\tau = (2 - K)V \quad dJ/d\tau = (2 - 1/K)J , \quad (3)$$

where τ is the RG parameter. The “starting” values for the dimensionless couplings at the energy scale of the spin gap [9] are given by $V(\tau = 0) = J(\tau = 0) \sim (\delta t)^2 / \Delta_s^2$, where δt is the microscopic interladder hopping parameter (see Fig. 1). Eq. (3) shows that the CDW order parameter increases faster than SC for $K < 1$, and, thus, it is the first one to reach strong coupling. The question is whether there is a way to reverse this result and have superconductivity “win” the competition against CDW, by still keeping $K < 1$.

* Corresponding Author: Graz university of Technology, Petersgasse 16, A-8010 Graz, Austria. Phone +43-316-873-8180, Fax: +43-316-873-8677, Email: arrigoni@itp.tu-graz.ac.at




Fig. 1. Schematic representation of the coupled ladder system with alternating on-site energies discussed in the text. The ladders can be described, e. g., by a single-band Hubbard model or by a $t - J$ model. The ladders are then coupled via a hopping $\delta t \ll \Delta_s$, and an alternating on-site energy $\pm \varepsilon$ has been introduced in order to frustrate CDW pinning, as discussed in the text.

The solution consists in introducing a frustration in the CDW periodicity between the ladders. To understand this idea, one should observe that quasi-long-ranged CDW correlations in a single ladder become long-ranged in an array of coupled ladders because of the pinning between the CDW's on different ladders. Pinning can occur only if the wavelength $\pi/(2K_F)$ of the CDW is commensurate (equal) between neighboring ladders.

Thus, CDW pinning can be frustrated by introducing different Fermi momenta on the ladders. This is readily obtained, e. g., by introducing an alternating on-site energy offset ε [10], as shown in Fig. 1. Interestingly, the model shown in Fig. 1 displays the stripe structure which has been observed in several high- T_c superconductors in various regions of their phase diagram [11]. In particular, the period 4 structure seems to be the most stable stripe structure, at least in the $\text{La}_{1.6-x}\text{Nd}_{0.4}\text{Sr}_x\text{CuO}_4$ compound.

The energy offset ε introduces two different Fermi momenta K_{F1} and K_{F2} , with $K_{F1} - K_{F2} \sim \varepsilon/v_F$, which are associated to two potentially relevant CDW's with different wavelengths. The RG flux of the associated couplings V_1 and V_2 are both given by the first of Eq. (3) up to $\tau = \bar{\tau} \equiv \ln \Delta_s/\varepsilon$. For $\tau > \bar{\tau}$ the RG equation is given by (we take for simplicity the same K on the two different kinds of ladders)

$$dV_{1/2}d\tau = (1 - K/2)V_{1/2} \quad . \quad (4)$$

This form is due to the fact that the V_i operator ($i = 1, 2$) coupling CDW on neighboring ladders continues to increase only on the ladder with $K_F = K_{Fi}$, while it is frozen on the other.

Combining Eq. (3) with Eq. (4) it is easy to see that the condition for J to become more relevant than V , i. e. for superconductivity to win against CDW order is [10] (we consider here the case of a not too small ε)

$$K > K_c = \sqrt{3} - 1 \approx 0.8 \quad , \quad (5)$$

i. e. one has SC also for values of $K < 1$. Under this condition, the SC (Kosterlitz-Thouless [12]) critical temperature T_c is obtained by integration of Eq. (3) up to $J(\tau^*) \sim 1$. One obtains [9]

$$T_c \sim \Delta_s J_0^{K/(2K-1)} \sim \Delta_s \left(\frac{\delta t^2}{\Delta_s^2} \right)^{K/(2K-1)} \quad . \quad (6)$$

Notice that, in contrast to the BCS case in which T_c is exponentially small in the small coupling parameter (in that case given by the effective attractive interaction), here T_c is only power-law small in δt .

In conclusion, we have presented a model in which a relatively “high” T_c is obtained by a purely repulsive interaction between the electrons. The model consists of an array of weakly-coupled two-leg ladders with alternating on-site energies. This alternation is important in order to prevent the system to develop into a CDW insulator before becoming a superconductor.

This work was supported, in part, by NSF grants No. DMR 01-10329 at UCLA (SAK), and DMR-01-32990 at the University of Illinois (EF), and by a Heisenberg grant (AR 324/3-1) from the DFG (EA).

References

- [1] D. J. Scalapino, *Nature (London)* **377**, 12 (1995).
- [2] E. Dagotto and T. M. Rice, *Science* **271**, 618 (1996).
- [3] C. A. Hayward, D. Poilblanc, R. M. Noack, D. J. Scalapino, and W. Hanke, *Phys. Rev. Lett.* **75**, 926 (1995).
- [4] R. M. Noack, S. R. White, and D. J. Scalapino, *Phys. Rev. Lett.* **73**, 882 (1994).
- [5] C. A. Hayward and D. Poilblanc, *Phys. Rev. B* **53**, 11721 (1996).
- [6] M. Troyer, H. Tsunetsugu, and T. M. Rice, *Phys. Rev. B* **53**, 251 (1996).
- [7] R. M. Noack, S. R. White, and D. J. Scalapino, *Physica C* **270**, 281 (1996), (cond-mat/9601047).
- [8] R. M. Noack, N. Bulut, D. J. Scalapino, and M. G. Zacher, *Phys. Rev. B* **56**, 7162 (1997).
- [9] This holds away from special values of the doping, where Δ_s is of the order of the exchange coupling. Special dopings are discussed in Ref. [10].
- [10] E. Arrigoni, E. Fradkin, and S. A. Kivelson, cond-mat/0309572 (to appear in *Phys. Rev. B*).
- [11] For a brief review of the evidence of widespread local stripe order in the cuprates, see V. J. Emery, S. A. Kivelson, and J. M. Tranquada, *Proc. Natl. Acad. Sci.* **96**, 8814 (1999).
- [12] In two dimensions the order is long ranged only at $T = 0$, otherwise one obtains a quasi-long-ranged phase below the Kosterlitz-Thouless temperature.