

Program LM TART for Electronic Structure Calculations

S. Y. Savrasov

Department of Physics, New Jersey Institute of Technology, Newark, NJ 07102, USA

A computer program LM TART for electronic structure calculations using full potential linear muffin-tin orbital method is described.

PACS numbers:

I. DESCRIPTION

Full potential linear muffin-tin orbital (Ref.¹) (FP-LMTO) program LM TART is designed to perform band structure, total energy and force calculations of solids using methods of density functional theory (DFT) (Ref.²). The development of LM TART has been initiated in 1986 in P.N. Lebedev Physical Institute. It was further extensively developed in Max-Planck Institut für Festkörperforschung, and its most recent contributions have been done in Departments of Physics of Rutgers University and of New Jersey Institute of Technology.

LM TART performs electronic structure calculations using popular local density approximation (LDA) and generalized gradient approximation (GGA) of density functional theory². To deal with strongly correlated systems it also implements LDA+U method³ and will include the capability to study materials using spectral density functional approach and dynamical mean field theory (DMFT)⁴ in the nearest future. For determining equilibrium crystal structure and phonon spectra LM TART computes total energies, forces⁵ as well as full wavevector dependent lattice dynamics⁶ using linear response theory. For studying optical spectral functions LM TART evaluates dipole matrix elements between various electronic states and can determine optical conductivity of materials.

The code implements atom centered local muffin-tin orbitals as a basis for representing one-electron wave functions in both bare and screened (tight-binding) representations with general choice of tail energies². This allows to increase accuracy in the calculations by using multiple LMTO expansions. Low lying core electronic states can be resolved as bands in separate energy panels. For heavy elements, spin-orbit coupling matrix elements can be taken into account using a variational procedure proposed by Andersen¹. Full three dimensional treatment of magnetization in relativistic calculations including LDA+U is implemented to study problems related to magnetic anisotropy and noncollinear magnetism. Effect of finite temperature can be modelled by introducing Fermi statistics for the one-electron states. Additionally, LM TART can determine tight-binding fits to the energy bands by extracting hopping integrals between various orbitals using tight-binding LMTO representation.

LM TART is written on FORTRAN 90 and uses dynamical memory scheme. No additional recompilation of the code is required when changing numbers of atoms,

spins, plane waves, etc. There are two basic regimes for working with this program: the self-consistent charge density calculation and calculation of physical properties such as electronic structure, optical properties, etc.. The simplest input to LM TART involves only atomic charges of the atoms as well as crystal structure.

LM TART works with two different approximations related to a shape of the potential: (i) atomic sphere approximation (ASA) and plane wave expansion (PLW). ASA uses overlapping atomic spheres, where the potential is expanded in spherical harmonics inside the spheres, but any contribution from the interstitial region is neglected. Such method is fast and provides reasonably good energy bands. However it is not sufficiently accurate to deal with distortions and phonons. PLW is a full potential approximation which uses non-overlapping muffin-tin spheres, where the potential is represented via spherical harmonics expansions, and the interstitial region where the potential is expanded in plane waves. The full potential regime provides the best accuracy at the price of increasing computational time. Finally, LM TART can be run in a special tight-binding regime by supplying hopping integrals between various orbitals.

Two extensions of LM TART are available as separate programs. One is a full potential linear response linear muffin-tin orbital package LMTO PHONONS which is designed to perform linear response calculations of the phonon spectra for arbitrary wave vectors q . Main features of this code include: (i) Computations of first order changes in the charge density and the potential due to displacements of nuclei for arbitrary wave vectors q . (ii) Calculations of the dynamical matrix and the phonon spectra. (iii) Calculations of electron-phonon interactions, Eliashberg spectral functions² $F^2(\omega)$ and transport properties such as electrical and thermal resistivities. The description of the method and complete set of references can be found in Ref.⁶. Numerous applications are given in Ref.⁷.

Also available is an extension of LM TART to compute dynamical susceptibilities and magnon spectra. This is a full potential linear response linear muffin-tin orbital package of programs LMTO MAGNONS designed for these purposes. As a by-product, spin wave spectra and their lifetimes are accessible as peaks in calculated imaginary spin susceptibility. A short description of the method can be found in Ref.⁸.

Software MIND Lab which runs under Microsoft Windows operating systems can be used to set up input files

and analyze output files of LM TART. It can also be used to run the code without learning its extensive input options. This software is necessary to visualize all data withdrawn from LM TART. Currently, MINDLab can perform the following tasks (i) Crystal group calculations. By setting atomic positions MINDLab shows crystal group operations found for given atomic configuration; (ii) Fat bands calculations. Fully colored visualization of separated orbital characters on top of the band structure can be performed by mouse click operations. (iii) Density of states calculations. MINDLab computes and visualizes densities of states, total and orbital resolved. (iv) Extractions of hopping integrals for tight-binding fits can be performed quickly with MINDLab. (v) Optical properties calculations. Dielectric functions $\epsilon_1(\omega)$, $\epsilon_2(\omega)$, and electron energy loss spectra can be computed and visualized by MINDLab. (vi) Visualizations of charge densities, full potentials, and Fermi surfaces in both two and three dimensions using OpenGL graphical library implemented within MINDLab. (vii) Three dimensional visualization of crystal structures and their distortions. (ix) Full three dimensional visualization of vector fields including velocity fields on top of the Fermi surfaces and magnetizations. (x) Simplified input for studying correlated electronic systems which currently implements LDA+U method and will include LDA+DMFT technique in the nearest future.

MINDLab works with LM TART as with an executable file which reads input data, performs band structure calculation, and stores output files. MINDLab controls this process, it prepares input for LM TART using dialog windows, and executes LM TART program as a separate thread. When the calculation is finished, LM TART notifies the MINDLab, and the output files can be visualized by simple mouse click operations. While running LM TART for complicated compounds can be rather slow (it depends how many atoms per unit cell is selected), MINDLab can be used to prepare the input, after which the files can be copied to a more powerful computer where the LM TART can be executed much faster. After run is performed, the output can be copied back to the PC and analyzed quickly by MINDLab. All input/output files of LM TART are formatted ASCII files, and therefore system independent.

II. APPLICATIONS

Several most recent applications of LM TART can be found in Refs.^{4,7,8,9,10}. They include calculations of phonons and electron-phonon interactions in novel superconductors, such as MgB₂, LiBC, MgCNi₃; studies of phonons in strongly correlated systems, e.g. NiO, Pu, computations of magnetic anisotropy energies of ferromagnets Fe, Co, Ni, CrO₂ evaluations of dynamical spin susceptibilities and spin wave spectra, and so on.

Here we illustrate one of the most recent calculation⁹ of phonon spectrum in MgCNi₃ using a linear response

regime of LM TART. The calculated phonon dispersions along major high symmetry lines of the cubic Brillouin zone are given on Fig. 1. The frequencies are seen to be span up to 1000 K, with some of the modes showing significant dispersion. In general, we distinguished three panels where the top three branches around 900 K are carbon based, the middle three branches around 600 K are Mg based and 9 lower branches are all Ni based.




FIG. 1: Calculated phonon spectrum of MgCNi₃ along major symmetry directions of the Brillouin zone of the cubic lattice using density functional linear response method. Some curves are widened proportionally to the phonon linewidths.

We have discovered a striking feature of this phonon spectrum connected to the presence of a low frequency acoustic mode which is very soft and is even seen to be unstable along some points in (00) direction in the Brillouin zone. This mode is essentially Ni based and corresponds to perpendicular movements of two Ni atoms towards octahedral interstitials of the perovskite structure. The latter is made of each of the four Ni atoms and two Mg atoms.

The persistence of the instability which does not occur for any of the high symmetry point needs a non-trivial frozen phonon analysis. As our polarization vectors prompt that the largest anharmonicity is expected for the XM plane ($q_z = 0$) of the Brillouin zone of the cubic lattice, we have performed three such calculations for the points $a = (\frac{1}{4}\frac{1}{4}0)\frac{2\pi}{a}$, $b = (\frac{1}{2}\frac{1}{4}0)\frac{2\pi}{a}$, and M. The results of these calculations reveal essentially anharmonic interatomic potentials. A shallow double well with a depth of the order of 20-40 K and the curvature at the equilibrium of the order of 40-50 iK exists for the a and b points which becomes vanishing at the M point. We have found such a behavior by both total energy and force calculations for the supercells of 20 atoms (a and b points) and 10 atoms (M point). Clearly, such a small depth on the temperature scale indicates that the distortions are dynamical and zero point motions would prevent the

appearance of the static long range order. Complete discussion of these calculations as well as the comparisons with existing experiments can be found in Ref.⁹.

Fig.1 also shows the calculated phonon linewidths γ_q by widening some representative phonon dispersion curves ω_q proportionally to γ_q . In particular, we found that some phonons have rather large linewidths. This, for example holds, for all carbon based higher lying vibrational modes. The strength of the coupling, γ_q ; for each mode can be obtained by dividing γ_q by $N(E_F)^2$, where $N(E_F)$ is the density of states at the Fermi level equal to 5.3 st./[eV*cell] in our calculation. Due to large $N(E_F)^2$; this unfortunately results in strongly suppressed coupling for all carbon modes which would favor high critical temperatures. The coupling, however, is relatively strong for the Ni based modes. For example, we can find γ 's of the order of 1/3 for the Ni based optical phonons around 250 K. Our resulting value of electron-phonon coupling constant is found 1.51. This assumes that MgCNi₃ is a strongly coupled electron-phonon superconductor.

III. DOWNLOADS

All LM TART based programs described above are available for downloading at the following URL:

<http://www.physics.njit.edu/~m.indlab>. Each program consists of the source code, manual, and examples. An extensive database of electronic structures for different materials is also available at this HTTP URL.

IV. ACKNOWLEDGEMENTS

Part of the development has been done in collaboration with my brother Dr. Dmitriy Savrasov. Special thanks to Prof. Ole Andersen and Dr. Ove Jepsen who are my LM TO teachers. I also greatly acknowledge Dr. Andrej Postnikov who has initiated writing of all manuals.

The development of these programs is currently supported by the NSF DMR Grants No. 02382188, 0342290, US DOE division of Basic Energy Sciences Grant No. DE-FG 02-99ER 45761, and by the Computational Material Science network operated by US DOE.

-
- ¹ O. K. Andersen, Linear Methods of Band Theory, Phys. Rev. B 13, 3050 (1975).
 - ² For a review, see, Theory of the Inhomogeneous Electron Gas, edited by S. Lundqvist and S. H. March (Plenum, New York, 1983).
 - ³ For a review and complete set of references see, e.g., V. Anisimov, F. Aryasetiawan, and A. I. Lichtenstein, LDA+U Method, J. Phys.: Condens. Matter 9, 767 (1997).
 - ⁴ S. Savrasov, G. Kotliar, and E. Abrahams, Electronic correlations in metallic Plutonium within dynamical mean-field picture, Nature 410, 793 (2001); Xi Dai, S. Savrasov, G. Kotliar, A. M. Iglori, H. Ledbetter and E. Abrahams, Calculated Phonon Spectra of Plutonium at High Temperatures, Science 300, 953 (2003); S. Y. Savrasov and G. Kotliar, "Linear Response Calculations of Lattice Dynamics in Strongly Correlated Systems, Phys. Rev. Lett. 90, 056401 (2003); S. Y. Savrasov and G. Kotliar, Spectral Density Functional for Electronic Structure Calculations, Phys. Rev. B 69, 245101 (2004).
 - ⁵ S. Y. Savrasov and D. Y. Savrasov, Full potential linear-muffin-tin-orbital method for calculating total energies and forces, Phys. Rev. B. 46, 12181 (1992).
 - ⁶ S. Y. Savrasov, A linearized direct approach for calculating static response in solids, Solid State Commun. 74, 69 (1990); S. Y. Savrasov, Linear-response calculations of lattice dynamics using muffin-tin basis sets, Phys. Rev. Lett. 69, 2819 (1992); S. Y. Savrasov, Linear-response theory and lattice dynamics: a muffin-tin-orbital approach, Phys. Rev. B 54, 16470 (1996).
 - ⁷ S. Y. Savrasov, D. Y. Savrasov and O. K. Andersen, Linear-response calculations of electron-phonon interactions, Phys. Rev. Lett. 72, 372 (1994); S. Y. Savrasov and D. Y. Savrasov, Electron-phonon interactions and related physical properties of metals from linear-response theory, Phys. Rev. B 54, 16487 (1996); S. Y. Savrasov and O. K. Andersen, Linear-response calculation of the electron-phonon coupling in doped CaCuO₂, Phys. Rev. Lett. 77, 4430 (1996).
 - ⁸ S. Y. Savrasov, Linear Response Calculations of Spin Fluctuations, Phys. Rev. Lett. 81, 2570 (1998).
 - ⁹ A. Ignatov, S. Y. Savrasov, T. A. Tyson, Superconductivity near the vibrational mode instability in MgCNi₃, Phys. Rev. B 68, 220504(R) (2003);
 - ¹⁰ S. Ostanin, V. Trubitsin, J. B. Staunton, S. Y. Savrasov, A density functional study of pressure induced superconductivity in P and its implication for spintronics, Phys. Rev. Lett. 91, 087002 (2003); J. M. An, S. Y. Savrasov, H. Rosner, W. E. Pickett, Extreme Electron-Phonon Coupling in Boron Based Layered Superconductors, Phys. Rev. B 66, 220502(R) (2002); I. Yang, S. Savrasov, and G. Kotliar, Importance of correlation effects on magnetic anisotropy in Fe and Ni, Phys. Rev. Lett. 87, 216405 (2001);