

Ground-State Degeneracy of Correlated Insulators with Edges

Gil Refael¹ and Hsiu-Hau Lin^{1,2,3}

¹*Kavli Institute of Theoretical Physics, University of California, Santa Barbara, CA 93106*

²*Department of Physics, National Tsing-Hua University, Hsinchu 300, Taiwan*

³*Physics Division, National Center for Theoretical Sciences, Hsinchu 300, Taiwan*

(Dated: March 23, 2022)

Using the topological flux insertion procedure, the ground-state degeneracy of an insulator on a periodic lattice with filling factor $\nu = p/q$ was found to be at least q -fold. Applying the same argument in a lattice with edges, we show that the degeneracy is modified by the additional edge density ν_E associated with the open boundaries. In particular, we demonstrate that these edge corrections may even make an insulator with integer bulk filling degenerate.

A fascinating aspect of quantum mechanics is the interplay between global topology and some physical properties, that are naively believed to depend only on local quantities. Restrictions that originate from *topological* arguments[1] often lead to elegant and non-trivial predictions of bulk properties in a non-perturbative fashion. One example is particle statistics dictating the possible ground states of a system. A stark demonstration of that is anyons with fractional statistics[2]. Another instance of this picture is the non-perturbative proof of Luttinger's theorem[3, 4]; it is revealed that the topology dictates the space enclosed within the Fermi surface of a Landau-Fermi liquid to be exactly the particle density in the conduction band, disregarding the details of the microscopic Hamiltonian. Beyond their elegance and appeal, topological ground-state properties have been recently made the centerpiece, primarily due to Kitaev, of a novel kind of quantum computation[5].

Predictions from topological arguments are often sensitive to the geometries of the bulk. For instance, the insulating Z_2 -gauge spin liquid[6] is four-fold or two-fold degenerate depending on the bulk geometry is a torus or a cylinder respectively. Another well-known example is the $\nu = 1/q$ fractional-quantum-Hall liquid[7] on the two-dimensional torus. A topological constraint due to these exotic excitations gives rise to the non-trivial q -fold degeneracy. Remarkably, by cutting the torus into a closed strip with two open edges, the ground-state degeneracy in the thermodynamic limit shoots up from q to infinity, with gapless excitations described by the chiral Luttinger liquid on either side of the Hall bar. It is then natural to expect that the topological arguments might deliver rather different messages when edges are present. Indeed, the quantum Hall system is not the only example of juicy edge physics. Other examples are Andreev bound states appearing on edges of superconducting lattices[8], spin-1/2 excitations of the Haldane phase located at the edge of $S = 1$ Heisenberg chain[9], and the ferromagnetic moment on a zigzag tip of a carbon nanotube[10].

In this Letter, we revisit a topological constraint in insulators - the correspondence between the filling factor and the ground state degeneracy. In

Ref. [11], Oshikawa extends the Lieb-Schultz-Mattis argument[12] and Laughlin's treatment of the quantized Hall conductance[13], to show that the ground state degeneracy of an insulator with filling fraction $\nu = p/q$ (where q and p are coprimes) is at least q -fold. This is established using the flux-insertion method applied to the system on the periodic and finite lattice. By taking appropriate thermodynamic limit, the ground state degeneracy is related to the intensive particle filling factor in the bulk. However, in the presence of open boundaries, the particle density is no longer uniform near the edges. The physics of the edge is captured through appropriate edge filling factors that are introduced in addition to the bulk density. In this Letter, we will show that the degeneracy of the ground state is determined by *both the bulk and edge filling factors* in the presence of edges.

For instance, a gapped insulator on a periodic lattice, made of spinless particles with bulk filling factor $\nu_B = 1$, may have no degeneracy. However, if the lattice has open boundary conditions along one direction, the particle density near the edge may differ from the bulk filling factor far from the edge. This difference is captured by the edge density ν_E , which arises from particles localized near the edge, as well as changes of the extended wave functions in the bulk. The edge state may break translational invariance and gives rise to ground-state degeneracy despite of the bulk state having perfect translational symmetry parallel to the edge. For example, for $\nu_E = 1/2$, the edge may form a CDW with period two. Such a state is doubly degenerate, although the bulk insulator is featureless.

Let us now review the flux insertion procedure in the context of a system with open boundaries in one direction (we follow Ref.[11] closely). Consider a lattice in d -dimensional space with open boundaries in one direction, $1 \leq x_1 \leq L_1$, and periodic in the remaining $d-1$ dimensions with finite length L_i , $i = 2, \dots, d$. The insulating ground state $|\Phi_0\rangle$ contains N particles that reside on the lattice sites. Making use of the periodicity of the lattice, say in the x_d direction, the ground state can be chosen to be an eigenstate of the momentum operator \hat{P}_d , carrying momentum P_d^0 in the x_d direction.

We now introduce a fictitious vector potential \vec{A} that couples with “unit charge” to all particles in the system. Protected by the gap, one can adiabatically insert a flux quanta $\Phi = hc$ through the ring in the x_d -direction and map $|\Psi_0\rangle$ to $|\Psi'_0\rangle$ of the same energy. Because the flux insertion can be achieved by the constant vector potential $A_d = \Phi/L_d$ in the direction of x_d , which commutes with \hat{P}_d , the momentum remains constant in the whole adiabatic procedure, $\hat{P}_d|\Psi'_0\rangle = P_d^0|\Psi'_0\rangle$.

But the flux insertion changes the Hamiltonian from $\mathcal{H}(\Phi = 0)$ to a different topological sector $\mathcal{H}(\Phi = hc)$. Before being able to compare $|\Psi'_0\rangle$ to $|\Psi_0\rangle$, we need to restore the Hamiltonian to the same topological sector $\mathcal{H}(\Phi = 0)$. This is done using the following large unitary gauge transformation:

$$U = \exp\left(\frac{2\pi i}{L_d} \sum_{\vec{x}} x_d \hat{n}_{\vec{x}}\right). \quad (1)$$

Now $U\mathcal{H}(hc)U^{-1} = \mathcal{H}(0)$, and therefore $U|\Psi'_0\rangle$ is an eigenket of the original Hamiltonian. Since our system is a gapped insulator, and the flux insertion was adiabatic, we expect that $U|\Psi'_0\rangle$ has the same energy as $|\Psi_0\rangle$ (with $\mathcal{H}(0)$ in mind).

The momentum of the newfound ground state can be evaluated straightforwardly, $\hat{P}_d U|\Psi'_0\rangle = (U\hat{P}_d + [\hat{P}_d, U])|\Psi'_0\rangle = (P_d^0 + 2\pi N/L_d)U|\Psi'_0\rangle$. So the momentum of $U|\Psi'_0\rangle$ is $P'_d = P_d^0 + 2\pi N/L_d$. If N and L_d are mutually prime, then $U|\Psi'_0\rangle$ and $|\Psi_0\rangle$ are *two* degenerate ground states. To relate the ground state degeneracy to the filling factor, one needs to take appropriate thermodynamic limit. In a fully periodic system, Oshikawa's argument follows by saying that

$$\Delta P = P'_d - P_d^0 = 2\pi \frac{N}{L_d} = 2\pi \nu C_d, \quad (2)$$

where $\nu = N/V = p/q$ is the filling factor of the lattice. The volume of the system is $V = \prod_{i=1}^d L_i$ and the transverse size at each particular x_i is $C_i = V/L_i$. The thermodynamic limit is taken by fixing ν (so that p and q are well defined without any correction) and choosing C_d to be coprime with q . Thus, by repeating the flux insertion, we can generate q -fold distinct ground states. If we assume that the ground state degeneracy D_g is a robust quantity, disregarding the details of the thermodynamic limit, the gauge argument leads to the constraint $D_g \geq q$.

Returning to the non-periodic system introduced above, we pick up the discussion from Eq. (2). Two different kinds of non-uniformity occur near the edges. First, the edges give rise to new states of particles localized near the edges; we define the *edge-state filling factor* $n_e = N_e/C_1$, where N_e is the number of edge particles. Second, bulk states (forming a continuum) that propagate throughout the lattice may be repelled from the edges. This repulsion creates a charge depletion near the edge; we define $n_\Delta = N_\Delta/C_1$, where N_Δ is the total

charge depleted near the edge. Both effects give rise to non-trivial corrections to the gauge argument.

Loosely speaking, the presence of edges divides the total number of particles into two groups: bulk and edge, labeled by N_B and N_E respectively. The precise distinction between bulk and edge states would be explained in later paragraphs. In light of this division, we need to restate Eq. (2) as

$$\Delta P = 2\pi \frac{N_B + N_E}{L_d} = 2\pi \frac{N_B}{L_d} + 2\pi n_e C_{1d}, \quad (3)$$

with $C_{1d} = V/(L_1 L_d)$. Before we can conclude anything about the ground-state degeneracy, we also need to account for the repulsion of bulk states from the edge.

As mentioned above, the edges may repel the bulk states, creating a charge depletion near the edges with density $n_\Delta = N_\Delta/C_1$. Now, in order to achieve the thermodynamic limit of an insulator with bulk filling ν_B and edge filling n_e , we need to consider a sequence of finite lattices with a total number of particles which is

$$\begin{aligned} N_0 &= N_B + N_E = (\nu_B V - N_\Delta) + n_e C_1 \\ &= \nu_B V + (n_e - n_\Delta) C_1 \end{aligned} \quad (4)$$

Note that $n_\Delta \equiv p_\Delta/q_\Delta$ is not necessarily an integer. To contrast, in a periodic system the thermodynamic limit would simply be taken by considering a sequence of lattices with $N_0 = \nu_B V$ particles.

Including both types of edge effects, the momentum difference between $|\Psi_0\rangle$ and its sibling $U|\Psi'_0\rangle$ is

$$\Delta P = 2\pi \nu_B C_d + 2\pi \nu_E C_{1d}, \quad (5)$$

with $\nu_B = p_B/q_B$ and $\nu_E = n_e - n_\Delta = p_E/q_E$. From the revised gauge argument in Eq. (5), we obtain the main result of this paper: the degeneracy of an insulating state in the presence of edges is given by

$$D_g \geq LCD(q_B, q_E) \geq q_B, \quad (6)$$

where LCD denotes *least common denominator*.

Before discussing possible extensions and generalizations of the result in Eq. (6), we resolve some subtleties in the derivation above. A central issue is the difficulty of distinguishing edge and bulk particles as in Eq. (3) in a strongly correlated system, where the concept of single particle states is rather vague. At first sight, this distinction may seem impossible to carry out in practice, and might be mistaken to be a theorist's whim; there is, however, a precise way of counting the number of edge and bulk states. This method utilizes a spectral function decomposition of the system's correlation function, which in principle we know with utmost precision. Following Lehman decomposition of the single-particle spectral function:

$$\begin{aligned} A(\vec{x}, \omega) &= \sum_m \delta(\omega - E_m) |\langle m | \psi^\dagger(\vec{x}) | \Psi_0 \rangle|^2 \\ &\pm \sum_m \delta(\omega + E_m) |\langle m | \psi(\vec{x}) | \Psi_0 \rangle|^2. \end{aligned} \quad (7)$$

Here $\psi(\vec{x})$ is the annihilation operator in the Schrödinger picture at lattice site \vec{x} . The \pm distinguishes between fermions and bosons. Without loss of generality, let us concentrate on fermions. Note that $|m\rangle$ are many-body excited states that show one particle excitations. The excitations correspond to one-particle states of free particles. In the following we simply refer to $|m\rangle$ as states.

Next, consider the $\omega < 0$ (hole) part of the spectral function $A(\vec{x}, \omega)$. In a finite system, it is given as a discrete sum of δ -functions multiplied by matrix elements, and it obeys the sum rule

$$\sum_{\vec{x}} \int_{-\infty}^0 A(\vec{x}, \omega) d\omega = N, \quad (8)$$

where N is the total number of particles in the lattice. If we take the limit $L_1 \rightarrow \infty$ while keeping the bulk density fixed, we expect that some of the δ -functions will merge into continuum, whereas the rest will remain sharp and separated. The first group is the excited *bulk states*, and the second group is the excited *edge states*. We stress that $|m\rangle$ are strongly correlated many-body states that show one particle excitations $\Delta Q = \pm 1$ relative to the ground state. More precisely, bulk states $|m_B\rangle$ scale as

$$|\langle m_B | \psi^\dagger(\vec{x}) | \Psi_0 \rangle|^2 \sim \mathcal{O}\left(\frac{1}{L_1}\right) \quad (9)$$

for any \vec{x} , as L_1 is scaled to infinity. Unlike bulk states, the edge states, $|m_E\rangle$, tend to an \vec{x} -dependent constant as $L_1 \rightarrow \infty$:

$$|\langle m_E | \psi^\dagger(\vec{x}) | \Psi_0 \rangle|^2 \sim \mathcal{O}(1). \quad (10)$$

Thus we can break the spectral function into a bulk (A_B) and edge (A_E) parts:

$$\begin{aligned} A_{B/E}(\vec{x}, \omega) &= \sum_{m_{B/E}} \delta(\omega - E_{m_{B/E}}) |\langle m_{B/E} | \psi^\dagger(\vec{x}) | \Psi_0 \rangle|^2 \\ &\pm \sum_{m_{B/E}} \delta(\omega + E_{m_{B/E}}) |\langle m_{B/E} | \psi(\vec{x}) | \Psi_0 \rangle|^2. \end{aligned} \quad (11)$$

The definition of N_B and N_E is hence

$$N_{B(E)} = \sum_{\vec{x}} \int_{-\infty}^0 d\omega A_{B(E)}(\vec{x}, \omega). \quad (12)$$

There are two subtleties in the above procedure. First, it is possible that an edge state hybridizes with a bulk state. Definition along the lines of Eqs. (11) and (12) specifies that such a state is an edge state; even though such a state is partially delocalized, it is associated with the edge due to the localized weight. The number of such states will scale with L_i for $i > 2$, but *not* with L_1 . Another subtlety occurs if there is an accidental

degeneracy between a bulk and edge states. In this case, in a finite system, it may be impossible to distinguish between the edge and bulk states and two hybridized states. This accidental degeneracy will surely be lifted by a different choice of system dimensions, and therefore needs not bother us.

Now that the derivation is complete, we would like to elaborate on some extensions of the edge argument. The first question we address is the effect of bulk-state depletion near the edge. Note that if n_Δ is an integer, the bound on the ground-state degeneracy, $D_g \geq LCD(q_B, q_e)$, only depends on ν_B and $n_e = p_e/q_e$. In this case, the depletion of bulk states does not play any role in determining ground-state degeneracy. In fact, for featureless insulators with low-energy excitations described by Fermi liquid theory, we can show that n_Δ is an integer.

Start with the ground state $|0\rangle$ of the featureless insulator on the periodic lattice (without edges). The insulator with open edge can be viewed as perturbing the original ground state $|0\rangle$ with hole excitations near the boundary. The translational invariance ensures that we can construct local quasi-hole states $|r\rangle = \psi(r)|0\rangle$, with the same spatial profile but located at different C_1 lattice sites on the edge. In general, these states, located at different lattice sites, would have non-vanishing overlaps. If one assumes the Fermi-liquid picture is at work here, these local (Wannier) orbitals would form a continuous band (with dispersion needs to be determined self-consistently). To ensure the new ground state (in the presence of open boundaries) is also insulating, these quasi-particles must form a *band insulator* at the edge, i.e. the number of holes in one unit cell, $n_\Delta = N_\Delta/C_1$, is an integer.

The argument also applies to a simple but exotic situation when the low-energy excitations carry fractional “charges” (not necessarily the ordinary electric charge), but are well described by Fermi-liquid like theory. Following similar argument, n_Δ is fractional, reflecting the breaking up of ordinary particles. While the argument does not hold when low-energy excitations are no longer described by Fermi liquid theory, in the above case, n_Δ serves as a useful precursor for identifying low-energy excitations in featureless insulators.

Another extension of the theorem is when the edges are far apart. It is then natural to divide the edge corrections, $n_e = \sum_i n_{ei}$ and $n_\Delta = \sum_i n_{\Delta i}$, associating with each edges. A simple example helps to demonstrate this point and the statements above; consider a segment of carbon nanotube with circumference L_y (in y direction) and zigzag edge at $x = 0$ and $x = L_x$. Choose the tight-binding hopping to be in the strong anisotropic regime, $|t_h| > 2|t|$, where t_h and t denote the hopping amplitudes along horizontal and (vertical) zigzag bonds. The interactions between particles are assumed to be weak, with a typical energy scale V .

Ignoring the weak interaction momentarily, the open

boundaries at $x = 0$ and $x = L_x$, give rise to single-particle edge states at $E = 0$ with momentum-dependent localization length $\xi(k_y) = \ln |(2t \cos k_y)/t_h|$. Each edge has L_y distinct states, corresponding to each quantized k_y . Since the bare hopping Hamiltonian is quadratic, the total number of states has to equal the number of particles in the lattice, or twice the number of unit cells (each carbon-nanotube unit cell contains two sites). Hence, the L_y edge states must come at the expense of L_y bulk states. This leads to $n_\Delta = 1$ at natural filling $\nu_B = 1$, and thus there is no effect on the degeneracy due to the bulk states repulsion at the edge.

Nevertheless, the edge-state filling factor, $n_e = n_{e1} + n_{e2}$, may be tuned to arbitrary fractions by removing or adding a small amount of electrons (proportional to the length L_y) which do not affect ν_B . Because the interaction V is weak (compared with the bulk gap Δ_B), the non-trivial mixing of single-particle state occurs within edge states and lifts the exact degeneracy at $E = 0$. The ground state is non-trivial and certainly depends on the specific form of interactions. But the revised gauge argument with the momentum shift in Eq. (5), leads to at least q_e -fold degeneracy of the ground state even though the bulk filling is an integer.

When considering particular physical systems, as in the above example, it is possible that the ground-state degeneracy will be larger than our rigorous result in Eq. (6). If the two edges are known by other means to be independent (as is probably the case in many large systems) and are also independent of the bulk state, then the degeneracy should be at least the product $q_{e1} \cdot q_{e2} \cdot LCD(q_\Delta, q_B)$. This degeneracy would become the number of low-lying states, if there is a weak interaction between the two edges. Other cases are also possible, most notably, independence of the edge states from the bulk, but not from each other. In this case the degeneracy is $LCD(q_{e1}, q_{e2}) \cdot LCD(q_\Delta, q_B)$. We add that for incommensurate edge filling in the thermodynamic limit $L \rightarrow \infty$, it is very likely that the low-energy physics of the system is described by a gapless liquid on the edge, living inside the gapped bulk spectrum (similar to the chiral edge states in quantum Hall liquid). Since these edge excitations sometimes are the only low-energy excitations, appropriate treatment of them is crucially important.

An interesting situation that may also be quite common is that the bulk and edge conspire to produce a uniform particle density even near the edge. This happens if the bulk states repelled from the edge are exactly compensated by particles trapped at the edge: $n_e = n_\Delta$. Our revised gauge argument is then reduced to Oshikawa's original argument with only the bulk filling determining the degeneracy.

The revised gauge argument can be generalized to the case of more than one non-periodic direction. We require

that at least one of the directions of the lattice is periodic to derive a generalized gauge argument, although this restriction may not be necessary in a physical system. We consider x_d as periodic. If all other sides of the attice terminate at $x_i = 0$ and $x_i = L$, there may be edge (surface) and wedge states, with filling numbers that scale as $N_\alpha \sim L^\alpha$, where $\alpha = 1, 2, \dots, d-1$. One then needs more filling factors n_α to specify the filling of the system. Also, one needs the depletion parameters $n_{\Delta_a}^b$ where $a = 1, \dots, d$ specifies the repelled 'bulk' state, and $b = 1, \dots, a-1$ is the number of restricted dimensions for the bulk states. To clarify this statement consider the example of a cube that is made periodic in the z direction. The bulk is three dimensional, and the depletion away from the surfaces ($x = 0, x = L, y = 0$ and $y = L$) amounts to $n_{\Delta_3}^1 \cdot L^2$ particles. Also, the wedges at $x = 0, y = 0, x = 0, y = L$ etc. may repel the bulk states with a volume of $n_{\Delta_3}^2 \cdot L$. Similarly, edge (surface) states on the faces of the cube may be repelled from the wedges with corrections $n_{\Delta_2}^1 \cdot L$.

In summary, using topological flux insertion procedure, we rederive the connection between the ground-state degeneracy in an insulator with its filling factors ν_B in the bulk and ν_E at the edge. As expected, the presence of open edges produces rich physical phenomena at the boundaries. In addition, the response of the bulk states to the edge may lend a different perspective into the correlated bulk physics.

We would like to thank Leon Balents, Matthew Fisher, Arun Paramekanti, Tami Pereg-Barnea, and particularly Masaki Oshikawa for insightful discussions. HHL is grateful for support from NSC-91-2120-M-007-001 and NSC-92-2112-M-007-039. GR thankfully acknowledges support from NSF fund PHY99-07949.

-
- [1] X. G. Wen, cond-mat/9506066.
 - [2] Y.-H. Chen, F. Wilczek, E. Witten, and B. I. Halperin, Int. J. of Mod. Phys. B, **3**, 1001 (1989).
 - [3] M. Oshikawa, Phys. Rev. Lett. **84**, 3370 (2000).
 - [4] A. Paramekanti and A. Vishwanath, cond-mat/0406619.
 - [5] A. Yu. Kitaev, Annals Phys. **303**, 2 (2003).
 - [6] T. Senthil and M.P.A. Fisher, Phys. Rev. B **62**, 7850 (2000).
 - [7] W.-G. Wen, Int. J. Mod. Phys. B **6**, 1711 (1992).
 - [8] C.-R. Hu, Phys. Rev. Lett. **72**, 1526 (1994).
 - [9] P. P. Mitra, B. I. Halperin, and I. Affleck, Phys. Rev. B, **45**, 5299 (1992).
 - [10] G. Refael, H.-H. Lin and M. Oshikawa, work in progress.
 - [11] M. Oshikawa, Phys. Rev. Lett. **84**, 1535 (2000).
 - [12] E. H. Lieb, T. Schultz and D. J. Mattis, Ann. Phys. (N.Y.) **16**, 407 (1961).
 - [13] R. B. Laughlin, Phys. Rev. B **23**, 5632 (1981).