

Multiple quantum dynamics in dipole-coupling spins in solid.

G.B. Furman, S.D. Goren, V. M. Meerovich, V. L. Sokolovsky, and S. William

Physics Department, Ben Gurion University, Beer Sheva, 84105, Israel

Abstract

A perturbation method deals with dipolar coupling spins in solids is presented. As example of application the method, the multile-quantum coherence dynamics in clusters of a linear chain of four nuclear spins and a ring of six spins coupled by dipole-dipole interaction are considered. The calculated 0Q- and 2Q intensities in a linear chain of four nuclear spins and 6Q intensity in a ring of six spins vs the duration of the preparation period agree well with the exact solutions (for linear chain of four nuclear spins) and simulation data (for linear chain of four nuclear spins and a ring of six spin).

Introduction

The dipole-dipole interaction (DDI) is widespread in nature and determines behavior of many physical systems. In solids state nuclear magnetic resonance (NMR), it is responsible for many specific phenomena and plays an important role in nuclear spin dynamics [1]. However in solids the evolution of a spin system under the DDI involves many spins and behavior of such system cannot be analyzed analytically. Even the numerical analysis becomes difficult because the number of states $N = 2^n$ is growing exponentially with the increasing of number of spins n . Therefore existing theories are based on phenomenological models and only macroscopic characteristics such as spin-spin relaxation times, the second and the fourth moments of resonance lines are taken into account [1]. These difficulties are very clearly displayed in multiple-quantum (MQ) spin dynamics. The MQ phenomena involve various multiple-spin transitions between the Zeeman energy levels and MQ coherence is formed at time $t > \omega_d^{-1}$, where ω_d is the characteristic frequency of DDI [9]. Hence, $\omega_d t > 1$ is not a small parameter, that did not allow one to use perturbation methods to study MQ dynamics. Indeed only simple exactly solvable models of a spin system such as two and three dipolar coupling spins 1/2 [11, 12] or one-dimensional linear chains of spins [13] were analyzed theoretically. The last achievement in this direction is the model with identical DDI coupling constants for all spin pairs [14, 15]. Note that the simplified calculations essential for the case of identical DDI coupling constant have been already mentioned [17]. This model describes only zero- and second-order coherences. But in real solids, DDI is characterized by the different coupling constants. In order to obtain important information on molecular structure and spin dynamics of these systems, the evolution of large number modes of MQ coherence has to be analyzed.

Another important purpose of the study of MQ coherence is the field of quantum information processing [6, 7, 8]. It was experimentally demonstrated the possibility of creating pseudopure spin states in clusters of coupled spins by MQ method [18] and that dynamics of the quantum entanglement is uniquely determined by the time evolution of MQ coherences [12].

Our objective is to adapt a perturbation approach to the problem of dipolar coupling spin dynamics in solids and to obtain the description of large number modes of MQ coherence. The Hamiltonian of the spin system with different DDI constants is divided into several parts, each of them is characterized by its constant. Our main idea is to take into

account in MQ dynamics the influence of the different items of the Hamiltonian with different degree of accuracy. The items corresponding to smaller DDI coupling constants are considered as a perturbation while the item with the largest constant (describing the interaction between nearest neighbours) is taken into account exactly. The proposed approach will be a power method to describe wide range of physical problems dealing with dynamics of dipolar coupling spins in solids. On the one hand, this approach uses the advantages of exactly solvable models [15, 16]. On the other hand, it simplifies calculations by using a perturbation technique.

Theory

Let us consider a system of nuclear spins $1/2$ placed in an external magnetic field $H_0 \parallel z$ axis and subjected to a suitable designed pulse sequence, thus permitting the observation of MQ coherences [10]. The effect of the sequence of irradiating pulses on the spin system can be represented by a unitary transformation propagator [10]

$$U(t) = e^{-it\mathcal{H}}, \quad (1)$$

where $\mathcal{H}$ is the effective time-independent Hamiltonian. First assume that the effective Hamiltonian can be divided into two parts with only two different DDI constants α and β : $\mathcal{H} = \mathcal{A} + \mathcal{B}$, where the operators $\mathcal{A}$ and $\mathcal{B}$ can be presented in the following form: $\mathcal{A} = \alpha A$ and $\mathcal{B} = \beta B$, $\alpha = \|\mathcal{A}\|$ and $\beta = \|\mathcal{B}\|$ are norms of operators $\mathcal{A}$ and $\mathcal{B}$, respectively ($\alpha > \beta$ and $[A, B] \neq 0$). The operators A and B do not include any DDI coupling constants. We present expression (1) in the form

$$e^{-it(\alpha A + \beta B)} = e^{-it\beta B} \sigma_A(t), \quad (2)$$

where operator $\sigma(t)$ obeys the differential equation

$$i \frac{d\sigma(t)}{dt} = \alpha A(t) \sigma_A(t), \quad (3)$$

with the initial condition

$$\sigma_A(0) = 1 \quad (4)$$

and $A(t) = e^{-it\beta B} A e^{it\beta B}$. Assume that $\alpha > \beta$. Solving (3), we limit ourselves by the first power of $\frac{\beta}{\alpha}$. Expanding the exponential term in the right hand of Eq. (2) in a series and

substituting in Eq. (3), we obtain

$$i \frac{d\sigma_A(t)}{dt} = \alpha (A - it\beta [B, A]) \sigma_A(t). \quad (5)$$

To solve Eq.(5) we will use the iterative method. We will search the solution of Eq.(5) as a series on parameter $(\alpha t)^n$:

$$\sigma_A(t) = \sum_{n=0}^{\infty} \sigma_A^{(n)}(t) \quad (6)$$

Taking into account that $\sigma_A^{(0)}(t) = 1$ the solution of Eq.(5) for $n = 1$ can be obtained: $\sigma_A^{(1)}(t) = (yA - \frac{1}{2}y^2\varepsilon [B, A])$

$$\sigma_A^{(1)}(t) = (-i\alpha t) A. \quad (7)$$

For $n = 2$, ones obtains

$$\sigma_A^{(2)}(t) = \frac{(-i\alpha t)^2}{2} \left(A^2 + \frac{\beta}{\alpha} [B, A] \right) \quad (8)$$

Keeping only linear in $\frac{\beta}{\alpha}$ terms the following expression for operator $\sigma_A^{(m)}(t)$ can be obtained

$$\sigma_A^{(m)}(y) = \frac{(-i\alpha t)^m}{m!} \left(A^m - \frac{\beta}{\alpha} \left((m-1) B A^{m-1} - \sum_{j=0}^{m-2} A^{m-1-j} B A^j \right) \right) \quad (9)$$

Utilizing (9) to expand the expression of the unitary transformation propagator (1) we find

$$U(t) = e^{-it(\alpha A + \beta B)} = \left(1 - \frac{\beta}{\alpha} \sum_{n=0}^{\infty} \frac{(i\alpha t)^{n+1}}{(n+1)!} \sum_{j=0}^n \frac{(-1)^j}{j!(n-j)!} A^j B A^{n-j} \right) e^{-i\alpha t A}. \quad (10)$$

Taking into account that

$$\sum_{j=0}^n \frac{(-1)^j}{j!(n-j)!} A^j B A^{n-j} = \frac{1}{n!} \{B, A^n\} \quad (11)$$

where the repeated commutators bracket $\{\dots\}$ is defined by

$$\{B, A^0\} = B, \quad \{B, A^{n+1}\} = [\{B, A^n\}, A], \quad (12)$$

we obtained that expression of the unitary transformation propagator takes a form

$$U(t) = e^{-it(\alpha A + \beta B)} = \left(1 - \frac{\beta}{\alpha} \sum_{n=0}^{\infty} \frac{(i\alpha t)^{n+1}}{(n+1)!} \{B, A^n\} \right) e^{-i\alpha t A}. \quad (13)$$

The summing over n in (13) results in

$$\sum_{n=0}^{\infty} \frac{(i\alpha t)^{n+1}}{(n+1)!} \{B, A^n\} = i\alpha \int dt e^{i\alpha t A} B e^{-i\alpha t A}. \quad (14)$$

Substituting Eq. (14) into (13) we obtain

$$e^{-it(\alpha A + \beta B)} = \left(1 - i\beta \int_0^t dx e^{-ixA} B e^{ixA}\right) e^{-i\alpha t A}, \quad (15)$$

which is a well-known formula for expansion of an exponential operator in a perturbation series [19, 20]. Expanding the operator exponent $e^{-i\alpha t A}$ of the left-hand of Eq. (13) gives

$$e^{-it(\alpha A + \beta B)} = \sum_{k=0}^{\infty} \frac{(-i\alpha t)^k}{k!} \left(A^k - \frac{\beta}{\alpha} \sum_{n=0}^{\infty} \frac{(i\alpha t)^{n+1}}{(n+1)!} \{B, A^n\} A^k \right) \quad (16)$$

After rearranging of the series (16) we obtain

$$e^{-it(\alpha A + \beta B)} = \left[1 - \frac{\beta}{\alpha} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(-1)^m (i\alpha t)^{n+m+1}}{n!m!(n+m+1)} A^n B A^m \right] e^{-i\alpha t A} \quad (17)$$

In the same way we obtain the expansion up to second order in the ratio $\frac{\beta}{\alpha}$:

$$\begin{aligned} e^{-it(\alpha A + \beta B)} = & \left\{ 1 - \frac{\beta}{\alpha} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \left[\frac{(-1)^m (i\alpha t)^{n+m+1}}{n!m!(n+m+1)} A^n B A^m \right. \right. \\ & \left. \left. - \left(\frac{\beta}{\alpha} \right) \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} \frac{(-1)^{m+k} (i\alpha t)^{k+l+m+n+2}}{n!m!k!l!(k+l+1)(k+l+m+n+2)} A^{l+m} B A^k B A^l \right] \right\} e^{-i\alpha t A} \end{aligned} \quad (18)$$

Expression (18) can be generalized to a case when the exponential operator contains arbitrary number of the non-commutative operators and can be extended to include various powers of the operators. For example, if the effective Hamiltonian operator includes three non-commutative operators $\mathcal{H} = \mathcal{A} + \mathcal{B} + \mathcal{C} = \alpha A + \beta B + \delta C$, where $\delta \leq \beta < \alpha$, we can obtain the expansion in series for a three-spin cluster replacing B by $B + \frac{\delta}{\beta} C$ in Eq. (18) .

Expressions (17) and (18) represent independent expansions in two parameters $\frac{\beta}{\alpha} < 1$ and $x = \alpha t$. Eqs. (17) and (18) appear to be complicated but in fact it is quite simple to use, as the following examples will illustrate.

Results and discussion

Let us consider a cluster which constitutes a linear chain of four nuclear spins 1/2 coupled by DDI. The MQ dynamics in the rotating frame is described by the time-independent

average Hamiltonian:

$$\mathcal{H} = \sum_{j < k} \mathcal{H}_{jk} = -\frac{1}{2} \sum_{j < k} d_{jk} (I_j^+ I_k^+ + I_j^- I_k^-) \quad (19)$$

and I_j^+ and I_j^- are the raising and lowering operators for spin j . The dipolar coupling constant, d_{jk} , for any pair of nuclei j and k , is given by

$$d_{jk} = \frac{\gamma^2 \hbar}{2r_{jk}^3} (1 - 3 \cos^2 \theta_{jk}), \quad (20)$$

where γ is the gyromagnetic ratio of the nuclei, r_{jk} is the internuclear spacing, and θ_{jk} is the angle that the vector $\vec{r}_{jk}$ makes with the external magnetic field. In the high-temperature approximation the density matrix at the end of the preparation period is given by

$$\rho(t) = e^{-i\mathcal{H}t} \rho(0) e^{i\mathcal{H}t} \quad (21)$$

where $\rho(0)$ is the initial density matrix in the high-temperature approximation

$$\rho(0) = \sum_{j=1}^4 I_j^z, \quad (22)$$

I_j^z is the projection of the angular momentum operator of spin j on the direction of the external field. The average Hamiltonian (19) can be divided into the three parts according to the number of different coupling constants $d_{12} > d_{13} > d_{14}$:

$$\mathcal{H} = \mathcal{H}_{12} + \mathcal{H}_{13} + \mathcal{H}_{14}, \quad (23)$$

where d_{12} is the coupling constant of the nearest neighbors, d_{13} is the coupling constant of the next nearest neighbors, and d_{14} is the coupling constant of the next-next-nearest neighbors. The experimentally observed values are the intensities of multiple-quantum coherences $J_{nQ}(t)$:

$$J_{nQ}(t) = \frac{1}{Tr \rho^2(0)} \sum_{p,q} \rho_{pq}^2(t) \text{ for } n = m_{zp} - m_{zq}. \quad (24)$$

where m_{zp} and m_{zq} are the eigenvalues of the initial density matrix (22). Using expansion (18) up to second order in $\frac{\beta}{\alpha} = \frac{d_{13}}{d_{12}}$ and $\frac{\delta}{\alpha} = \frac{d_{14}}{d_{12}}$ and keeping terms up to eighth order in $x = d_{12}t$, one can determine the normalized 0-quantum (J_{0Q}) and 2-quantum (J_{2Q}) intensities:

$$J_{0Q} = 1 - \frac{3}{2}x^2 \left[1 + \frac{1}{3} \left(\frac{\delta}{\alpha} \right)^2 + \frac{2}{3} \left(\frac{\beta}{\alpha} \right)^2 \right] + x^4 \left[\frac{7}{6} + \frac{2}{3} \left(\frac{\delta}{\alpha} \right)^2 + 2 \left(\frac{\beta}{\alpha} \right)^2 \right] \\ - x^6 \left[\frac{2}{5} + \frac{1}{3} \left(\frac{\delta}{\alpha} \right)^2 + \frac{4}{3} \left(\frac{\beta}{\alpha} \right)^2 \right] + \frac{2}{21}x^8 \left[\frac{47}{60} + \left(\frac{\delta}{\alpha} \right)^2 + \frac{22}{5} \left(\frac{\beta}{\alpha} \right)^2 \right] \quad (25)$$

and

$$J_{2Q} = -x^2 \left[\frac{3}{4} + \frac{1}{2} \left(\frac{\beta}{\alpha} \right)^2 \right] + x^4 \left[\frac{7}{12} + \frac{1}{3} \left(\frac{\delta}{\alpha} \right)^2 + \left(\frac{\beta}{\alpha} \right)^2 \right] \\ - x^6 \left[\frac{1}{5} + \frac{1}{6} \left(\frac{\delta}{\alpha} \right)^2 + \frac{2}{3} \left(\frac{\beta}{\alpha} \right)^2 \right] + \frac{1}{21}x^8 \left[\frac{47}{60} + \left(\frac{\delta}{\alpha} \right)^2 + \frac{22}{5} \left(\frac{\beta}{\alpha} \right)^2 \right]. \quad (26)$$

In the case where the parameter x is not small we have to take exactly into account of the terms containing x . By summation of series in (12) over n , m , l , and k we obtain exact analytical expressions which give the time dependence of intensities of 0 - quantum coherence

$$J_{0Q} = 1 - 2J_{2Q} \quad (27)$$

and of 2-quantum coherence

$$J_{2Q} = -\frac{1}{4} + \frac{1}{4} \cos(x\sqrt{5}) \cos(x) - \frac{x\sqrt{5}}{50} \sin(x\sqrt{5}) \cos(x) \left[\left(\frac{\delta}{a} \right)^2 + 3 \left(\frac{\beta}{a} \right)^2 \right] \\ - x^2 \left\{ \frac{\cos(x\sqrt{5}) \cos(x)}{5} \left[\left(\frac{\beta}{a} \right)^2 + \frac{3}{4} \left(\frac{\delta}{a} \right)^2 \right] + \frac{\sqrt{5}}{20} \left(\frac{\delta}{a} \right)^2 \sin(x\sqrt{5}) \sin(x) \right\} \quad (28)$$

in which the interaction between the nearest neighbours is taken into account exactly. Now let us compare the time dependence of the MQ intensities given by (25) - (28) with those obtained using an exact solution of the evolution equation and using computer simulation. The exact solution gives the following expressions for J_{0Q}

$$J_{0Q} = 1 - 2J_{2Q} \quad (29)$$

and for J_{2Q}

$$J_{2Q} = -\frac{1}{4} + \frac{1}{8} \cos(K_+x) \cos \frac{P_+x}{2} \cos \frac{P_-x}{2} + \frac{1}{8} \cos(K_-x) \cos(Px) \quad (30)$$

where

$$K_{\pm} = \left(\frac{\delta}{\alpha}\right) \pm 1 \quad (31)$$

$$P = \left(\left(\frac{\delta}{\alpha}\right)^2 + 2 \left(\frac{\delta}{\alpha}\right) + 5 + 4 \left(\frac{\beta}{\alpha}\right)^2 \right)^{\frac{1}{2}} \quad (32)$$

$$P_{\pm} = R_{+} \pm R_{-} \quad (33)$$

$$R_{\pm} = \left(\left(\frac{\delta}{\alpha}\right)^2 - 2 \left(\frac{\delta}{\alpha}\right) + 5 \pm 8 \frac{\beta}{\alpha} + 4 \left(\frac{\beta}{\alpha}\right)^2 \right) \quad (34)$$

Figs. 1 and 2 show the evolution of the normalized 0Q and 2Q coherences in the linear chain of four nuclear spins 1/2 coupled by DDI, where $\frac{\beta}{\alpha} = \frac{1}{8}$, $\frac{\delta}{\alpha} = \frac{1}{27}$ and at $t = 0$ the spin system is in thermal equilibrium determinates by matrix (16). These figures present also the results of computer simulation performed using the MATLAB package. One can see that expressions (25) and (26) gives a good agreement with the exact analytical and numerical results for $x \leq 1$, while expressions (27) and (28) agree up to $x = 5$. Note that our results coincide also with the dependences obtained in paper [21].

As a second example let us consider the MQ dynamics in a ring of six spins coupled by dipole-dipole interaction. The ratios of the dipole-dipole coupling constants are given by $\frac{\beta}{\alpha} = \frac{1}{3\sqrt{3}}$ and $\frac{\delta}{\alpha} = \frac{1}{8}$, where α is the coupling constant of the nearest neighbors, β is the coupling constant of the next-nearest neighbors, δ is the coupling constant of the next-next-nearest neighbors. Calculation using expansion (18) up to second order in ratios of $\frac{\beta}{\alpha}$ and $\frac{\delta}{\alpha}$ and eighth order in $x = \alpha t$ gives zero intensity for 6Q coherence. The non-zero intensity can be obtained by taking into account exactly of terms related to the interaction between the nearest neighbors. Using Eq. (18) we obtain the analytical expression of the intensities of 6Q coherence calculated up to second order in ratio of $\frac{\beta}{\alpha}$ and $\frac{\delta}{\alpha}$

$$J_{6Q} = -\frac{\left(\frac{\delta}{\alpha}\right)^2}{497664} [84x(2 \cos x + \cos 2x) + 88 \sin x - 5(32 \sin 2x + \sin 4x)]^2 \quad (35)$$

It is interesting that J_{6Q} coherence up to second order does not depend on the interaction between the next-nearest neighbors. The validity of the time dependence of J_{6Q} according to Eq. (35) was tested by comparison with numerical results. Fig. 3 shows that the prediction of Eq. (35) gives a reasonable agreement with the simulation data up to $x \leq 7$, which

coincides with the estimation using the dipolar coupling constant: $x < \frac{\alpha}{\delta} = 8$. The results of our simulation, shown in Fig. 3, coincide with the dependences obtained in Refs. [13, 18].

Conclusion

In conclusion, the perturbation method was developed which is based on the expansion of the operator exponent in a perturbation series. It allows us to apply the perturbation approach to the description of the MQ spin dynamics in solids. The exact and perturbation analytical expressions were obtained which describe 0Q and 2Q dynamics in a cluster in the form of a linear chain of four nuclear spins coupled by DDI in solids. The calculated 0Q- and 2Q intensities versus the duration of the preparation period agree with the simulation data [21]. The obtained analytical expression for 6Q dynamics in a ring of six spins coupled by DDI is in a good agreement with the results of numerical simulation. The developed method can be extended to include various power of the operators and applied to the description of a wide range of physical problems that deal with dynamics of dipolar coupling spins in solids.

Acknowledgments

The authors are grateful to E. B. Fel'dman and I. I. Maximov (Institute of Problems of Chemical Physics, Chernogolovka) and A. K. Khitrin (Kent State University, Kent) for useful discussions. This research was supported by a Grant from the U.S.-Israel Binational Science Foundation (BSF).

-
- [1] A. Abragam, Principle of nuclear magnetism, Oxford University Press, London, 1961.
 - [2] R. Bruschweiler and R. R. Ernst, Chem. Phys. Lett. 264, 393 (1997).
 - [3] J. S. Waugh, Mol. Phys. 95, 731 (1998).
 - [4] E. B. Fel'dman and S. Lacelle, J. Chem. Phys. 108, 4709 (1998).
 - [5] U. Haeberlen, *High Resolution NMR Spectroscopy in Solids: Selective Averaging*, Academic Press, New York, San Francisco, London, 1976.
 - [6] D. G. Cory, A. F. Fahmy, T. F. Havel, Proc. Natl. Acad. Sci U.S.A., 94, 1634 (1997).
 - [7] N. Gershenfeld and I. L. Chuang, Science, 275, 350 (1997).
 - [8] B. Kane, Nature (London), 393, 133 (1998).
 - [9] J. B. Murdoch, W. S. Warren, D. P. Weitekamp, and A. Pines, J. Magn. Reson., 60, 205 (1984).

- [10] J. Baum, M. Munovitz, A. N. Garroway, A.Pines, J.Chem.Phys. **83**, 2015 (1985).
- [11] A. K. Roy and K. K. Gleason, J. Magn. Res. A, **120**, 139 (1996).
- [12] S. I. Doronin, Phys.Rev. A, **68**, 052306 (2003).
- [13] S. I. Doronin, I.I.Maksimov, E.B.Fel'dman, J. Exp. Theor.Phys. **91**,597 (2000).
- [14] J. Baugh, A. Kleinhammes, D. Han, Q. Wang, and Y. Wu, Science, **294**, 1505 (2001).
- [15] M. G. Rudavets and E. B.Fel'dman, JETP Letter, **75**, 635 (2002).
- [16] A. R. Kessel, R. R. Nigmatullin, N. M. Yakovleva, e-print quant-ph/0212138.
- [17] I. J. Lowe and R. E. Norberg Phys. Rev. **107** , 46 (1957).
- [18] Jae-Seung Lee and A. K. Khitrin, e-print quant-ph/0402132.
- [19] R Bellman, *Introduction to Matrix Analysis* (New York: McGraw-Hill),1960.
- [20] R. M. Wilcox, J. Math. Phys., **8**, 962 (1967).
- [21] G. Cho and J. P. Yesinowski, J.Chem.Phys., **100**, 15716 (1996).

Captions for figures.

Figure 1

Fig. 1. Time dependences (in units of $\frac{1}{\alpha}$) of the normalized intensities of 0Q coherence in a linear chain of four nuclear spins 1/2 coupled by DDI. Solid line is the exact solution (Eq.(29)), dotted line is the calculation using Eq.(25), dashed line is the calculation using Eq.(27), open circles present the simulation results of [21], and down solid triangles are our numerical simulation.

Figure 2

Fig. 2. Time dependences (in units of $\frac{1}{\alpha}$) of the normalized intensities of 2Q coherence in a linear chain of four nuclear spins 1/2 coupled by DDI. Solid line is the exact solution (Eq.(30), dotted line is the calculation using Eq.(28), dashed line is the calculation using Eq.(26), open circles present the simulation results of [21], and down solid triangles are our numerical simulation.

Figure 3

Fig.3

Time dependences of the intensities of 6Q coherence in a ring of six spins. The dashed line is the prediction of Eq. (35) up to second order in ratio of $\frac{\delta}{\alpha}$, and the solid line is the simulation data from [13, 18].