

Dephasing and breakdown of adiabaticity in the splitting of Bose Einstein Condensates

L. Pezzé^{1,2}, A. Smerzi^{1,2}, G.P. Berman¹, A.R. Bishop¹ and L.A. Collins¹

1) *Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

2) *Istituto Nazionale per la Fisica della Materia BEC-CRS*

and Dipartimento di Fisica, Università di Trento, I-38050 Povo, Italy

(Dated: March 23, 2022)

We study the quantum dynamics of a BEC condensate trapped in a double-well potential with a rising interwell barrier. We analytically find the characteristic time scales of the splitting process and compare our results with numerical analyses available in the literature. In first stage of the dynamics, the condensate follows adiabatically the rising of the interwell barrier. At a critical time t_{ad} , small amplitude fluctuations around the average trajectory increase exponentially fast, signaling the break-down of adiabaticity. We have found a highly non-trivial dependence of the dephasing time t_D , defined by $\sigma_\phi(t_D) = 1$, where $\sigma_\phi(t)$ is the dynamical quantum phase spreading, on t_{ad} and on the ramping time of the interwell barrier.

PACS numbers:

Introduction: From the early observations of a Bose Einstein Condensate (BEC) there has been a growing interest in theoretical and experimental studies of a condensate in a double-well trap. There are several goals related to these studies: i) to understand the analog of the Josephson effect in this type of system; ii) to clarify the meaning of the phase in quantum mechanics; and iii) to create interferometers working at the Heisenberg limit. The recent experimental realization [1] of a stable double-well trap is renewing the interest in these topics. Even though residual sources of noise are still limiting the coherence life time of the two coupled systems, the experimental progress is very promising [2] for future developments and possible technological applications.

In this paper we study, in a two-mode approximation, the dynamical splitting of a condensate trapped in a double-well while ramping up the interwell barrier. This problem has raised some controversy in the past. In [3], Javanainen and Wilkens (JW) analyzed the condensate splitting in a two-stage stages model: first, an initial condensate is partially split in two parts by ramping a potential barrier in the middle. The ramping rate is $R_I \ll \omega_p$, where ω_p is the Josephson plasma frequency. Because of this inequality the process is strictly adiabatic. It ends in a regime when it is possible to neglect the particle exchange between the two wells. In the second stage, the barrier is risen to infinity at a rate $R_{II} \gg \omega_p$, and then the system is left alone for a time t . There has been a debate between JW and Leggett and Sols (LS) [4] about the rate at which the two halves of the condensate lose their phase memory in this second stage. The crucial point was the estimation of the ground state phase fluctuations. In [5], Javanainen and Ivanov (JI) clarified the issue by analyzing numerically a two-mode model of a realistic continuous splitting.

In the present paper, we study the splitting problem within an analytically solvable model. We recover simple analytical estimates of the various time scales of the problem, which agree quite well with the JI estimates. Moreover, we emphasize that even after the loss of adiabaticity it is still not possible to neglect the tunneling exchange between the wells. As a consequence, the dephasing time, in which the system loses memory about the relative phase between the two condensates, has a complicated dependence on the ramping time of the interwell barrier. This has important consequences when studying the realizability of a BEC interferometer.

The main results of this paper are the analytical expression of the time of breakdown of adiabaticity and the dephasing time as a function of both the initial condition and the typical time scale of the splitting process.

The Quantum-Phase model. In this section, we review the Quantum-Phase model (QPM) of a condensate trapped in a symmetric two-well trap at zero temperature. The second quantization Hamiltonian of a system of bosons interacting with a δ -pseudopotential is given by

$$\hat{H}(t) = \int dz \hat{\Psi}^\dagger(z, t) \left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial z^2} + V(z, t) \right) \hat{\Psi}(z, t) + \frac{g}{2} \int dz \hat{\Psi}^\dagger(z, t) \hat{\Psi}^\dagger(z, t) \hat{\Psi}(z, t) \hat{\Psi}(z, t), \quad (1)$$

where $\hat{\Psi}$ is the bosonic field operator, and $V(z, t)$ is the time-dependent external double-well potential; $g = \frac{4\pi\hbar^2 a}{m}$ is the strength of the interparticle interaction, with a being the s -wave scattering length. The two-mode ansatz reads

$$\hat{\Psi}(z, t) = \psi_1(z, t) \hat{a}_1 + \psi_2(z, t) \hat{a}_2, \quad (2)$$

where $\psi_{1,2}(z, t)$ can be constructed as sum and difference of the first symmetric and antisymmetric Gross-Pitaevskii dynamical wave-functions in the double well trap. The operator $\hat{a}_{1,2}^\dagger$ ($\hat{a}_{1,2}$) creates (destroys) a particle in the modes 1,2, respectively. In the following, we will decouple the “external dynamics” (the evolution of the wave function) and

the ‘‘internal dynamics’’ (the evolution of the operators) [6]. The important time scale of the external dynamics is given by the trapping oscillation period $\tau_z = 2\pi/\omega_z$, where ω_z is the trap frequency. If the barrier is raised on a time scale $\Delta t \gg \tau_z$, then the final wave function $\psi_{1,2}(z, t)$ will correspond to the ground state of the 1,2 well respectively. If $\Delta t \ll \tau_z$, on the other hand, the splitting process excites the system, increasing the condensate energy. Menotti *et al.* in [6] indicate the revival time of coherence, τ_r [7, 8], as an upper bound for the internal dynamics. In fact, when $\Delta t \gg \tau_r$ the phase coherence is lost during the splitting process, and the two final condensates will be independently exhibiting no phase coherence. In current experiments, $\tau_z \ll \tau_r$, with τ_r longer than the life time of the condensate. Therefore, raising the potential barrier at a rate $\tau_z \ll \Delta t \ll \tau_r$, we can decouple the internal and the external dynamics. In the two-mode approximation the Hamiltonian of the system is [5, 9]

$$\hat{H} = \frac{E_c}{4} \left(\hat{a}_1^\dagger \hat{a}_1^\dagger \hat{a}_1 \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2^\dagger \hat{a}_2 \hat{a}_2 \right) - \frac{E_j}{N} \left(\hat{a}_1^\dagger \hat{a}_2 + \hat{a}_2^\dagger \hat{a}_1 \right). \quad (3)$$

The operator $\hat{N} = \hat{n}_1 + \hat{n}_2 = \hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2$ is the total number of particles, and it commutes with \hat{H} . The quantity E_j is the ‘‘Josephson coupling energy’’, and E_c is the ‘‘one-site energy’’:

$$E_c = 2g \int dz |\psi_1(z, t)|^2 = 2g \int dz |\psi_2(z, t)|^2, \quad (4)$$

$$E_j = -N \int dz \left[\frac{\hbar^2}{2m} \frac{\partial \psi_1(z, t)}{\partial z} \frac{\partial \psi_2(z, t)}{\partial z} + \psi_1(z, t) V(z, t) \psi_2(z, t) \right]. \quad (5)$$

It is convenient to study Eq.(3) in the Bargmann phase-states representation [9]. We write a general state in the Hilbert space of the two-mode system as

$$|\psi\rangle = \int_{-\pi}^{+\pi} \frac{d\phi}{2\pi} \psi(\phi) |\phi\rangle, \quad (6)$$

where ϕ is the relative phase between the two modes, and

$$|\phi\rangle = \sum_{n=-N/2}^{N/2} \frac{e^{in\phi}}{\sqrt{n!}} |n\rangle \quad (7)$$

are un-normalized vectors of the overcomplete Bargmann base, written in the relative number of particles n . In the Bargmann representation the action of any operators on $|\psi\rangle$ can be represented in terms of differential operators acting on the associated $\psi(\phi)$. The main consequence of the overcompleteness is the non-standard inner product between Bargmann vectors (7) $\langle \phi | \theta \rangle \approx \cos^N \left(\frac{\phi - \theta}{2} \right)$. It affects the inner product between states (6) written in the Bargmann representation.

In the limit $E_j \ll N^2 E_c$, and rescaling the time as $t \rightarrow t E_c / 2\hbar$, the dynamical equation for the 2π -periodic phase amplitude $\Psi(\phi, t)$ is

$$i \frac{\partial \Psi(\phi, t)}{\partial t} = - \frac{\partial^2 \Psi(\phi, t)}{\partial \phi^2} - \Gamma(t) \cos(\phi) \Psi(\phi, t), \quad (8)$$

with $\Gamma(t) = 2E_j(t)/E_c$. In this paper, we study the dynamical evolution of the wave function $\Psi(\phi, t)$ when the initial condensate is split with a symmetric double-well potential. This experiment has been recently realized [1], with the two wells ramped apart linearly in time. The distance between the center of the wells evolves according to $d(t) = d_0 + d_{fin} t / \Delta t_R$, where d_0 and d_{fin} are the initial and final distances, respectively, and Δt_R is the total ramping time. With this setup it is possible to find, both in WKB approximation [10] and by a numerical 1-D simulation, that the Josephson coupling energy evolves in time with the exponential law $E_j(t) = E_j(0) e^{-t/\tau}$, where the effective ramping time $\tau = \Delta t_R \hbar / \sqrt{2m(V_0 - \mu)d_0^2}$ depends of the particle mass m , the initial height of the potential barrier V_0 , and the chemical potential μ . In the two-mode approximation the one-site energy E_c remains constant during the dynamics. We notice that E_j scales exponentially with the interwell distance only when the condensates are well separated. During the initial splitting, when the chemical potential is close to the interwell barrier, the dynamics remains adiabatic. The adiabaticity will break down at a large separation of the two condensates, in the deep tunneling regime, which will be the focus of the next sections.

The two-mode model has been extensively discussed in the literature. In general, it is expected to work when the ground state and the first excited state are close to each other in energy and well separated from higher energy modes. It works in the limit of weak atom-atom interaction [5] and small atom number, $Ng \ll \omega_z$, where ω_z is the

frequency of the trap in the separation direction. It also has limited validity in the case of a low potential barrier when it is not allowed to neglect higher excitation modes. However, it becomes increasingly accurate by rising the potential barrier. In this case, in fact, the two lower lying modes become closer in energy and separated from the higher ones. As pointed out by Menotti *et al.* [6], a fundamental condition for the validity of the two-mode model is the different time scale which characterizes the “internal” and “external” dynamics. This decoupling is at the basis of the two-mode ansatz (2). Through a gaussian variational model, and numerical simulations of the mode functions [6], it is possible to conclude that, to a good approximation and for reasonable regimes, the wave function dynamics are independent of the operator dynamics.

Numerical Solution. We first solved numerically equation (8). Figure (1) presents a plot of the normalized $|\Psi(\phi, t)|^2$ at different times. Superimposed on the phase amplitude, the blue line presents the cosine potential in arbitrary units. At $t = 0$ (Fig. (1,A)) the phase amplitude $\Psi(\phi, t)$ is in its ground state, which, for sufficiently high values of Ej , is well approximated by a Gaussian. At $t > 0$, the height of the potential barrier decreases exponentially (here we choose $\tau = 5$ msec), and the phase amplitude spreads (Fig. (1,B)) until it touches the borders at $\pm\pi$. The wiggles in Fig. (1,C)) arise from the interference between the two overlapping tails in the region around $\phi \sim \pi$, which eventually spread over the full region Fig. (1,D). The period of the oscillations of the interference pattern depends on the velocity of spreading of the phase amplitude: the more adiabatic the expansion, the smaller the number of oscillations.




FIG. 1: Profile of the phase amplitude $|\Psi(\phi)|^2$ at different times: A) $t = 0$, corresponds to the ground state at the beginning of the dynamic; B) $t = t_{ad} = 23$ msec, which corresponds to the breakdown of adiabaticity as calculated by Eq. (25); C) $t = t_D = 200$ msec, corresponds to the dephasing time ($\sigma_\phi(t_D) = 1$) as calculated by the equation (39); D) $t = 500$ msec. The blue line represent the cosine potential (in rescaled units) decreasing with $\tau = 5$ msec.

Variational Approach: We study Eq. (8) with a time-dependent variational approach [11]. The time evolution of the variational parameters is characterized by the minimization of an action with the effective Langragian

$$L(q_i, \dot{q}_i) = i\langle \Psi \dot{\Psi} \rangle - \langle \Psi \hat{H} \Psi \rangle, \quad (9)$$

with q_i being the time-dependent parameters of the phase amplitude $\Psi(\phi, q_i(t))$. This provides the familiar Lagrange equations

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} = \frac{\partial L}{\partial q_i}. \quad (10)$$

We choose the time-dependent variational phase amplitude

$$\Psi(\phi, t) = \frac{1}{(2\pi\sigma_\phi^2(t))^{1/4}} \exp\left(-\frac{\phi^2}{4\sigma_\phi^2(t)} + i\frac{\delta(t)}{2}\phi^2\right), \quad (11)$$

with the condition that the width $\sigma_\phi(t) \ll 2\pi$ during the dynamics. At the beginning of the dynamics, the phase amplitude is rather narrow and, feeling only the quadratic part of the $\cos(\phi)$ potential, it can be approximated by

the gaussian (11). During the first stage of the dynamics, the phase amplitude will follow the instantaneous ground state of the system. A breakdown of adiabaticity will occur at time t_{ad} , which will depend on the ramping time τ_R . The Gaussian ansatz will fail when the wave-function touches the borders $\phi = \pm\pi$, namely when $\sigma_\phi(t) \gtrsim 1$. With the variational ansatz (11) the equation of motion for the width of the phase amplitude becomes:

$$\ddot{\sigma}_\phi = \frac{1}{\sigma_\phi^3} - 2\sigma_\phi \Gamma(t) e^{-\sigma_\phi^2/2}. \quad (12)$$

Adiabatic Variational Solution. We can calculate the adiabatic solution from equation (12) by imposing the adiabaticity condition $\dot{\sigma}_\phi = 0$. In this limit, we have $\Gamma \gg 1$, and the dynamics are governed by the particle exchange through the two wells, which keeps the phase coherence between the two condensates. We obtain the equation

$$\sigma_{ad}(t) = \sqrt{4W\left(\sqrt{\frac{1}{4\sqrt{2}\Gamma(t)}}\right)} = \sqrt{4W\left(\frac{1}{8}\sqrt{\frac{E_c}{E_j(0)}} e^{t/2\tau}\right)}, \quad (13)$$

where $W(x)$ is the Lambert-W function [12]. In Figure (3) we compare the adiabatic solution with the numerical one for the values $\tau = 5$ msec and $\tau = 20$ msec.

Linear approximation We now solve equation (12) seeking a solution of the form $\sigma_\phi(t) = \sigma_{ad}(t) + \varepsilon(t)$. We expect that there will be a time at which the solution $\varepsilon(t)$ becomes a significant correction to $\sigma_{ad}(t)$: this will give the criterion for the breakdown of adiabaticity. Replacing $\sigma_\phi(t) = \sigma_{ad}(t) + \varepsilon(t)$ in equation (12), where $\sigma_{ad}(t)$ is the adiabatic solution (13), we obtain a second-order differential equation for $\varepsilon(t)$

$$\ddot{\varepsilon}(t) = -\left(\frac{4 + \sigma_{ad}^2(t)}{\sigma_{ad}^4(t)}\right) \varepsilon(t) + O(\varepsilon(t)^2), \quad (14)$$

where we neglected quadratic terms in $\varepsilon(t)$. We are interested in studying this equation for times $0 \leq t \leq t_{ad}$: in this range we can assume $\sigma_{ad}^2(t) \ll 1$, and equivalently $E_j(t) \gg E_c$ and $\Gamma(t) \gg 1$. In the regime $\sigma_{ad} \ll 1$, we approximate $(4 + \sigma_{ad}^2(t))/\sigma_{ad}^4(t) \sim 4/\sigma_{ad}^4(t)$. In the same approximation we expand Eq. (13) in a Taylor series for $1/\Gamma(t) \approx 0$, neglecting quadratic terms. Accounting for the rescaling of the time, the equation (14) becomes

$$\ddot{\varepsilon}(t) = -\frac{4E_c E_j(0)}{\hbar^2} e^{-t/\tau} \varepsilon(t). \quad (15)$$

This is the equation of a harmonic oscillator with a time dependent (exponentially-decreasing) frequency. It is possible to find the solution of this equation in terms of Bessel functions:

$$\varepsilon(t) = C J_0\left(2\sqrt{\frac{4E_c E_j(0)}{\hbar^2}} \tau e^{-t/2\tau}\right) + D Y_0\left(2\sqrt{\frac{4E_c E_j(0)}{\hbar^2}} \tau e^{-t/2\tau}\right), \quad (16)$$

where J_0 is the zero-order Bessel function of the first kind and Y_0 is the zero-order Bessel function of the second kind [13], while C and D are constants which depend on the initial conditions $\varepsilon(0)$, and $\dot{\varepsilon}(0)$. Choosing as initial condition $\varepsilon(0) = 0$, and $\dot{\varepsilon}(0) \neq 0$, we have

$$C = -\frac{\dot{\varepsilon}(0)}{\sqrt{A}} \left(\frac{Y_0(2\sqrt{A}\tau)}{Y_1(2\sqrt{A}\tau)J_0(2\sqrt{A}\tau) - Y_0(2\sqrt{A}\tau)J_1(2\sqrt{A}\tau)} \right), \quad (17)$$

$$D = \frac{\dot{\varepsilon}(0)}{\sqrt{A}} \left(\frac{J_0(2\sqrt{A}\tau)}{Y_1(2\sqrt{A}\tau)J_0(2\sqrt{A}\tau) - Y_0(2\sqrt{A}\tau)J_1(2\sqrt{A}\tau)} \right), \quad (18)$$

where $A = 4E_c E_j(0)/\hbar^2$. Figure (2) presents a plot of the functions $J_0(2\sqrt{A}\tau e^{-t/2\tau})$ and $Y_0(2\sqrt{A}\tau e^{-t/2\tau})$ for $A = 1$ and for an arbitrary time τ .

Initially, the solutions are oscillating with a $\pi/2$ phase difference:

$$J_0(2\sqrt{A}\tau e^{-t/2\tau}) \approx \sqrt{\frac{1}{\pi\sqrt{A}\tau}} e^{t/4\tau} \cos\left(2\sqrt{A}\tau e^{-t/2\tau} - \frac{\pi}{2}\right), \quad (19)$$




FIG. 2: Behavior of the functions $J_0(2\tau e^{-t/\tau})$ (blue line) and $Y_0(2\tau e^{-t/\tau})$ (red line). The dashed green line represents the exponential asymptotes $e^{t/4\tau}$.

$$Y_0(2\sqrt{A}\tau e^{-t/2\tau}) \approx \sqrt{\frac{1}{\pi\sqrt{A}\tau}} e^{t/4\tau} \sin\left(2\sqrt{A}\tau e^{-t/2\tau} - \frac{\pi}{2}\right). \quad (20)$$

The amplitude of the oscillations increase exponentially as $e^{-t/4\tau}$. For large times ($t \gtrsim 4\tau$) the functions approach their asymptotic behavior: J_0 tends to a constant value

$$J_0(2\sqrt{A}\tau e^{-t/2\tau}) \rightarrow 1 \quad (21)$$

independently of A and τ , and Y_0 diverges linearly in time

$$Y_0(2\sqrt{A}\tau e^{-t/2\tau}) \rightarrow -\frac{2}{\pi} \ln(2\sqrt{A}\tau) \frac{t}{2\tau}. \quad (22)$$

The times t_J and t_Y in the figure (2) indicate the last oscillation of the Bessel functions Y_0 and J_0 . They are defined by the equations

$$2\sqrt{A}\tau e^{-t_Y/2\tau} = 0.8935, \quad (23)$$

$$2\sqrt{A}\tau e^{-t_J/2\tau} = 2.4048. \quad (24)$$

We interpret this divergence at $t > 4\tau$ as the signature of breakdown of adiabaticity. From equations (23) and (24) it is natural to define the time of breakdown of adiabaticity t_{ad} by the relation

$$2\sqrt{A}\tau e^{-t_{ad}/2\tau} = c, \quad (25)$$

where c is an arbitrary number ~ 1 . Since $A = 4E_c E_j(0)/\hbar^2$, $E_J(t) = E_J(0)e^{-t/\tau}$, and introducing the Josephson oscillation frequency $\omega_j(t) = \sqrt{E_c E_j(t)}/\hbar$, we can rewrite Eq. (25) as

$$\frac{1}{\omega_j(t_{ad})} = \frac{4}{c} \tau, \quad (26)$$

which gives

$$t_{ad} = 2\tau \ln\left(\omega_j(0) \frac{4}{c} \tau\right). \quad (27)$$

Eq. (26) coincides with the definition of breakdown of adiabaticity suggested with a different heuristic argument by Javanainen in [5] with $\alpha = 4/c$. In his work, Javanainen estimates numerically the parameter $\alpha = 2\pi$. This value is in reasonable agreement with our analytical calculations. We have $c = 2/\pi = 0.63$, which is in good agreement with our estimations. The more important physical result suggested by equation (26) is the fact that the constant c does not depend of the time scale τ . This has a physical meaning [4]. Initially the system is in its ground state, and we suppose $E_j(0) \gg E_c$; the phase dispersion is $\sigma_\phi(0) \ll 1$, and the system feels only the quadratic part of the $\cos(\phi)$ potential. By ramping the two wells, the quantity $E_j(t)$ decreases with time scale τ . As long as $\omega_j(t) \gg 1/\tau$, the system adjusts itself in such a way that it is always in the ground state, and the change of the potential is adiabatic. Over the time t_{ad} defined by the equation (26), the frequency $\omega_j(t)$ becomes so small that it is impossible for the system to adjust in the ground state following the decreasing of the tunneling rate. The evolution of the system is no longer adiabatic.

Figure (3) shows a comparison between the numerical solution (blue line) of the dynamical equation (8) and the variational adiabatic solution (red line) given by the equation (13). The figures (3A, B) refer to the cases $\tau = 5$ msec and $\tau_R = 20$ msec, respectively. In these figures the time t_{ad} is defined by equation (25) with $c = 1$.




FIG. 3: Comparison between different solutions of the equation (8) for two cases: A) $\tau_R^{eff} = 5$ msec, and B) $\tau_R^{eff} = 20$ msec. The blue line represents the numerical solution, and the red line represents the variational adiabatic solution given by the equation (13). We also indicate the time of breakdown of adiabaticity as given by the eq. (25): $t_{ad} = 21$ msec for A), and $t_{ad} = 111$ msec for B).

Dephasing Time: We define the dephasing time t_D as the time needed for the phase dispersion to become of order 1. Roughly, at $t = t_D$ the phase amplitude $\Psi(\phi, t)$ reaches the borders $\phi = \pm\pi$, and we lose every information about the phase. It might be useful to recall that in a single experiment a well defined phase will actually be measured. However, in different experiments, repeated in identical conditions, the measured phases would differ, with a mean square fluctuation σ_ϕ . The main result of this section will be the analytical estimate of t_D . We first notice that (comparing the red and blue lines in the figure (4)) we can approximate well the numerically exact phase amplitude with a gaussian even for $t > t_{ad}$. We can therefore expect that the gaussian variational ansatz can give a good estimate of t_D . Thus, we approximate the QPM \hat{H}_ϕ with the Hamiltonian of a harmonic oscillator with a time dependent frequency:

$$H_\phi = -\frac{E_c}{2\hbar} \frac{\partial^2}{\partial \phi^2} - \frac{E_j(0) e^{-t/\tau}}{2\hbar} \phi^2. \quad (28)$$

The quantum evolution of the system governed by Eq.(28) can be calculated exactly in the Wigner phase space. We first need to calculate the solution of the classical equation of motion

$$\ddot{\phi}(t) = -\frac{E_c E_j(0)}{\hbar^2} e^{-t/\tau} \phi(t), \quad (29)$$

which can be expressed in terms of Bessel functions of the first and the second kind. With $A = E_c E_j(0)/\hbar^2$, we have (similarly with equation (15) and (16)):

$$\phi(t) = C J_0(2\sqrt{A}\tau e^{-t/2\tau}) + D Y_0(2\sqrt{A}\tau e^{-t/2\tau}), \quad (30)$$




FIG. 4: Comparison between the numerical solution of equation (8) (red line), the variational approach obtained by solving the equation (12) (blue line), and the harmonic oscillator approximation obtained by substituting $\cos \phi \rightarrow -\phi^2/2$ in Eq. (8)

where C and D depend of the initial conditions $\phi_0 \equiv \phi(0)$ and $\dot{\phi}_0 \equiv \dot{\phi}(0)$:

$$C = \frac{\phi_0}{J_0(2\sqrt{A}\tau)} - \frac{Y_0(2\sqrt{A}\tau)}{J_0(2\sqrt{A}\tau)} \frac{\frac{\dot{\phi}_0}{\sqrt{A}} J_0(2\sqrt{A}\tau) - \phi_0 J_1(2\sqrt{A}\tau)}{Y_1(2\sqrt{A}\tau) J_0(2\sqrt{A}\tau) - Y_0(2\sqrt{A}\tau) J_1(2\sqrt{A}\tau)}, \quad (31)$$

$$D = \frac{\frac{\dot{\phi}_0}{\sqrt{A}} J_0(2\sqrt{A}\tau) - \phi_0 J_1(2\sqrt{A}\tau)}{Y_1(2\sqrt{A}\tau) J_0(2\sqrt{A}\tau) - Y_0(2\sqrt{A}\tau) J_1(2\sqrt{A}\tau)}. \quad (32)$$

We can now calculate the phase dispersion $\sigma_\phi^2(t) = \langle \phi(t, \phi_0, \dot{\phi}_0)^2 \rangle - \langle \phi(t, \phi_0, \dot{\phi}_0) \rangle^2$, where the brackets indicate the integration over the initial conditions ϕ_0 and $\dot{\phi}_0$ averaged with the Wigner transform $P(\phi_0, \dot{\phi}_0, t)$. For an harmonic oscillator the Wigner transform [14] becomes

$$P(\phi_0, \dot{\phi}_0, t) = \frac{1}{\pi E_c} \int_{-\infty}^{+\infty} d\xi \frac{e^{-\frac{(\phi_0 - \xi)^2}{4\sigma_\phi^2}}}{(2\pi\sigma_\phi^2)^{\frac{1}{4}}} \frac{e^{-\frac{(\phi_0 + \xi)^2}{4\sigma_\phi^2}}}{(2\pi\sigma_\phi^2)^{\frac{1}{4}}} e^{2i\frac{\hbar}{E_c}\xi\dot{\phi}_0} \quad (33)$$

$$= \frac{e^{-\frac{\phi_0^2}{2\sigma_\phi^2}}}{(2\pi\sigma_\phi^2)^{\frac{1}{4}}} \frac{e^{-\frac{\dot{\phi}_0^2}{2\sigma_\phi^2}}}{(2\pi\sigma_\phi^2)^{\frac{1}{4}}}, \quad (34)$$

where $\sigma_{\dot{\phi}_0} = E_c/(2\hbar\sigma_\phi)$ is the width of the distribution $P(\phi_0, \dot{\phi}_0, t)$ in the variable $\dot{\phi}_0$ canonically conjugated to ϕ_0 . Notice that, in equation (33), we have integrated between $\pm\infty$, and not between $\pm\pi$, which is consistent with the condition $\sigma_{\phi_0}, \sigma_{\dot{\phi}_0} \ll 2\pi$. Since the Wigner transform (33) is symmetric in ϕ_0 and $\dot{\phi}_0$, we have that $\langle \phi(t, \phi_0, \dot{\phi}_0) \rangle = 0$, and

$$\begin{aligned} \sigma_\phi(t)^2 &= \langle \phi(t, \phi_0, \dot{\phi}_0)^2 \rangle \\ &= \int_{-\infty}^{+\infty} d\phi_0 d\dot{\phi}_0 \left(C(\phi_0, \dot{\phi}_0) J_0(2\sqrt{A}\tau e^{-t/2\tau}) + D(\phi_0, \dot{\phi}_0) Y_0(2\sqrt{A}\tau e^{-t/2\tau}) \right)^2 P(\phi_0, \dot{\phi}_0, t) \\ &= \left(C^* J_0(2\sqrt{A}\tau e^{-t/2\tau}) + D^* Y_0(2\sqrt{A}\tau e^{-t/2\tau}) \right)^2, \end{aligned} \quad (35)$$

where $C^*(D^*) = \int_{-\infty}^{+\infty} d\phi_0 d\dot{\phi}_0 C(\phi_0, \dot{\phi}_0) (D(\phi_0, \dot{\phi}_0)) P(\phi_0, \dot{\phi}_0, t)$. If we define $K(2\sqrt{A}\tau) \equiv Y_1(2\sqrt{A}\tau)J_0(2\sqrt{A}\tau) - Y_0(2\sqrt{A}\tau)J_1(2\sqrt{A}\tau)$, we can write

$$(C^*)^2 = \frac{\sigma_\phi^2(0)}{J_0^2(2\sqrt{A}\tau)} \left[1 + \frac{Y_0(2\sqrt{A}\tau)J_1(2\sqrt{A}\tau)}{K(2\sqrt{A}\tau)} \right]^2 + \frac{\sigma_\phi^2(0)}{A} \frac{Y_0^2(2\sqrt{A}\tau)}{K^2(2\sqrt{A}\tau)}, \quad (36)$$

$$(D^*)^2 = \frac{\sigma_\phi^2(0)}{A} \frac{J_0^2(2\sqrt{A}\tau)}{K^2(2\sqrt{A}\tau)} + \sigma_\phi^2(0) \frac{J_1^2(2\sqrt{A}\tau)}{K^2(2\sqrt{A}\tau)}. \quad (37)$$

Notice that in the limit $\tau \rightarrow 0$, we recover the free evolution dynamics as given by equation (41). As shown by figure (2), the harmonic oscillator approximation remain valid even for $t > t_{ad}$. The dephasing of the phase amplitude occurs at $t_D > \tau$. We can therefore use the asymptotic expansion $t \gg \tau$ for the Bessel functions $J_O(2\sqrt{A}\tau e^{-t/2\tau}) \rightarrow 1$ and $Y_O(2\sqrt{A}\tau e^{-t/2\tau}) \rightarrow \frac{2}{\pi} \ln(\sqrt{A}\tau) - \frac{t}{\pi\tau}$. We can rewrite eq. (35) as

$$\sigma_\phi(t)^2 = \left(C^* + D^* \left(\frac{2}{\pi} \ln(\sqrt{A}\tau) - \frac{t}{\pi\tau} \right) \right)^2. \quad (38)$$

The dephasing time t_D is defined by the relation $\sigma_\phi(t_D) = 1$, from which we obtain

$$t_D = 2\tau \ln(\sqrt{A}\tau) + \pi\tau \left(\frac{1 + C^*}{D^*} \right). \quad (39)$$

This is the central result of our paper. Figure (5) presents the dependences t_D and t_{ad} as a function of τ .




FIG. 5: Dependences of t_D (blue line) and t_{ad} (red line) as the functions of τ . The dephasing time is given by Eq. (39), the time of breakdown of adiabaticity is given by Eq. (25).

Holding time: In [1] the two potential wells are separated with a ramping time Δt_R . At the end of the ramping, the condensates are held in the trap for a time Δt_{hold} . We can study the phase dispersion during this time by using the variational approximation (12) with a constant parameter Γ . If we have $\Delta t_R = \beta\tau$ where the $\beta \gtrsim 1$, we can neglect the potential energy term in Eq. (12), which becomes

$$\ddot{\sigma}_\phi = \frac{1}{\sigma_\phi^3}. \quad (40)$$

The phase-width evolves freely in time. Accounting for the rescaling of the time, the solution of this equation is given by

$$\sigma_\phi^2(t) = \sigma_\phi^2(\Delta t_R) + \frac{E_c^2}{4\hbar^2 \sigma_\phi^2(\Delta t_R)} (t - \Delta t_R)^2, \quad (41)$$

where $\sigma_\phi^2(\Delta t_R)$ is the gaussian width at $t = \Delta t_R$ (end of the ramping). For $t \lesssim t_{\Delta t_R} + 2\sigma_\phi^2(\Delta t_R)\hbar/E_c$, we have that $\sigma_\phi(t)$ is almost constant, while for $t \gg \Delta t_R + 2\sigma_\phi^2(\Delta t_R)\hbar/E_c$ the width $\sigma_\phi(t)$ is a linear function of time. In figure (6) we present the evolution of $\sigma_\phi(t)$ for different values of Δt_R . In this figure $\tau = 5$ msec, $E_c = 0.001 \times \hbar$ kHz, and $E_c = 100 \times \hbar$ kHz. To take into account the boundaries $\phi = \pm\pi$, we compare the numerical solution with the quantity

$$\bar{\sigma}_\phi^2(t) = \int_{-\pi}^{+\pi} d\phi |\psi(\phi)|^2 \phi^2, \quad (42)$$

where $|\psi(\phi)|^2 \sim \exp(-\phi^2/2\sigma_\phi^2(t))$ and $\sigma_\phi^2(t)$ is given by Eq. (41).

In the model of Legget and Sols [4, 15, 16] the free evolution takes place with the breakdown of adiabaticity.




FIG. 6: Evolution of $\sigma_\phi(t)$ for different values of Δt_R . The red line represents the case $\Delta t_R = \infty$ corresponding to an unlimited ramping the two potential wells. The other lines refer to different choices of finite Δt_R . We approximate the evolution of $\sigma_\phi(t)$ after the ramping of the wells by the free evolution given by equation (41), and we take into account the boundaries $\pm\pi$ through the equation (42).

This model assumes that $\Delta t_R = t_{ad}$. We can check this model by looking at Fig. (41): the red line represents the numerical solution of equation (8), for $\tau = 5$ msec, the blue line represents the free expansion occurring at the breakdown of adiabaticity t_{ad} as defined by eq. (25). We can see that the model of Legget and Sols gives a reasonable estimation of the phase spreading, however it over-estimates the dephasing time.

Repeated Interference Experiments: It has been recently shown by Castin and Dalibard [17] and Javanainen and Yoo [18] that the phase of a condensate is established by measurement. Two BECs, initially in number "Fock" states, will interfere and have a definite relative phase. A different phase, however, will be measured in different experiments, so that averaging over the ensemble, no interference is observed. To illustrate the effect of dephasing, we present here a simple qualitative analysis. We consider the interference between two independent (not overlapping) condensates in the Thomas Fermi (TF) approximation. Initially the two condensates are in equilibrium, the condensate 1 being centered at $x = -d/2$, and the condensate 2 at $x = d/2$. We assume that at $t = 0$ the order parameter is described by the linear combination

$$\hat{\Psi}(\vec{r}) = \hat{\Psi}_1(\vec{r}) + e^{i\Phi} \hat{\Psi}_2(\vec{r}), \quad (43)$$

where $\Psi_1(\vec{r})$ and $\Psi_2(\vec{r})$ are the equilibrium wave functions (order parameters) of the two condensates, respectively, which we assume are well separated in space:

$$\Psi_1(\vec{r})\Psi_2(\vec{r}) \approx 0. \quad (44)$$

At $t = 0$ the trap is switched off. We neglect the interaction between the two condensates but we account for the atom-atom interaction in each single condensate, which is important during the free expansion. The total density $\rho = |\Psi|^2$ of the overlapping condensates exhibits modulation of the form

$$\rho(\vec{r}, t, \phi) = \rho_1(\vec{r}, t) + \rho_2(\vec{r}, t) + 2\sqrt{\rho_1(\vec{r}, t)\rho_2(\vec{r}, t)} \cos(S_1(\vec{r}, t) - S_2(\vec{r}, t) + \phi), \quad (45)$$




FIG. 7: Contour plot of the interference fringes calculated in the TF approximation. The density is averaged over several experiments, according to Eq. (47). The quantum dynamics determines a phase uncertainty σ : A) $\sigma = 0$, B) $\sigma = 1$, C) $\sigma = 2$, D) $\sigma = 3$.

where $\Psi_{1,2}(\vec{r}, t) = \sqrt{\rho_{1,2}(\vec{r}, t)} e^{S_{1,2}(\vec{r}, t)}$. By recalling the quadratic behavior of the phase in the TF approximation, we obtain asymptotically

$$S_1(\vec{r}, t) - S_2(\vec{r}, t) = \frac{md}{\hbar t} x. \quad (46)$$

In a single experiment, the interference pattern is characterized by the straight line fringes which are orthogonal to the x -axis (radial axis of the two cigar shaped parallel condensates). We obtain fringes perpendicular to the x -axis with spacing $\hbar t/md$ between two consecutive fringes. To experimentally test the quantum phase dynamics it would be necessary to average over several identical interferometric realizations. The relative phase will be chosen randomly with a gaussian distribution of width σ . The ensemble averaged density is therefore $\rho(\vec{r}, t) = \int_{-\pi}^{+\pi} d\phi \rho(\vec{r}, t; \phi) |\Psi(\phi, t)|^2 = \frac{1}{2}[\rho_1(\vec{r}, t) + \rho_2(\vec{r}, t) + \rho_{int}(\vec{r}, t)]$, with $\rho_{1,2}(\vec{r}, t) \equiv |\psi_{1,2}(\vec{r}, t)|$ the densities of each released condensate, and

$$\rho_{int}(\vec{r}, t) = 2\sqrt{\rho_1(\vec{r}, t)\rho_2(\vec{r}, t)} \int_{-\pi}^{+\pi} d\phi \cos\left(\frac{md}{\hbar t} x + \phi\right) \frac{e^{-\frac{\phi^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} \quad (47)$$

$$= 2\sqrt{\rho_1(\vec{r}, t)\rho_2(\vec{r}, t)} \cos\left(\frac{md}{\hbar t} x\right) e^{-\sigma^2/2}. \quad (48)$$

Conclusions: We have studied the dephasing and adiabaticity time scales in the splitting of a single Bose Einstein Condensate. We have focused on the quantum behavior of the system within a two-mode model, solving a Schrödinger-like equation in the phase variable within a variational gaussian ansatz. We have obtained the exact behavior in the adiabatic and free expansion regimes, and we have obtained analytical estimations for the dephasing time and the time when the dynamics depart from the adiabatic evolution. We have compared our analytical results with both full numerical solutions of the two-mode Hamiltonian, and the existing literature. Finally, we have shown how the spreading of phase distribution can affect the visibility of interference fringes in an ensemble of repeated experiments.

This work was supported by the Department of Energy under the contract W-7405-ENG-36 and DOE Office of Basic Energy Sciences.

[1] Y. Shin *et al.* *Phys. Rev. Lett.* **92**, 050405 (2004).

[2] L.A. Collins, L.Pezzé, A. Smerzi, G.P. Berman and A.R. Bishop *Phys. Rev. A*, to be published (2004)

[3] J. Javanainen and M. Wilkens, *Phys. Rev. Lett.* **78**, 4675 (1997)

- [4] A.J. Leggett and F.Sols, *Phys. Rev. Lett.* **81**, 1345 (1998).
- [5] J. Javanainen and M. Yu. Ivanov, *Phys. Rev. A* **60**, 2351 (1999).
- [6] C. Menotti, J.R. Anglin, J.I. Cirac and P. Zoller *Phys. Rev. Lett.* **63**, 023601 (2001)
- [7] M. Lewenstein and L. You, *Phys. Rev. Lett.* **77**, 3489 (1996)
- [8] P. Villain, M. Lewenstein, R. Dum, Y. Castin, L. You, A. Imamoglu, T.A.B. Kennedy, *J. Mod. Opt.* **44**, 1775 (1997)
- [9] J.R. Anglin, P. Drummond, A. Smerzi, *Phys. Rev. A* **64**, 063605 (2001).
- [10] A. Messiah, *Quantum Mechanics*, North-Holland, Amsterdam (1965), pag. 231
- [11] A. Smerzi and S. Raghavan, *Phys. Rev. A* **61**, 063601 (2000).
- [12] See the web site <http://mathworld.wolfram.com/LambertW-Function.html>
- [13] M. Abramowitz and I. A. Stegun, *Book of Mathematical Functions* (National Bureau of Standards, Washington, DC, 1984).
- [14] M. Hillery, R.F. O Connel, M.O. Scully, E.P. Wigner, *Phys. Rep.* **106**, 121 (1984)
- [15] A.J. Leggett and F. Sols, *Found. Phys.* **21**, 353 (1991)
- [16] I. Zapata, F. Sols and A.J. Leggett, *Phys. Rev. A* **57**, 28 (1998)
- [17] Y. Castin and J. Dalibard, *Phys. Rev. A* **55**, 4330 (1997)
- [18] J. Javanainen and S.M. Yoo, *Phys. Rev. Lett.* **76**, 161 (1996)