

Comment on "Charge expulsion and electric field in superconductors"

Tomio Koyama
Institute for Materials Research,
Tohoku University, Sendai 980-8577, Japan

March 22, 2004

Abstract

We propose a generalized London theory that can correctly describe the longitudinal and transverse responses of conventional superconductors to an electromagnetic field. The continuity equation is satisfied by use of a special gauge for the scalar and vector potentials. Our phenomenological theory provides a simple example of the Anderson-Higgs mechanism.

In ref.[1] Hirsch claimed that hole superconductors should carry a surface charge in the ground state even without an applied electric field and calculated phenomenologically the inhomogeneous charge appearing inside a superconductor. In this comment we point out that the phenomenological calculation given in ref.[1] for the charge distribution is not correct, which includes a misunderstanding of the electrodynamics of a superconductor. Since a similar misunderstanding is also seen in recent articles [2, 3], it will be worth presenting the correct phenomenological theory based on the generalized London theory for the electrodynamics of a superconductor.

In ref.[1] the charge density $\rho(\mathbf{r})$ in a superconductor is assumed to satisfy the equation,

$$\rho(\mathbf{r}) = -\frac{1}{4\pi\lambda_L^2} \nabla^2 A_0(\mathbf{r}); \quad (1)$$

under the Lorenz gauge $\nabla \cdot \mathbf{A}(\mathbf{r}) + \frac{1}{c} \partial_t A_0(\mathbf{r}) = 0$ with $(A_0; \mathbf{A})$ being the scalar and vector potentials.

In eq.(1) λ_L is the London penetration depth, which indicates that the screening length of a longitudinal electric field in the superconducting state is equal to the London penetration depth, not the Thomas-Fermi screening length λ_{TF} . This point is stressed in ref.[1, 2, 3] as the remarkable electric properties in the superconducting state. However, eq.(1) cannot be accepted, because it is against the general understanding that the charge screening length is still given by λ_{TF} in the superconducting state. The charge density $\rho(\mathbf{r}; t)$ induced by an external electric field in the superconducting state can be calculated microscopically within the linear response theory, using the formula, $\rho(\mathbf{r}; t) = e^2 \int dt^0 \int d\mathbf{r}^0 K(\mathbf{r} - \mathbf{r}^0; t - t^0) A_0(\mathbf{r}^0; t^0)$, where $K(\mathbf{r} - \mathbf{r}^0; t - t^0)$ is the density-density correlation function, i.e., $K(\mathbf{r} - \mathbf{r}^0; t - t^0) = i(t - t^0) \langle [\hat{n}(\mathbf{r}; t); \hat{n}(\mathbf{r}^0; t^0)] \rangle$. In the lowest order (single-loop) approximation for $K(\mathbf{r} - \mathbf{r}^0; t - t^0)$ of a s-wave BCS superconductor one obtains the equation in the static limit,

$$\rho(\mathbf{r}) = e^2 \sum_{\mathbf{q}} K(\mathbf{q}) A_0(\mathbf{q}) e^{i\mathbf{q} \cdot \mathbf{r}}; \quad (2)$$

where

$$K(\mathbf{q}) = \sum_{\mathbf{k}} \frac{E_{\mathbf{k}} E_{\mathbf{k}+\mathbf{q}}}{E_{\mathbf{k}} E_{\mathbf{k}+\mathbf{q}} (E_{\mathbf{k}} + E_{\mathbf{k}+\mathbf{q}})^2}; \quad (3)$$

with $E_{\mathbf{k}} = \sqrt{\frac{\hbar^2 \mathbf{k}^2}{2m} + \Delta^2}$. From eq.(3) it follows

$$K(0) = \sum_{\mathbf{k}} \frac{\Delta^2}{E_{\mathbf{k}}^3} = 2N(0); \quad (4)$$

in the limit $q \rightarrow 0$, where $N(0)$ is the density of states at the Fermi level. Note that the gap function disappears after the k -integration. Then, one finds the equation valid in the long wavelength region,

$$\langle \mathbf{r} \rangle = -2e^2 N(0) A_0(\mathbf{r}) = -\frac{1}{4} \frac{1}{v_F^2} A_0(\mathbf{r}) : \quad (5)$$

Note that the charge screening length coincides with the Thomas-Fermi length $\lambda_{TF} = 1/(8e^2 N(0))$. From this observation one understands that the phenomenological theory should be constructed in terms of eq.(5) instead of eq.(1), together with the London equation. In this case the problem is that what gauge should be chosen, since the continuity equation is not fulfilled if one uses the Lorenz gauge, which indicates that the Lorenz gauge should be abandoned. Note that the substitution of eq.(5) and the London equation into the continuity equation, $\partial_t \tilde{\phi} + \nabla \cdot \mathbf{j} = 0$, leads to the equation as follows,

$$\tilde{\phi} - \tilde{A}(\mathbf{r};t) + \frac{1}{2} \frac{1}{v_F} \frac{1}{c} \partial_t A_0(\mathbf{r};t) = 0 : \quad (6)$$

We propose that eq.(6) should be regarded as the gauge condition for $(A_0; \tilde{A})$. Notice that the ratio of the screening lengths in eq.(6) can be rewritten as $\lambda_L^2 = \lambda_{TF}^2 = c^2 = \frac{1}{3} v_F^2 = \frac{1}{3} v_{GB}^2$, with v_F being the Fermi velocity, if one assumes $\lambda_{TF} = \frac{1}{2} \lambda_L$ and $\lambda_L = \frac{1}{3} \lambda_{GB}$ in the free electron model. Note that $v_{GB} = \frac{1}{3} v_F$ coincides with the velocity of the Goldstone boson in the superconducting state. The gauge function $\tilde{\phi}(\mathbf{r};t)$, which may be considered as the phase of the order parameter, i.e., the Goldstone boson, can be introduced by the gauge transformation, $\tilde{A} \rightarrow \tilde{A} - \frac{\hbar c}{e} \nabla \tilde{\phi}$ and $A_0 \rightarrow A_0 + \frac{\hbar}{e} \partial_t \tilde{\phi}$ with $e = 2e$ so that the new gauge field also satisfies eq.(6). It is also noted that the Maxwell-London equations in our theory can be derived from the gauge-invariant Lagrangian,

$$L = \frac{1}{8} \frac{1}{e^2} \frac{1}{\lambda_L^2} \frac{c^2}{v_{GB}^2} E A_0 + \hbar \partial_t \tilde{\phi}^2 - e \tilde{A}^2 - \hbar c \nabla \tilde{\phi}^2 + \frac{E^2}{8} + \frac{B^2}{8} : \quad (7)$$

This Lagrangian describes the massive plasma modes. Under the gauge condition (6) one can obtain the

dispersion relations, $\epsilon_T(p^2) = \epsilon_p^2 (1 + \frac{1}{\lambda_L^2} p^2)$ for the transverse one and $\epsilon_L(p^2) = \epsilon_p^2 (1 + (v_{GB}^2 = c^2) \frac{1}{\lambda_L^2} p^2)$ for the longitudinal one with $\epsilon_p = c = \lambda_L$. In the neutral case i.e., $e = 0$, the Lagrangian is reduced to the one for the massless scalar field, $L = \hbar c^2 (8 e^2 \frac{1}{\lambda_L^2}) v_{GB}^2 \partial_t \tilde{\phi}'^2 - \tilde{\phi}'^2$. Then, eq.(7) indicates that the Goldstone boson $\tilde{\phi}'$ is absorbed into the gauge field $(A_0; \tilde{A})$, i.e., the Anderson-Higgs mechanism [4]. Thus, our phenomenological theory provides the consistent description for the electrodynamics of a superconductor, in which the charge screening length is given by the Thomas-Fermi length. The gauge condition given in eq.(6) was proposed in ref.[5] and is called the phason gauge.

References

- [1] J.E. Hirsch, Phys.Rev.B 68 184502 (2003).
- [2] J.Govaerts, D.Bertrand and G.Stenuit, Supercond.Sci.Technol 14 463 (2001).
- [3] J.E.Hirsch, Phys.Rev.B 69 214515 (2004).
- [4] See, for example, S.Weinberg, The Quantum Theory of Fields (Cambridge University Press, Cambridge 1996) chapt.21.
- [5] H.Matsumoto and H.Umezawa, Fortschr.Phys. 24 357 (1976).