

Nonextensive statistical mechanics, anomalous diffusion and central limit theorems *

Constantino Tsallis

*Santa Fe Institute, 1399 Hyde Park Road,
Santa Fe, New Mexico 87501, USA*

and

*Centro Brasileiro de Pesquisas Fisicas,
Rua Xavier Sigaud 150,
22290-180 Rio de Janeiro-RJ, Brazil.*

(Dated: March 23, 2022)

We briefly review Boltzmann-Gibbs and nonextensive statistical mechanics as well as their connections with Fokker-Planck equations and with existing central limit theorems. We then provide some hints that might pave the road to the proof of a new central limit theorem, which would play a fundamental role in the foundations and ubiquity of nonextensive statistical mechanics. The basic novelty introduced within this conjectural theorem is the *generalization of the hypothesis of independence* of the N random variables being summed. In addition to this, we also advance some nonlinear dynamical (possibly exact) relations which generalize the concepts of Lyapunov exponents, entropy production per unit time, and their interconnection as first proved by Pesin for chaotic systems.

I. INTRODUCTION

As well known, *thermodynamics* is the basic branch of physics which focuses on the generic connections between variables (temperature, pressure, volume, energy, entropy and many others) that play an important role in the description of the *macroscopic* world. Boltzmann and Gibbs provided a magnificent connection of thermodynamics with the *microscopic* world [1, 2]. This connection, normally referred to as *Boltzmann-Gibbs (BG) statistical mechanics* (or simply *statistical mechanics* since it was basically the only one to be formulated along more than one century), turns out to be the appropriate one for ubiquitous systems in nature. It is based on the following axiomatic expression for the entropy:

$$S_{BG} \equiv -k \sum_{i=1}^W p_i \ln p_i, \quad (1)$$

with

$$\sum_{i=1}^W p_i = 1, \quad (2)$$

where p_i is the probability associated with the i^{th} microscopic state of the system, and k is Boltzmann constant. In the particular case of equiprobability, i.e., $p_i = 1/W$ ($\forall i$), Eq. (1) yields the celebrated *Boltzmann principle* (as referred to by Einstein himself [3]):

$$S_{BG} = k \ln W. \quad (3)$$

From now on, and without loss of generality, we shall take k equal to unity.

For continuous variables, the *BG* entropy is written as

$$S_{BG} \equiv - \int dx p(x) \ln p(x) \quad (x \in \mathbb{R}^d), \quad (4)$$

with

$$\int dx p(x) = 1. \quad (5)$$

If x happens to carry physical units, we write Eq. (4) as follows:

$$S_{BG} \equiv - \int dx p(x) \ln \left[\left(\prod_{r=1}^d \sigma_r \right) p(x) \right], \quad (6)$$

where $\sigma_r > 0$ carries the same units as the r^{th} component of the d -dimensional variable x . Clearly, when x carries no units (i.e., when $x \in \mathbb{R}^d$), we take $\sigma_r = 1$ ($\forall r$). In fact, everytime this is possible (and it *is* possible most of the time), we shall adapt the physical units in such a way that $\sigma_r = 1$ ($\forall r$) *even when x does have physical units*. Consistently, unless otherwise specified, we shall use Eq. (4) for the general continuous case. If we consider the particular case $p(x) = \sum_{i=1}^W p_i \delta(x - x_i)$ (with $\sum_{i=1}^W p_i = 1$, $\{x_i\}$ being some set of values, and $\delta(z)$ being Dirac's delta distribution), Eq. (4) recovers Eq. (1).

For quantum systems, the *BG* entropic form is written as

$$S_{BG} \equiv -\text{Tr} \rho \ln \rho, \quad (7)$$

with

$$\text{Tr} \rho = 1, \quad (8)$$

ρ being the density operator or matrix. When the $W \times W$ matrix ρ is diagonalized, it shows the set $\{p_i\}$ in its diagonal. In what follows, depending on the context, we shall use either the discrete form (Eqs. (1) and (2)), or

*Invited paper to appear in *Milan Journal of Mathematics* (2005).

the continuous form (Eqs. (4) and (5)), or the matricial form (Eqs. (7) and (8)).

In spite of its tremendous power and usefulness, the *BG* concepts and statistical mechanics appear to be *not universally applicable*. Indeed, there is a plethora of natural and artificial systems (see, for instance, [4] and references therein) for which they do not provide the adequate mathematical frame for handling physically relevant quantities. This fact started being explicitly recognized at least as early as in 1902 by Gibbs himself: see page 35 of [2], where he addresses anomalies related to systems such as gravitation. A formalism becomes therefore desirable which would address such anomalous systems. A vast class of them (although surely not *all* of them) appears to be adequately discussed within a generalization of the *BG* theory, frequently referred to as *nonextensive statistical mechanics*. This theory was first introduced in 1988 [5], and then refined in 1991 [6] and 1998 [7]. It is based on the following generalization of S_{BG} :

$$S_q \equiv \frac{1 - \sum_{i=1}^W p_i^q}{q - 1} \quad (q \in \mathbb{R}; S_1 = S_{BG}) . \quad (9)$$

Expressions (4), (6) and (7) are respectively generalized into

$$S_q \equiv \frac{1 - \int dx [p(x)]^q}{q - 1} , \quad (10)$$

$$S_q \equiv \frac{1 - \int d \left[x / \left(\prod_{r=1}^d \sigma_r \right) \right] \left[\left(\prod_{r=1}^d \sigma_r \right) p(x) \right]^q}{q - 1} , \quad (11)$$

and

$$S_q \equiv \frac{1 - \text{Tr } \rho^q}{q - 1} \quad (12)$$

For equiprobability (i.e., $p_i = 1/W$, $\forall i$), Eq. (9) yields

$$S_q = \ln_q W , \quad (13)$$

with the *q-logarithm* function defined as [9]

$$\ln_q z \equiv \frac{z^{1-q} - 1}{1 - q} \quad (z \in \mathbb{R}; z > 0; \ln_1 z = \ln z) . \quad (14)$$

Its inverse function, the *q-exponential*, is given by [9]

$$e_q^z \equiv [1 + (1 - q)z]^{1/(1-q)} \quad (e_1^z = e^z) \quad (15)$$

if the argument $1 + (1 - q)z$ is positive, and equals zero otherwise.

In Section II we briefly review a few known results concerning the probability distributions which extremize the entropy, and are ultimately associated with macroscopically stationary states. In Section III we briefly review available results related to Fokker-Planck equations. In Section IV we review the standard and the

Lévy-Gnedenko central limit theorems, and argue about the possible formulation of a new theorem that would generalize the standard limit theorem. Such a theorem would provide an important mathematical cornerstone for nonextensive statistical mechanics and its ubiquity in nature. Providing hints that could help formulating and proving the theorem constitutes the main reason of the present paper.

II. STATISTICAL MECHANICS

Entropy is necessary to formulate statistical mechanics but it is not sufficient. Indeed, we must also introduce the concept of *energy*. The easiest (and more frequently used) way to do so is addressing the so called (by Gibbs) *canonical ensemble*. It corresponds to the ubiquitous physical situation in which the system of interest is in contact with a (large by definition) *thermostat* which, at equilibrium (or at the physically relevant stationary state, more generally speaking), imposes to the system its temperature. The system is typically described by a quantum Hamiltonian, and is characterized by the spectrum of energies $\{E_i\}$ ($i = 1, 2, \dots, W$) defined as the eigenvalues associated with the Hamiltonian and its boundary conditions. The probability distribution at the relevant stationary state is the one which extremizes the entropy under the norm and the energy constraints.

A. Boltzmann-Gibbs statistical mechanics

At thermal equilibrium we must optimize S_{BG} (as given by Eq. (1)) with the norm constraint as given by Eq. (2), and the energy constraint given as follows:

$$\sum_{i=1}^W p_i E_i = U_{BG} , \quad (16)$$

where U_{BG} is given and referred to as the *internal energy*. By following the Lagrange method, we define the quantity

$$\Phi_{BG} \equiv S_{BG} + \alpha \left(\sum_{i=1}^W p_i - 1 \right) - \beta \left(\sum_{i=1}^W p_i E_i - U_{BG} \right) , \quad (17)$$

where α and β are the Lagrange parameters (their signs have been chosen following tradition). The extremizing condition $\delta \Phi_{BG} / \delta p_j = 0$ yields

$$p_j = e^{\alpha - 1 - \beta E_j} \quad (j = 1, 2, \dots, W) . \quad (18)$$

The use of Eq. (2) allows the elimination of the parameter α . We then obtain the celebrated Boltzmann-Gibbs weight

$$p_i = \frac{e^{-\beta E_i}}{Z_{BG}} \quad (i = 1, 2, \dots, W) , \quad (19)$$

where β is connected with the thermostat *temperature* T through $\beta \equiv 1/T$, and the *partition function* is defined as follows:

$$Z_{BG}(\beta) \equiv \sum_{j=1}^W e^{-\beta E_j}. \quad (20)$$

It is precisely the present Eqs. (19) and (20) that are referred to as *dogma* by the mathematician Takens! [8].

Clearly, if we replace the probability distribution (19) into Eq. (16), we obtain the thermodynamically important relation between the inverse temperature β and the internal energy U_{BG} .

As a subsidiary comment, whose relevance will become transparent later on, let us remark that the *BG* weight (19) can be seen as the solution of the *linear* ordinary differential equation

$$\frac{dy}{dx} = ay \quad (y(0) = 1). \quad (21)$$

Indeed, its solution is given by

$$y = e^{ax}, \quad (22)$$

which reproduces Eq.(19) through the identification $(x, a, y) \equiv (E_i, -\beta, Z_{BG} p_i)$.

B. Nonextensive statistical mechanics

We want now to optimize S_q (as given by Eq. (9)) with the norm constraint still given by Eq. (2), and the energy constraint generalized as follows (see [7]):

$$\frac{\sum_{i=1}^W p_i^q E_i}{\sum_{i=1}^W p_i^q} = U_q, \quad (23)$$

(referred to as *q-expectation value* or *q-mean value*) or, equivalently,

$$\sum_{i=1}^W p_i^q (E_i - U_q) = 0. \quad (24)$$

The functional (17) is generalized into

$$\Phi_q \equiv S_q + \alpha \left[\sum_{i=1}^W p_i - 1 \right] - \beta \left[\sum_{i=1}^W p_i^q (E_i - U_q) \right], \quad (25)$$

and the extremizing condition $\delta \Phi_q / \delta p_j = 0$ yields

$$p_j = \left(\frac{q}{\alpha} \right)^{1/(1-q)} e_q^{-\beta(E_j - U_q)} \quad (j = 1, 2, \dots, W). \quad (26)$$

The use of Eq. (2) allows, as before, the elimination of the parameter α , thus obtaining the generalized weight

$$p_i = \frac{e_q^{-\beta(E_i - U_q)}}{Z_q} \quad (i = 1, 2, \dots, W), \quad (27)$$

with

$$Z_q(\beta) \equiv \sum_{j=1}^W e_q^{-\beta(E_j - U_q)}. \quad (28)$$

This probability distribution corresponds to a *maximum* (*minimum*) of S_q for $q > 0$ ($q < 0$). For $q = 0$, the entropy is constant, namely $S_0 = W - 1$, and the distribution is given by $p_i = [1 - \beta(E_i - U_0)] / \sum_{j=1}^W [1 - \beta(E_j - U_0)]$ (we recall the *cutoff* of the q -exponential function for $q < 1$, i.e., the states for which $1 - \beta(E_i - U_0) < 0$ do *not* contribute).

As a subsidiary comment, let us remark that the weight (27) can be seen as the solution of the *nonlinear* ordinary differential equation

$$\frac{dy}{dx} = ay^q \quad (y(0) = 1). \quad (29)$$

Indeed, the solution is given by

$$y = e_q^{ax}, \quad (30)$$

which reproduces Eq.(27) through the identification $(x, a, y) \equiv (E_i - U_q, -\beta, Z_q p_i)$.

C. The Gaussian case

Let us now assume a stochastic real variable x such that

$$\langle x \rangle \equiv \int_{-\infty}^{\infty} dx x p(x) = 0, \quad (31)$$

and

$$\langle x^2 \rangle \equiv \int_{-\infty}^{\infty} dx x^2 p(x) = D, \quad (32)$$

where $D > 0$ is given. The maximization of entropy (4) with constraints (5) and (32) yields the celebrated Gaussian distribution, more precisely

$$p(x) = \sqrt{\frac{\beta}{\pi}} e^{-\beta x^2} \quad (33)$$

where β is a Lagrange parameter. By using Eq. (32), we immediately obtain $\beta = 1/2D$, hence

$$p(x) = \frac{e^{-x^2/(2D)}}{\sqrt{2\pi D}} \quad (34)$$

We straightforwardly verify that constraint (31) is satisfied as well.

A more general case is to assume the following constraints:

$$\langle x \rangle \equiv \int_{-\infty}^{\infty} dx x p(x) = C, \quad (35)$$

and

$$\langle (x - \langle x \rangle)^2 \rangle \equiv \int_{-\infty}^{\infty} dx (x - \langle x \rangle)^2 p(x) = D. \quad (36)$$

The extremizing distribution is then given by

$$p(x) = \sqrt{\frac{\beta}{\pi}} e^{-\beta(x - \langle x \rangle)^2}, \quad (37)$$

or, equivalently,

$$p(x) = \frac{e^{-(x - C)^2/(2D)}}{\sqrt{2\pi D}}. \quad (38)$$

D. The q -generalization of the Gaussian case

Let us assume that the constraints are now even more general, namely

$$\langle x \rangle_q \equiv \frac{\int_{-\infty}^{\infty} dx x [p(x)]^q}{\int_{-\infty}^{\infty} dx [p(x)]^q} = C_q, \quad (39)$$

and

$$\langle (x - \langle x \rangle_q)^2 \rangle_q \equiv \frac{\int_{-\infty}^{\infty} dx (x - \langle x \rangle_q)^2 [p(x)]^q}{\int_{-\infty}^{\infty} dx [p(x)]^q} = D_q, \quad (40)$$

where $P(x) \equiv [p(x)]^q / \int dy [p(y)]^q$ is, in the literature, referred to as the *escort distribution*.

The associated distribution extremizing S_q is now given by

$$p(x) = A_q \sqrt{\beta} e^{-\beta(x - \langle x \rangle_q)^2}, \quad (41)$$

where we have [10] $A_q = \sqrt{(q-1)/\pi} \Gamma(1/(q-1)) / \Gamma((3-q)/[2/(q-1)])$ for $q > 1$, and $A_q = \sqrt{(1-q)/\pi} \Gamma((5-3q)/[2(1-q)]) / \Gamma((2-q)/(1-q))$ for $q < 1$, $\Gamma(z)$ being the Riemann function.

The use of constraint (40) straightforwardly provides $\beta = 1/[(3-q)D_q]$, which, replaced in Eq. (41), yields

$$p(x) = \begin{cases} \frac{A_q}{\sqrt{(3-q)D_q}} \frac{1}{[1 + \frac{q-1}{(3-q)D_q} (x - C_q)^2]^{1/(q-1)}} & (q > 1) \\ \frac{A_q}{\sqrt{(3-q)D_q}} [1 - \frac{1-q}{(3-q)D_q} (x - C_q)^2]^{1/(1-q)} & (q < 1) \end{cases} \quad (42)$$

In the $q < 1$ case, the support is compact, and the distribution vanishes outside the interval $|x - C_q| \leq \sqrt{(3-q)D_q/(1-q)}$.

If $\langle x \rangle_q = 0$, distributions (41) and (42) take respectively the simple forms

$$p(x) = A_q \sqrt{\beta} e^{-\beta x^2}, \quad (43)$$

and

$$p(x) = \frac{A_q}{\sqrt{(3-q)D_q}} e^{-x^2/[(3-q)D_q]}, \quad (44)$$

which appear frequently in various contexts.

It can be seen [11] that these distributions are analytic extensions (to real values of q) of the Student's t -distribution and the r -distribution, for $q > 1$ and $q < 1$ respectively. Consistently, there is an asymptotic long *power-law tail* for $q > 1$, and a *compact support* for $q < 1$. There is an upper bound for q , namely $q = 3$, imposed by the norm constraint (5). More precisely, the admissible values of q are $q < 3$. It deserves to be mentioned that also constraint (40) has an upper bound in order to be finite, which happens to be *precisely the same*, i.e. $q = 3$.

III. DIFFUSION AND FOKKER-PLANCK EQUATIONS

Normal diffusion (i.e., the one which satisfies $\langle x^2 \rangle \propto t$, typical of Brownian motion [12]) is characterized by the heat equation (the simplest form of the Fokker-Planck equation)

$$\frac{\partial p(x, t)}{\partial t} = D \frac{\partial^2 p(x, t)}{\partial x^2} \quad (D > 0). \quad (45)$$

By assuming, at $t = 0$, the paradigmatic form

$$p(x, 0) = \delta(x), \quad (46)$$

we obtain the following exact solution:

$$p(x, t) = \frac{e^{-x^2/2Dt}}{\sqrt{2\pi Dt}} \quad (t \geq 0) \quad (47)$$

There is of course an infinity of possible generalizations of Eq. (45), which is *linear* and defined through *integer derivatives*. We address here two important such generalizations, both of them associated with *anomalous diffusion*, i.e., violating the relation $\langle x^2 \rangle \propto t$. The first one remains linear but replaces the second derivative of the right term by a *fractional derivative*; it yields $\langle x^2 \rangle \rightarrow \infty$ ($\forall t > 0$). The second one is *nonlinear* but preserves the integer derivatives; it yields $\langle x^2 \rangle \propto t^\alpha$ ($\alpha \neq 1$).

The first one is as follows:

$$\frac{\partial p(x, t)}{\partial t} = D \frac{\partial^\gamma p(x, t)}{\partial x^\gamma} \quad (D > 0; 0 < \gamma < 2). \quad (48)$$

The exact solution associated with the $t = 0$ condition (46) is given by

$$p(x, t) = \frac{1}{(Dt)^{1/\gamma}} L_\gamma(x/(Dt)^{1/\gamma}) \quad (t \geq 0), \quad (49)$$

where $L_\gamma(z)$ is the Lévy distribution of index γ . The $\gamma \rightarrow 2$ limit obviously corresponds to the Gaussian solution (47). The $\gamma = 1$ particular case corresponds to the Cauchy-Lorentz distribution

$$p(x, t) = L_1(x/(Dt)) = \frac{1}{\pi} \frac{Dt}{(Dt)^2 + x^2} \quad (50)$$

For all other values of γ different from 1 and 2, the Lévy distribution has no direct analytic expression, and is expressed only through its Fourier transform.

The second generalization of Eq. (45) is as follows:

$$\frac{\partial p(x, t)}{\partial t} = D \frac{\partial^2 [p(x, t)]^\nu}{\partial x^2} \quad (\nu \in \mathbb{R}). \quad (51)$$

The exact solution associated with the $t = 0$ condition (46) is given [13, 14] by

$$p(x, t) = \frac{A_q}{\sqrt{(3-q)(Dt)^{1/(3-q)}}} e_q^{-x^2/[(3-q)(Dt)^{2/(3-q)}]} \quad (t \geq 0) \quad (52)$$

with

$$q = 2 - \nu < 3, \quad (53)$$

with $D > 0$ if $\nu > 0$, and $D < 0$ if $\nu < 0$. If we rewrite the diffusion coefficient as $D \equiv \bar{D}/\nu$, Eq. (51) can be rewritten as follows:

$$\frac{\partial p(x, t)}{\partial t} = \bar{D} \frac{\partial^2 [p(x, t)]^\nu - 1}{\partial x^2 \nu}, \quad (54)$$

which, in the limit $\nu \rightarrow 0$, becomes

$$\frac{\partial p(x, t)}{\partial t} = \bar{D} \frac{\partial^2 [\ln p(x, t)]}{\partial x^2}, \quad (55)$$

whose solution is known to be the Cauchy-Lorentz distribution.

If we start from any $t = 0$ distribution different from (46), the solution $p(x, t)$ does not coincide with Eq. (52), but, nevertheless, it asymptotically approaches [15] Eq. (52) for $t \rightarrow \infty$. In other words, the solution (52) constitutes an *attractor* in the space of the distributions, i.e., it is a solution which is *robust*. For all $q < 3$, x scales [14] with $t^{1/(3-q)}$. Consequently, in all cases for which the second moment of $p(x, t)$ is *finite* (i.e., for $q < 5/3$) we obtain $\langle x^2 \rangle \propto t^{2/(3-q)}$. What happens for $0 \leq 5/3 < 3$ is that the prefactor of $t^{2/(3-q)}$ diverges. In this case, it is convenient to focus on the *width* of the distribution, or, equivalently, on $\langle x^2 \rangle_q$.

To close these remarks about Fokker-Planck equations, let us write down a quite general one, namely

$$\frac{\partial^\delta p(x, t)}{\partial t^\delta} = D \frac{\partial^\gamma [p(x, t)]^{2-q}}{\partial |x|^\gamma} \quad ((\delta, \gamma, q) \in \mathbb{R}^3), \quad (56)$$

where we have used, for convenience, q rather than ν (related through Eq. (53)). The regions we address are $0 < \delta \leq 1$, $0 < \gamma \leq 2$ and $q < 3$. Not surprising, the solutions for arbitrary (δ, γ, q) are not known. But, nevertheless, we depict the $\delta = 1$ regions in Fig. 1. This will clarify the particular cases that are related to the central limit theorems we are interested in.

IV. CENTRAL LIMIT THEOREMS

In one of its simplest versions, the standard central limit theorem may be formulated as follows. Let $\{x_l\}$




FIG. 1: $\delta = 1$ regions that are relevant for the Central Limit Theorems (CLT). The solution of Eq. (56) for $(\gamma, q) = (2, 1)$ is a Gaussian distribution (which optimizes the entropy S_{BG} , and whose variance is finite). For $q = 1$ and $0 < \gamma < 2$, the solutions of Eq. (56) are Lévy distributions (which asymptotically decay as $1/|x|^{1+\gamma}$, and whose variance diverges). For $\gamma = 2$ and $q < 3$, the solutions are q -Gaussians (which optimize the entropy S_q , and whose variance is finite for $q < 5/3$ and diverges for $5/3 \leq q < 3$; their q -variance is finite for all $q < 3$; they have a compact support for $q < 1$ and asymptotically decay as $1/|x|^{2/(q-1)}$ for $1 < q < 3$). Within the standard CLT, we sum *independent* variables whose variance is *finite*; within the Lévy-Gnedenko CLT, we sum *independent* variables whose variance *diverges*; within the conjectural CLT, we sum *specialy correlated* variables whose q -variance is *finite*.

($l = 1, 2, \dots, N$) be a set of *independent* random variables, each of them satisfying the *same* symmetric (with regard to $x = 0$) distribution $p(x)$. Let also $p(x)$ be such that $\langle x^2 \rangle \equiv \int_{-\infty}^{\infty} dx x^2 p(x)$ is *finite*. Let us define now the sum variable

$$Z \equiv \sum_{l=1}^N x_l \quad (57)$$

The question to which the central limit theorem answers is *what is the probability distribution of the random variable Z when $N \rightarrow \infty$?* This answer happens to be very simple, namely a *Gaussian* (in the properly rescaled variables). If we call $p(Z, N)$ this distribution (with $p(Z, 1) = p(x)$), to exhibit the Gaussian we must use as abscissa Z/\sqrt{N} , and as ordinate $\sqrt{N}p(Z, N)$. Then, we gradually see emerging the Gaussian as $N \rightarrow \infty$. And, *what specific Gaussian?* The one whose second moment *precisely coincides* with $\langle x^2 \rangle$.

This theorem clearly is closely related to normal diffusion, hence to Eq. (45) and to its solution (47), where t plays the role of N .

Let us now address the other central limit theorem which is known, namely the Lévy-Gnedenko one. In its simplest version, it can be formulated as follows. As before, the set $\{x_l\}$ ($l = 1, 2, \dots, N$) is constituted of *independent* random variables, each of them satisfying

the *same* symmetric (with regard to $x = 0$) distribution $p(x)$. But now, we have the case where $\langle x^2 \rangle$ *diverges*. The question remains the same, namely *what is the probability distribution of the random variable Z when $N \rightarrow \infty$?* The answer now is: a *Lévy distribution* $L_\gamma(Z)$. *With what value for its index γ ?* All Lévy distributions decay, for $|x| \rightarrow \infty$, as $1/|x|^{1+\gamma}$. This behavior is preserved for the distribution of Z while N increases. Let us be more specific: the typical case is that for which $p(x)$ decays as $1/|x|^\mu$ with $1 < \mu < 3$. Then $\gamma = \mu - 1$. And *what specific Lévy distribution L_γ ?* The one which has *precisely the same coefficient* of the asymptotically dominant term. In other words, let us consider the *finite* value $\lim_{|x| \rightarrow \infty} [p(x)|x|^\mu]$. The particular Lévy distribution which is asymptotically approached when $N \rightarrow \infty$ is the one which has the *same* value for $\lim_{|x| \rightarrow \infty} [L_\gamma(x)|x|^{1+\gamma}]$. To see, on a graphic representation, the gradual emergence of the Lévy distribution while N increases, the abscissa must be $Z/N^{1/\gamma}$, and the ordinate must be $N^{1/\gamma}p(Z, N)$.

This theorem is closely related to the anomalous diffusion characterized by Eq. (48) and by its solution (49), where, as before, t plays the role of N .

Let us finally address the *conjectural* theorem that we are focusing in this paper. It is of course the one to be associated with the nonlinear Eq. (51) and its solution (52). *What hypothesis is to be violated in the two preceding and well studied theorems? The hypothesis of independence!* Indeed, we believe that the variables $\{x_i\}$ are to be assumed somehow *collectively correlated* in such a persistent manner that the correlation does *not* disappear even in the $N \rightarrow \infty$ limit. For the standard central limit theorem, the quantity which is preserved is the second moment $\langle x^2 \rangle \equiv \int_{-\infty}^{\infty} dx x^2 p(x)$. For the Lévy-Gnedenko theorem, the quantity which is preserved is the coefficient $\lim_{|x| \rightarrow \infty} [p(x)|x|^\mu]$. For this conjectural theorem, some quantity is expected to be preserved. Could it be $\langle x^2 \rangle_q \equiv \int_{-\infty}^{\infty} dx x^2 [p(x)]^q / \int_{-\infty}^{\infty} dx [p(x)]^q$?

It should be transparently clear at this point that we have *no definitive arguments* for proving this conjectural theorem. Various converging paths are nevertheless available that might inspire a (professional or *amateur*) mathematician the way to prove it. Galileo used to say that knowing a result is not neglectable in order to prove it! It is our best hope that his saying does apply in the present case! *So, what are these converging paths?* Although naturally intertwined, let us expose them along six different lines.

1. The q -product hint

The following generalization of the product operation has been recently introduced [16]. It is called *q -product* and is defined through

$$X \otimes_q Y \equiv [X^{1-q} + Y^{1-q} - 1]^{1/(1-q)} \quad (q \in \mathbb{R}) \quad (58)$$

We shall address the case where $X \geq 1$ and $Y \geq 1$ [17]. This product has the following properties:

- (i) $X \otimes_1 Y = XY$;
- (ii) $\ln_q(X \otimes_q Y) = \ln_q X + \ln_q Y$ (whereas $\ln_q(XY) = \ln_q X + \ln_q Y + (1-q)(\ln_q X)(\ln_q Y)$);
- (iii) $1/(X \otimes_q Y) = (1/X) \otimes_{2-q}(1/Y)$;
- (iv) $X \otimes_q (Y \otimes_q Z) = (X \otimes_q Y) \otimes_q Z = X \otimes_q Y \otimes_q Z = (X^{1-q} + Y^{1-q} + Z^{1-q} - 2)^{1/(1-q)}$;
- (v) $X \otimes_q 1 = X$;
- (vi) $X \otimes_q Y = Y \otimes_q X$;
- (vii) For fixed (X, Y) , $X \otimes_q Y$ monotonically increases with q .

Property (ii) is particularly important since it is directly relevant to the *extensivity* of S_q that will be addressed soon.

Notice that property (iii) involves, like the solution of Eq. (51) with Eq. (53), a $q \leftrightarrow (2-q)$ transformation.

We may apply this product to the number $W_{A_1+A_2+\dots+A_N}^{eff}$ of *allowed* states in a composed system whose subsystems A_1, A_2, \dots, A_N have respectively $W_{A_1}, W_{A_2}, \dots, W_{A_N}$ possible states (by *allowed* we mean that their probability is essentially *nonzero*). If $q = 1$ we have the total number $W_{A_1+A_2+\dots+A_N} = \prod_{i=1}^N W_{A_i}$ of states that are not only possible a priori, but even generically allowed. But, if $q \neq 1$, say $q < 1$, we expect correlations to inhibit (even forbidden occasionally) *some* of the states, i.e., to be associated with a probability close (in some sense) to zero. In this case, the *effective* total number $W_{A_1+A_2+\dots+A_N}^{eff}$ of allowed states is expected to be *smaller* than $W_{A_1+A_2+\dots+A_N}$. We expect to have basically

$$\begin{aligned} W_{A_1+A_2+\dots+A_N}^{eff} &\sim W_{A_1} \otimes_q W_{A_2} \otimes_q \dots \otimes_q W_{A_N} \\ &= \left[\left(\sum_{i=1}^N W_{A_i}^{(1-q)} \right) - (N-1) \right]^{1/(1-q)} \\ &< W_{A_1+A_2+\dots+A_N} = \prod_{i=1}^N W_{A_i}. \end{aligned} \quad (59)$$

In particular, for $q = 0$, we have

$$W_{A_1+A_2+\dots+A_N}^{eff} = \left(\sum_{i=1}^N W_{A_i} \right) - N + 1. \quad (60)$$

If the N subsystems are all equal, we have that

$$\ln_q W^{eff}(N) \sim N \ln_q W(1), \quad (61)$$

the changement of notation clearly being $W^{eff}(N) \equiv W_{A_1+A_2+\dots+A_N}^{eff}$ and $W(1) \equiv W_{A_1}$. Consequently, for $q = 1$, we have

$$W^{eff}(N) = W(N) = [W(1)]^N, \quad (62)$$

whereas, for $q < 1$, we have

$$\begin{aligned} W^{eff}(N) &\sim \left\{ N \left[[W(1)]^{1-q} - 1 \right] + 1 \right\}^{1/(1-q)} \\ &< W(N) = [W(1)]^N. \end{aligned} \quad (63)$$

We therefore see that, for $q = 1$, the number $W^{eff}(N)$ of nonzero-probability states grows *exponentially* with N , whereas, for $q < 1$, $W^{eff}(N)$ grows like a *power-law* with N , more precisely like $N^{1/(1-q)}$. For $q = 0$, it grows *linearly* with N , more precisely $W^{eff}(N) = N[W(1) - 1] + 1$.

2. The hint of the extensivity of S_q

We have recently argued (see [18] and references therein) that special correlations may exist between the N subsystems A_1, A_2, \dots, A_N of a composed system such that one (and only one) value of the index q exists which ensures the *additivity* of S_q . The trivial case of course is that of S_{BG} . We consider the case of *independency*, i.e., the joint probabilities given by

$$p_{i_1 i_2 \dots i_N}^{A_1 + A_2 + \dots + A_N} = \prod_{l=1}^N p_{i_l}^{A_l}, \quad \forall (i_1, i_2, \dots, i_N). \quad (64)$$

We straightforwardly verify that

$$S_{BG}(A_1 + A_2 + \dots + A_N) = \sum_{l=1}^N S_{BG}(A_l) \quad (65)$$

where

$$S_{BG}(A_1 + A_2 + \dots + A_N) \equiv - \sum_{i_1=1}^{W_{A_1}} \sum_{i_2=1}^{W_{A_2}} \dots \sum_{i_N=1}^{W_{A_N}} p_{i_1 i_2 \dots i_N}^{A_1 + A_2 + \dots + A_N} \ln p_{i_1 i_2 \dots i_N}^{A_1 + A_2 + \dots + A_N} \quad (66)$$

and

$$S_{BG}(A_l) \equiv - \sum_{i_l=1}^{W_{A_l}} p_{i_l}^{A_l} \ln p_{i_l}^{A_l}, \quad \forall l. \quad (67)$$

We can also verify

$$\ln[1 + (1-q)S_q(A_1 + A_2 + \dots + A_N)] = \sum_{l=1}^N \ln[1 + (1-q)S_q(A_l)]. \quad (68)$$

If the subsystems are all equal, we have

$$S_{BG}(N) = N S_{BG}(1), \quad (69)$$

the notation being self-explanatory.

If we consider $N = 2$ in Eq. (68) we obtain

$$S_q(A_1 + A_2) = S_q(A_1) + S_q(A_2) + (1-q)S_q(A_1)S_q(A_2). \quad (70)$$

Therefore, S_{BG} is said to be *extensive* (or *additive*). Consistently, S_q is, unless $q = 1$, *nonextensive* (or *nonadditive*). It is in fact from this property that the statistical mechanics we are talking about has been named *nonextensive*. As we shall exhibit in what follows, this early

denomination might (unfortunately) be somewhat misleading. Indeed, S_q is, for $q \neq 1$, nonextensive *if* the subsystems are (explicitly or tacitly) assumed *independent*. But, if they are specially correlated, S_q can in fact be extensive for some special value of $q \neq 1$.

Consider now the following set of joint probabilities (clearly corresponding to *nonindependent* subsystems):

$$p_{11\dots 1}^{A_1 + A_2 + \dots + A_N} = \left(\sum_{l=1}^N p_1^{A_l} \right) - N + 1, \quad (71)$$

$$p_{1\dots l_i \dots 1}^{A_1 + A_2 + \dots + A_N} = p_{i_l}^{A_l} \quad (i_l \neq 1; \forall l), \quad (72)$$

and zero otherwise. Consequently, we have only $W_{A_1 + A_2 + \dots + A_N}^{eff} = \left(\sum_{l=1}^N W_{A_l} \right) - N + 1$ joint probabilities which are generically nonzero, consistently with Eq. (60). We straightforwardly verify

$$S_0(A_1 + A_2 + \dots + A_N) = \sum_{l=1}^N S_0(A_l), \quad (73)$$

where

$$S_0(A_1 + A_2 + \dots + A_N) = W_{A_1 + A_2 + \dots + A_N}^{eff} - 1, \quad (74)$$

and

$$S_0(A_l) = W_{A_l} - 1 \quad (\forall l). \quad (75)$$

Therefore, for the particular correlations involved in Eqs. (71) and (72), it is for $q = 0$ (and *not* for $q = 1$) that S_q is additive.

Unfortunately, we do not know the general explicit form of the set of joint probabilities for an arbitrary number N of (not necessarily equal) subsystems, corresponding to $q \neq 0, 1$. We do know however, for $0 \leq q \leq 1$, some special cases, such as the $N = 2, 3$ generic *binary* subsystems. They are presented and analyzed in [18]. We also know, for binary equal subsystems, exact recurrence relations that enable the calculation of the entire set of 2^N joint probabilities associated with N subsystems given the entire set of 2^{N-1} joint probabilities associated with $N - 1$ subsystems. This calculation is rather lengthy and will be the object of a separate paper.

Now we present here (in a form which is slightly different, and possibly more transparent, than the one used in [18]) the case of $N = 2$ equal binary subsystems. Consider the set of joint probabilities indicated in Table I. Let us impose the additivity of the entropy, i.e., $S_q(2) = 2S_q(1)$, where

$$S_q(2) \equiv \frac{1 - [p^2 + \kappa]^q - 2[p(1-p) - \kappa]^q - [(1-p)^2 + \kappa]^q}{q - 1} \quad (76)$$

and

$$S_q(1) \equiv \frac{1 - p^q - (1-p)^q}{q - 1}. \quad (77)$$

The relation between the *correlation* κ , p and q is then given by

$$2[p^q + (1-p)^q] - [p^2 + \kappa]^q - 2[p(1-p) - \kappa]^q - [(1-p)^2 + \kappa]^q = 1. \quad (78)$$

(With the notation $f_q(p) \equiv p^2 + \kappa$, this relation coincides with the one indicated in [18]). In Fig. 2, typical κ versus p curves are presented. The *lower* curve corresponds to $p^2 + \kappa = 0$ and to $(1-p)^2 + \kappa = 0$. The *upper* curve corresponds to $p(1-p) - \kappa = 0$. The lower curve undoubtedly corresponds to $q = 0$. The upper curve could in principle correspond to *both* cases $q \rightarrow -\infty$ and $q \rightarrow \infty$. Indeed, for $\kappa = p(1-p)$, we have that $p^2 + \kappa = p$ and $(1-p)^2 + \kappa = 1-p$, hence $S_q(2) = S_q(1)$. Since we must also satisfy $S_q(2) = 2S_q(1)$, only two possibilities emerge a priori, and these are $S_q(1) = 0$ (which corresponds to $q \rightarrow \infty$), and $S_q(1) \rightarrow \infty$ (which corresponds to $q \rightarrow -\infty$). Monotonicity with regard to q suggests that the upper curve should correspond to $q \rightarrow \infty$. But on the other hand, having, among the four joint probabilities that are a priori possible in the present $N = 2$ system, *two* zeros (which definitively is what corresponds to the upper curve) appears as *more* restrictive than having *one* zero (which definitively is what corresponds to the lower curve). The generic *four* nonzero-probability case corresponds to $q = 1$, and the generic *three* nonzero-probability case corresponds to $q = 0$. Consequently, it appears as reasonable that the generic *two* nonzero-probability case would correspond to $q \rightarrow -\infty$. This kind of paradoxal situation appears to need further discussion to be clarified. The situation would probably be clear-cut if we had — but we do not! —, for arbitrary values of N arbitrary subsystems, the general answer to be associated with arbitrary q . This point thus remains open.

$A \setminus B$	1	2	
1	$p^2 + \kappa$	$p(1-p) - \kappa$	p
2	$p(1-p) - \kappa$	$(1-p)^2 + \kappa$	$1-p$
	p	$1-p$	1

TABLE I: Joint probabilities for two binary subsystems A and B . The marginal probabilities are indicated as well. The correlation κ and the probability p are such that all terms $p^2 + \kappa$, $p(1-p) - \kappa$, and $(1-p)^2 + \kappa$ remain within the interval $[0, 1]$. The case of independency corresponds to $\kappa = 0$.

It is worthy stressing a logical consequence of what has been said up to now. Unless $q = 1$, we *cannot* simultaneously have equal probabilities for the set of joint probabilities of the *allowed* states *and* for the associated marginal probabilities. For example, for the $q = 0$ case of two equal binary subsystems, equal probabilities for the joint set means $p_{11}^{A+B} = p_{12}^{A+B} = p_{21}^{A+B} = 1/3$ and $p_{22}^{A+B} = 0$, hence $p = 2/3$ (which differs from $1-p = 1/3$), whereas equal probabilities for the marginal sets means $p = 1/2$,




FIG. 2: Curves $\kappa(p)$ which, for typical values of q , imply additivity of S_q . For $-1/4 \leq \kappa \leq 0$ we have $\sqrt{-\kappa} \leq p \leq 1 - \sqrt{-\kappa}$. For $0 \leq \kappa \leq 1/4$ we have $(1 - \sqrt{1-4\kappa})/2 \leq p \leq (1 + \sqrt{1-4\kappa})/2$.

hence $p_{11}^{A+B} = 0 \neq p_{12}^{A+B} = p_{21}^{A+B} = 1/2$. The $q = 1$ case *simultaneously* allows for $p_{11}^{A+B} = p_{12}^{A+B} = p_{21}^{A+B} = p_{22}^{A+B} = 1/4$ and $p = 1-p = 1/2$. It is clear that, if such behavior (i.e., generic impossibility of equal probabilities for *both* joint and marginal ones simultaneously) persists up to the thermodynamic limit $N \rightarrow \infty$, it will have heavy consequences for the macroscopic statistics of the system.

Let us recall at this stage that, depending on whether the subsystems that we are composing are or are not independent, it is S_{BG} or a different entropy which is *additive*. If they are correlated in the special form that has been illustrated above, and that persists up to the thermodynamic limit $N \rightarrow \infty$, it is S_q with a specific value of $q \neq 1$ which becomes *extensive* [19]. The corresponding statistical mechanics should consistently be based on S_q (or on a directly related one, such as say S_{2-q}) and *not* anymore on S_{BG} . We may summarize this scenario through the following statement: *Unless the composition law of the subsystems is specified, the question whether an entropy (or some similar quantity) is or is not extensive has no sense*. Allow us a quick digression. The situation is totally analogous to the quick or slow motion of a body. Ancient greeks considered motion to be an absolute property. It was not until Galileo that it was clearly perceived that motion has no sense *unless the referential is specified*.

3. The hint of the q -generalization of the Pascal triangle

Very recently, Suyari and Tsukuda [20] used the concept of q -product to consistently generalize various results that are widely known in the context of BG statistical mechanics and its mathematical structure. The generalization of the $n!$ (n -factorial) operation, as well as of the binomial (and even multinomial) coefficients was performed. As a corollary, the Pascal triangle itself was q -generalized as well. It is known that the coefficients of the n -th line of the Pascal triangle yield (after appropriate centralization and rescaling) the Gaussian distribution in the $n \rightarrow \infty$ limit. This is well known to be a simple consequence of the standard central limit theorem as applied to independent binary variables. Suyari claims that, in the $n \rightarrow \infty$ limit of the q -generalized Pascal triangle, what is obtained is precisely the q -Gaussian distributions!

4. The hint of the scale-free networks

The mathematical study of *random networks* (or *random graphs*) started many decades ago. But quite recently, it has acquired great interest due to the fact that such structures appear to be ubiquitous in physical, social, internet and other complex phenomena (see [21, 22] and references therein). A central quantity of such structures is the so-called *connectivity* or *degree distribution*, defined as the probability distribution of the number of links that are connected to the same site (or node); being more explicit, what one counts is the percentage of nodes that have a given number of links. For the important class of networks that are referred to as *scale-free* ones, this distribution is systematically found to be precisely a q -exponential. Many examples do exist in the literature. Three recent illustrations can be found in [23, 24, 25].

These scale-invariant structures exhibit *hubs* and *sub-hubs*, and so on. If embedded into a d -dimensional space, they tend to have *zero* Lebesgue measure. They appear to be like fractals, which implies that most of the locations of the d -dimensional space are forbidden (like the flights of any air company are expected to start and end *only* at the airports where that company operates, and not in any place of the territory!).

5. The nonlinear dynamical hint

We review at this point a few nonlinear dynamical results which, although not directly related to the possible *third* central limit theorem we are seeking, provide what we consider important connections within the mathematical structure which sustains nonextensive statistical mechanics.

We focus on classical nonlinear dynamical systems, either conservative (e.g., Hamiltonian systems) or dissipative. The basic ideas are quite general, but their presen-

tation becomes easier if we illustrate them on a simple, one-dimensional, system. Let us address unimodal one-dimensional maps, such as for example the z -logistic map defined through

$$x_{t+1} = 1 - a|x_t|^z \quad (t = 0, 1, 2, \dots; \quad z \geq 1; \quad 0 \leq a \leq 2; \quad -1 \leq x_t \leq 1) \quad (79)$$

The *sensitivity* ξ to the initial conditions is defined as follows:

$$\xi(t) \equiv \lim_{\Delta x(0) \rightarrow 0} \frac{\Delta x(t)}{\Delta x(0)}, \quad (80)$$

where $\Delta x(t)$ is the discrepancy at time t of two values of x_0 initially separated by $\Delta x(0)$. The typical behavior of ξ is given by

$$\xi(t) = e^{\lambda_1 t}, \quad (81)$$

where λ_1 is referred to as the *Lyapunov exponent* (the subindex 1 will become clear in a few lines). The following ordinary differential equation is satisfied: $d\xi/dt = \lambda_1 \xi$.

For most values of the *control parameter* a (see Eq. (79)), λ_1 is nonzero. When it is positive, we shall say that there is *strong chaos*. When it is negative, we have *regular orbits* as attractors (e.g., fixed points, cycles-2, cycles-3, cycles-4, etc). But there is an infinity of values of a for which λ_1 vanishes. Examples are (i) all the values of a for which there is a *doubling-period bifurcation* from one cycle to its double; (ii) all the values of a for which there is a *tangent bifurcation*; (iii) all the values of a for which there is an *edge to chaos*. All these cases are interesting, but by far the richest one is case (iii), referred to as *weak chaos*. Indeed, it is there that we expect the appearance of complexity (in physics, biology, economics, linguistics, and elsewhere), since it is the frontier between considerable order (regular orbits) and considerable disorder (strong chaos). So, the following question arises: *what is the behavior of $\xi(t)$ when λ_1 vanishes?* The typical behavior is as follows:

$$\xi(t) = e_q^{\lambda_q t}, \quad (82)$$

where we have $q > 1$ (with $\lambda_q > 0$) for the above cases (i) and (ii), and $q < 1$ (with $\lambda_q > 0$) for the above case (iii). The following ordinary differential equation is satisfied: $d\xi/dt = \lambda_q \xi^q$. The behavior (82) was conjectured in 1997 [26] and was proved in 2003 [27]. For example, for $z = 2$, the first edge of chaos appears at $a = a_c \equiv 1.4011\dots$. Its associated values are $q_{sen} = 0.2445\dots$ and $\lambda_{q_{sen}} = 1/(1 - q_{sen}) = 1.3236\dots$, where the subscript *sen* stands for *sensitivity*.

Let us now address another very important quantity, namely the *entropy production* K per unit time. It is basically defined as follows. Take the admissible phase space (the interval $[-1, 1]$ for the variable x in the map (79), for instance), and make a partition of it in W little parts, identified through $i = 1, 2, \dots, W$. Then consider

N_0 initial values of x_0 within one of the W intervals. As a function of time, the points within the initial window will possibly spread around, in such a way that we have $N_i(t)$ points in the i -th interval ($\sum_{i=1}^W N_i(t) = N_0$). We define next the set of probabilities $p_i \equiv N_i(t)/N_0$ ($\forall i$), and finally we calculate the entropy $S_q(t)$ through Eq. (9) for any chosen value of q . By construction we have $S_q(0) = 0$. We then define the entropy production per unit time as follows

$$K_q \equiv \lim_{t \rightarrow \infty} \lim_{W \rightarrow \infty} \lim_{N_0 \rightarrow \infty} \frac{S_q(t)}{t}. \quad (83)$$

The rate K_1 coincides, in most cases, with the so called Kolmogorov-Sinai entropy rate. The rate K_q represents its q -generalization. Consistently with the Pesin theorem, we expect, whenever $\lambda_1 \geq 0$,

$$K_1 = \lambda_1. \quad (84)$$

This is indeed verified (see, for instance, [28]). For example, for $(z, a) = (2, 2)$ in (79), we obtain $K_1 = \lambda_1 = \ln 2$. What happens however at the edge of chaos (say at $a_c(z)$)? Can we state something more informative than just $K_1 = \lambda_1 = 0$? *Yes, we can.* It can be numerically shown [29] and analytically proved [30] that

$$K_{q_{sen}} = \lambda_{q_{sen}}. \quad (85)$$

This interesting relation (which refers in fact to *upper bounds*: see details in [30]) basically generalizes the celebrated Pesin theorem, and was also conjectured in 1997 [26]. To be more specific, what happens is that there is a value of q (and only one), noted q_{sen} , such that $K_q = 0$ for $q > q_{sen}$, and $K_q \rightarrow \infty$ for $q < q_{sen}$. And for $q = q_{sen}$ we obtain a finite value $K_{q_{sen}}$ which precisely coincides with $\lambda_{q_{sen}}$. When $\lambda_1 > 0$, $q_{sen} = 1$ (strong chaos); when $\lambda_1 = 0$ and the system is at the edge of chaos, $q_{sen} < 1$ and generically $\lambda_{q_{sen}} > 0$ (weak chaos). For example, for the edge of chaos of the universality class to which the map (79) belongs, q_{sen} increases from $-\infty$ to slightly below unity when z increases from 1 to ∞ .

What happens when our nonlinear dynamical system has more than one, say μ nonnegative Lyapunov exponents $\{\lambda_1^{(m)}\}$ ($m = 1, 2, \dots, \mu$)? From the Pesin theorem, we certainly expect

$$K_1 = \sum_{m=1}^{\mu} \lambda_1^{(m)}. \quad (86)$$

What happens then if the system is at an edge of chaos, where $\lambda_1^{(m)} = 0, \forall m$? We expect [31] the following (conjectural) behavior for the Lebesgue measure ξ associated with the dynamically expanding directions:

$$\xi \simeq \prod_{m=1}^{\mu} \xi^{(m)} = \prod_{m=1}^{\mu} \exp_{q_{sen}^{(m)}}(\lambda_{q_{sen}^{(m)}}^{(m)} t), \quad (87)$$

where the (\simeq) sign might become just ($=$) under some simplifying hypothesis (like orthogonality of the directions along which the expansions associated with the $\lambda_1^{(m)}$'s occur).

Since we essentially expect, for $t \rightarrow \infty$,

$$S_{q_e} \sim \ln_{q_e} \xi, \quad (88)$$

(the subscript e stands for *entropy*), we possibly have the following relation:

$$S_{q_e} \sim \ln_{q_e} \left[\prod_{m=1}^{\mu} \exp_{q_{sen}^{(m)}}(\lambda_{q_{sen}^{(m)}}^{(m)} t) \right]. \quad (89)$$

Using the definition (83) we obtain the following interesting relation:

$$K_{q_e} = \lim_{t \rightarrow \infty} \frac{\ln_{q_e} \left[\prod_{m=1}^{\mu} \exp_{q_{sen}^{(m)}}(\lambda_{q_{sen}^{(m)}}^{(m)} t) \right]}{t}. \quad (90)$$

Two important cases must be distinguished, namely strong and weak chaos. Strong chaos corresponds to $\lambda_1^{(m)} > 0, \forall m$. In this case, we have $q_e = q_{sen}^{(m)} = 1, \forall m$, hence Eq. (90) straightforwardly recovers relation (86), consistently with the Pesin theorem. Weak chaos corresponds to $\lambda_1^{(m)} = 0, \forall m$. In this case, by focusing on the $t \rightarrow \infty$ asymptotic region, we have

$$K_{q_e} = \frac{\prod_{m=1}^{\mu} \left[(1 - q_{sen}^{(m)}) \lambda_{q_{sen}^{(m)}}^{(m)} \right]^{\frac{1 - q_e}{1 - q_{sen}^{(m)}}}}{1 - q_e} \times \lim_{t \rightarrow \infty} \frac{t^{(1 - q_e) \sum_{m=1}^{\mu} \frac{1}{1 - q_{sen}^{(m)}}}}{t}, \quad (91)$$

consequently (since q_e must be chosen so that K_{q_e} is finite)

$$\frac{1}{1 - q_e} = \sum_{m=1}^{\mu} \frac{1}{1 - q_{sen}^{(m)}}, \quad (92)$$

and

$$K_{q_e} = \frac{\prod_{m=1}^{\mu} \left[(1 - q_{sen}^{(m)}) \lambda_{q_{sen}^{(m)}}^{(m)} \right]^{\frac{1 - q_e}{1 - q_{sen}^{(m)}}}}{1 - q_e}. \quad (93)$$

Eq. (93) can be rewritten in a more symmetric form, namely

$$[(1 - q_e) K_{q_e}]^{\frac{1}{1 - q_e}} = \prod_{m=1}^{\mu} \left[(1 - q_{sen}^{(m)}) \lambda_{q_{sen}^{(m)}}^{(m)} \right]^{\frac{1}{1 - q_{sen}^{(m)}}}. \quad (94)$$

If $\mu = 1$, Eq. (92) recovers $q_e = q_{sen}$, and Eq. (94) recovers Eq. (85).

Another interesting particular case is when $q_{sen}^{(m)} = q_{sen} (\forall m)$, and $\lambda_{q_{sen}^{(m)}}^{(m)} = \lambda_{q_{sen}} (\forall m)$. From Eqs. (92) and (94) it follows then

$$1 - q_e = \frac{1 - q_{sen}}{\mu} \quad (95)$$

and

$$K_{q_e} = \mu \lambda_{q_{sen}}. \quad (96)$$

An example which verifies Eq. (95) can be found in [31].

One more particular case which is interesting is when the expansion is *linear* in time for *all* directions, i.e., $q_{sen}^{(m)} = 0$ ($\forall m$). We then have, from Eqs. (92) and (94),

$$q_e = 1 - \frac{1}{\mu} \quad (97)$$

and

$$K_{q_e} = \mu \left[\prod_{m=1}^{\mu} \lambda_0^{(m)} \right]^{1/\mu}. \quad (98)$$

Relation (97) has already emerged in the literature through various forms. A first example concerns $\mu = 1$, hence we expect $q_e = 0$. This is precisely what can be verified [32] for the Casati-Prosen triangle map [33], which is a two-dimensional, conservative (hence $\mu = 1$), mixing, ergodic one, with *linear* instability. In addition to $q_e = 0$, it has been verified that $K_e = \lambda_0$, in accordance with Eq. (98). A second example is the μ -dimensional lattice Lotka-Volterra, for which it has precisely been verified [34, 35] the result (97). A third example is a specific d -dimensional Boltzmann lattice model [36]. Its Hamiltonian-like behavior leads to $\mu = d/2$, which, from Eq. (97), implies $q_e = 1 - 2/d$. It is precisely this result that the authors [36] have obtained by imposing Galilean invariance to the dynamical equations of their model.

It is clear that the interesting (and possibly exact) relations (92) and (94) remain to be proved. At the present stage they constitute but conjectures.

6. The empirical hint

Last but not least, let us present a very pragmatic, epistemological-like, reason. There are, in the literature, already so many natural and artificial systems (and their number constantly increases) whose central quantities are well fitted by q -exponentials and q -Gaussians, that one is compelled to believe that only a limit theorem could explain such an ubiquity.

V. CONCLUSIONS

In this final section, let us summarize the scenario within which we are working. It appears that the entropic form $S(N; \{p_i\})$ to be used for constructing a statistical mechanics that is naturally compatible with usual macroscopic thermodynamics should be *generically extensive*, i.e., such that

$$0 \leq \lim_{N \rightarrow \infty} \frac{S(N; \{p_i\})}{N} < \infty, \quad (99)$$

the equality occurring only for the case of certainty or (thermodynamically) close to it (the typical example is a system in thermal equilibrium at *zero temperature*). Within this (axiomatic) viewpoint, we can distinguish the following cases:

(i) The N subsystems (typically physical elements such as particles free to move translationally and/or rotationally, localized spins, and similar entities) are (probabilistically) *independent* (the paradigmatic case is that of ideal gases; their total energy is strictly proportional to N , i.e., additive, hence *extensive*). Then we must use S_{BG} since $S_{BG}(N; \{p_i\}) \propto N$ ($\forall \{p_i\}$).

(ii) The N subsystems are *locally correlated* (the paradigmatic case is that of *short-range-interacting* many-body Hamiltonian systems; their total energy is asymptotically proportional to N , i.e., it is *extensive* once again). Then, once again, we must use S_{BG} since it satisfies Eq. (99), $\forall \{p_i\}$.

(iii) The N subsystems are *globally correlated* in the special manner addressed in this paper (the paradigmatic case appears to be that of *long-range-interacting* many-body Hamiltonian systems; their total energy asymptotically increases with N faster than linearly, i.e., it is *nonextensive*). Then, we must use S_q for that unique value of q which guarantees Eq. (99), $\forall \{p_i\}$.

(iv) The N subsystems are *globally correlated* in a manner which is more complex (or just as complex but in a different manner) than the one addressed in this paper. Then we must use an entropy which is *not* included in the family S_q for any value of q . Such an entropy would have to be either more general than S_q (see for instance [37] for the so called *superstatistics* [38]), or just of a completely different type (see for instance [39, 40, 41]).

The cases (i) and (ii), which we may call *simple* in the sense of *plectics* [42], belong to the world within which the concepts used in *BG* statistical mechanics have, since more than one century, been profusely shown to be the appropriate ones.

The cases (iii) and (iv), which we may call *complex* in the sense of *plectics* [42], belong to the world within which the concepts used in nonextensive statistical mechanics (as well as in its possible generalizations or alternatives) have been shown and keep being shown (since more than one decade, by now) to be the appropriate ones.

The cause for a system to be *simple* or *complex* in the above sense lies basically (see [3, 43]) on its microscopic dynamics in the *full* space of microscopic possibilities (Gibbs' Γ -space for many-body Hamiltonian systems). It is believed to be so because it is this dynamics which is expected to determine the possible persistent correlations that would be responsible for the geometrical structure within which the system tends to “live”, given its initial conditions.

If its elementary nonlinear dynamics is controlled by *strong* sensitivity to the initial conditions (i.e., at least one *positive* Lyapunov exponent if the system is a classical one), we expect the system to be *simple*. For virtually any initial condition, the system quickly visits the

neighborhood of virtually all possible states; it does so in such a way that the probability of any small part of the (macroscopically) admissible full space asymptotically becomes proportional to its size (Lebesgue measure). Its main time-dependent functions (typically) *exponentially* depend on time, the number of allowed states *exponentially* increases with N , and, if the system is Hamiltonian, its stationary state (called *thermal equilibrium*) is characterized by an energy distribution given by the celebrated *BG* weight. Summarizing, its paradigmatic ordinary differential equation is $dy/dx \propto y$.

If the elementary nonlinear dynamics of the system is nonregular and controlled by *weak* sensitivity to the initial conditions (i.e., *no positive* Lyapunov exponents if the system is a classical one), we expect the system to be *complex*. For given initial conditions, the system essentially visits a network of states (a *scale-free network* for many if not all q -systems) whose typical Lebesgue measure is zero. The particular network depends from the initial conditions and is highly inhomogeneous (like, as mentioned before, the network of airports on which a specific air company operates), but its geometry (both topology and metrics) is basically the same for virtually all initial conditions. To recover a homogeneous occupancy of the full space we are obliged to make *averages over all the initial conditions* (whereas no such thing is necessary for *simple* systems). The main time-dependent functions of the system typically depend on time slower than exponentially (typically like *power-laws*, more precisely like q -exponentials for most if not all the q -systems), the number of allowed states increases with

N like a power-law, and, if the system is Hamiltonian, its stationary or quasi-stationary (metastable) state (out of thermal equilibrium) is expected to be characterized by an energy distribution given by a q -exponential weight, which asymptotically approaches a power-law for high energies. Summarizing, its paradigmatic ordinary differential equation is $dy/dx \propto y^q$.

The standard central limit theorem plays a crucial role for the *simple* systems. We expect a similar theorem to exist for the *complex* systems of the q -class. The proof of such a theorem would be priceless.

In addition to the above, we have presented here more two conjectures (Eqs. (92) and (94)) concerning the entropy production per unit time for a nonlinear dynamical system at the edge of chaos, having μ vanishing Lyapunov exponents. The entropy S_q which increases *linearly* with time (when the system is exploring its phase space) appears to be that whose entropic index is q_e as given by Eq. (92) [31]. The associated entropy production K_{q_e} per unit time appears to be as given by Eq. (94). The proof of these two connected conjectures (including, naturally, the precise conditions for their validity) also remains as a mathematically open problem.

Acknowledgments

We acknowledge useful discussions with S. Abe, C. Anteneodo, F. Baldovin, S. Curilef, M. Gell-Mann, R. Hersh, L. Moyano, C. Pagani and Y. Sato, as well as partial financial support by Pronex/MCT, Faperj and CNPq (Brazilian agencies).

-
- [1] L. Boltzmann, *Weitere Studien über das Wärme Gleichgewicht unter Gas molekülen* [Further Studies on Thermal Equilibrium Between Gas Molecules], Wien, Ber. **66**, 275 (1872),
 - [2] J.W. Gibbs, *Elementary Principles in Statistical Mechanics - Developed with Especial Reference to the Rational Foundation of Thermodynamics* (C. Scribner's Sons, New York, 1902; Yale University Press, New Haven, 1948; OX Bow Press, Woodbridge, Connecticut, 1981). In page 35: "In treating of the canonical distribution, we shall always suppose the multiple integral in equation (92) [the partition function, as we call it nowadays] to have a finite value, as otherwise the coefficient of probability vanishes, and the law of distribution becomes illusory. This will exclude certain cases, but not such apparently, as will affect the value of our results with respect to their bearing on thermodynamics. It will exclude, for instance, cases in which the system or parts of it can be distributed in unlimited space [...]. It also excludes many cases in which the energy can decrease without limit, as when the system contains material points which attract one another inversely as the squares of their distances. [...]. For the purposes of a general discussion, it is sufficient to call attention to the assumption implicitly involved in the formula (92)."
 - [3] A. Einstein, *Theorie der Opaleszenz von homogenen Flüssigkeiten und Flüssigkeitsgemischen in der Nähe des kritischen Zustandes*, Annalen der Physik **33**, 1275 (1910). ["Usually W is put equal to the number of complexions... In order to calculate W , one needs a *complete* (molecular-mechanical) theory of the system under consideration. Therefore it is dubious whether the Boltzmann principle has any meaning without a complete molecular-mechanical theory or some other theory which describes the elementary processes. $S = \frac{R}{N} \log W + \text{const.}$ seems without content, from a phenomenological point of view, without giving in addition such an *Elementartheorie*." (Translation: Abraham Pais, *Subtle is the Lord...*, Oxford University Press, 1982)].
 - [4] M. Gell-Mann and C. Tsallis, eds., *Nonextensive Entropy - Interdisciplinary Applications* (Oxford University Press, New York, 2004); H.L. Swinney and C. Tsallis, eds., *Anomalous Distributions, Nonlinear Dynamics and Nonextensivity*, Physica D **193** (2004).
 - [5] C. Tsallis, *Possible generalization of Boltzmann-Gibbs statistics*, J. Stat. Phys. **52**, 479 (1988); for full bibliography see <http://tsallis.cat.cbpf.br/biblio.htm>.
 - [6] E.M.F. Curado and C. Tsallis, *Generalized statistical mechanics: connection with thermodynamics*, J. Phys. A **24**, L69 (1991); Corrigenda: **24**, 3187 (1991) and **25**, 1019 (1992).
 - [7] C. Tsallis, R.S. Mendes and A.R. Plastino, *The role*

of constraints within generalized nonextensive statistics, *Physica A* **261**, 534 (1998).

- [8] F. Takens, in *Structures in Dynamics - Finite Dimensional Deterministic Studies*, eds. H.W. Broer, F. Dumortier, S.J. van Strien and F. Takens, (North-Holland, Amsterdam, 1991). In page 253: "The values of p_i are determined by the following *dogma* : if the energy of the system in the i^{th} state is E_i and if the temperature of the system is T then: $p_i = e^{-E_i/kT}/Z(T)$, where $Z(T) = \sum_i e^{-E_i/kT}/Z(T)$, (this last constant is taken so that $\sum_i p_i = 1$). This choice of p_i is called the *Gibbs distribution*. We shall give no justification for this dogma; even a physicist like Ruelle disposes of this question as "deep and incompletely clarified"."
- [9] C. Tsallis, *What are the numbers that experiments provide?*, *Quimica Nova* **17**, 468 (1994).
- [10] D. Prato and C. Tsallis, *Nonextensive foundation of Lévy distributions*, *Phys. Rev. E* **60**, 2398 (1999).
- [11] A.M.C. de Souza and C. Tsallis, *Student's t- and r-distributions: Unified derivation from an entropic variational principle*, *Physica A* **236**, 52 (1997).
- [12] A. Einstein, *Über die von der molekulär kinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen*, *Annalen der Physik* **17**, 549 (1905).
- [13] A.R. Plastino and A. Plastino, *Non-extensive statistical mechanics and generalized Fokker-Planck equation*, *Physica A* **222**, 347 (1995).
- [14] C. Tsallis and D.J. Bukman, *Anomalous diffusion in the presence of external forces: exact time-dependent solutions and their thermostistical basis*, *Phys. Rev. E* **54**, R2197 (1996).
- [15] E.M.F. Curado and F.D. Nobre, *Derivation of nonlinear Fokker-Planck equations by means of approximations to the master equation*, *Phys. Rev. E* **67**, 021107 (2003); F.D. Nobre, E.M.F. Curado and G. Rowlands, *A procedure for obtaining general nonlinear Fokker-Planck equations*, *Physica A* **334**, 109 (2004).
- [16] L. Nivanen, A. Le Mehaute and Q.A. Wang, *Generalized algebra within a nonextensive statistics*, *Rep. Math. Phys.* **52**, 437 (2003); E.P. Borges, *A possible deformed algebra and calculus inspired in nonextensive thermostistics*, *Physica A* **340**, 95 (2004).
- [17] A simple generalization also covering possibly negative real numbers X and Y is the following one: $X \otimes_q Y \equiv [\text{sign}(X)][\text{sign}(Y)][|X|^{1-q} + |Y|^{1-q} - 1]^{1/(1-q)}$.
- [18] C. Tsallis, *Is the entropy S_q extensive or nonextensive?*, in *Complexity, Metastability and Nonextensivity*, eds. C. Beck, A. Rapisarda and C. Tsallis (World Scientific, Singapore, 2005), to appear [cond-mat/0409631]; Y. Sato and C. Tsallis, *On the extensivity of the entropy S_q for systems with $N \leq 3$ specially correlated subsystems*, *International Journal of Bifurcation and Chaos*, to appear: *Proceedings of the Summer School and Conference on Complexity in Science and Society*, ed. T. Bountis (Patras and Olympia, 14-26 July, 2004) [cond-mat/0411073].
- [19] To avoid ambiguities let us recall that an entropy $S(N)$ of a system composed of N subsystems is said *additive* if, under a given composition law for the N subsystems, $S(N) \propto N$. It is said *extensive* if $\lim_{N \rightarrow \infty} S(N)/N$ is finite. Consequently, if an entropy is additive, it also is extensive. The opposite is not true. The usual thermodynamic requirement for entropy is to be *extensive* (i.e., asymptotically additive), not necessarily *additive* (i.e., strictly additive). So, if the subsystems are independent, we must use S_{BG} since it is additive, hence extensive. If the subsystems are *locally* correlated (the paradigmatic case is that of *short-range-interacting* many-body Hamiltonian systems), we must also use S_{BG} since, although not being strictly additive, it is extensive. If the subsystems are *globally* correlated (the paradigmatic case appears to be that of *long-range-interacting* many-body Hamiltonian systems), then we must use another entropy as the basis for constructing a statistical mechanics. If this global correlation is of the type focused on in this paper, then S_q with the appropriate value of q is extensive, and is the one to be used. If the global correlation is of some different nature, even more complex than the one focused on in this paper, a still different entropic form might be necessary.
- [20] H. Suyari and M. Tsukada, *Law of error in Tsallis statistics*, *IEEE Transaction on Information Theory* (2004), in press [cond-mat/0401540]; H. Suyari, *q-Stirling's formula in Tsallis statistics*, cond-mat/0401541; H. Suyari, *Mathematical structure derived from the q-multinomial coefficient in Tsallis statistics*, cond-mat/0401546.
- [21] D.J. Watts and S.H. Strogatz, *Collective dynamics of small-world networks*, *Nature* **393**, 440 (1998).
- [22] R. Albert and A.-L. Barabasi, *Statistical mechanics of complex networks*, *Rev. Mod. Phys.* **74**, 47 (2002).
- [23] R. Albert and A.-L. Barabasi, *Topology of evolving networks: Local events and universality*, *Phys. Rev. Lett.* **85**, 5234 (2000); the analytically exact distribution which is found in this paper is, although written in a slightly different form, a q -exponential with $q = \frac{2m(2-r)+1-p-r}{m(3-2r)+1-p-r}$, where (m, p, r) are parameters of the specific growth model.
- [24] J.P.K. Doye, *Network topology of a potential energy landscape: A static scale-free network*, *Phys. Rev. Lett.* **88**, 238701 (2002); the distributions that are presented in this paper can be shown to be numerically very close to a q -exponential with $q \simeq 2$.
- [25] D.J.B. Soares, C. Tsallis, A.M. Mariz and L.R. da Silva, *Preferential attachment growth model and nonextensive statistical mechanics*, cond-mat/0410459.
- [26] C. Tsallis, A.R. Plastino and W.-M. Zheng, *Power-law sensitivity to initial conditions - New entropic representation*, *Chaos, Solitons and Fractals* **8**, 885 (1997).
- [27] F. Baldovin and A. Robledo, *Sensitivity to initial conditions at bifurcations in one-dimensional nonlinear maps: Rigorous nonextensive solutions*, *Europhys. Lett.* **60**, 518 (2002); F. Baldovin and A. Robledo, *Universal renormalization-group dynamics at the onset of chaos in logistic maps and nonextensive statistical mechanics*, *Phys. Rev. E* **66**, R045104 (2002).
- [28] V. Latora and M. Baranger, *Kolmogorov-Sinai entropy rate versus physical entropy*, *Phys. Rev. Lett.* **82**, 520 (1999).
- [29] V. Latora, M. Baranger, A. Rapisarda and C. Tsallis, *The rate of entropy increase at the edge of chaos*, *Phys. Lett. A* **273**, 97 (2000).
- [30] F. Baldovin and A. Robledo, *Nonextensive Pesin identity. Exact renormalization group analytical results for the dynamics at the edge of chaos of the logistic map.*, *Phys. Rev. E* **69**, 045202(R) (2004).
- [31] G.F.J. Ananos, F. Baldovin and C. Tsallis, *Anomalous sensitivity to initial conditions and entropy pro-*

- duction in standard maps: *Nonextensive approach*, cond-mat/0403656.
- [32] G. Casati, C. Tsallis and F. Baldovin, *Linear instability and statistical laws of physics*, in preparation.
 - [33] G. Casati and T. Prosen, *Mixing property of triangular billiards*, Phys. Rev. Lett. **83**, 4729 (1999); G. Casati and T. Prosen, *Triangle map: A model of quantum chaos*, Phys. Rev. Lett. **85**, 4261 (2000).
 - [34] G.A. Tsekouras, A. Provata and C. Tsallis, *Nonextensivity of the cyclic lattice Lotka Volterra model*, Phys. Rev. E **69**, 016120 (2004).
 - [35] C. Anteneodo, *Entropy production in the cyclic lattice Lotka-Volterra model*, Eur. Phys. J. B (2004), in press [cond-mat/0402248].
 - [36] B.M. Boghosian, P.J. Love, P.V. Coveney, I.V. Karlin, S. Succi and J. Yepez, *Galilean-invariant lattice-Boltzmann models with H-theorem*, Phys. Rev. E **68**, 025103(R) (2003).
 - [37] C. Tsallis and A.M.C. Souza, *Constructing a statistical mechanics for Beck-Cohen superstatistics*, Phys. Rev. E **67**, 026106 (2003); see also A.M.C. Souza and C. Tsallis, *Stability of the entropy for superstatistics*, Phys. Lett. A **319**, 273 (2003); A.M.C. Souza and C. Tsallis, *Stability analysis of the entropy for superstatistics*, Physica A **342**, 132 (2004).
 - [38] C. Beck and E.G.D. Cohen, *Superstatistics*, Physica A **322**, 267 (2003); see also C. Beck, *Superstatistics: Theory and Applications*, in *Nonadditive entropy and nonextensive statistical mechanics*, ed. M. Sugiyama, Continuum Mechanics and Thermodynamics **16**, 293 (Springer-Verlag, Heidelberg, 2004); E.G.D. Cohen, *Superstatistics*, in *Anomalous Distributions, Nonlinear Dynamics and Nonextensivity*, eds. H.L. Swinney and C. Tsallis, Physica D **193**, 35 (2004).
 - [39] E.M.F. Curado, *General aspects of the thermodynamical formalism*, in *Nonextensive Statistical Mechanics and Thermodynamics*, eds. S.R.A. Salinas and C. Tsallis, Braz. J. Phys. **29**, 36 (1999); E.M.F. Curado and F.D. Nobre, *On the stability of analytic entropic forms*, Physica A **335**, 94 (2004).
 - [40] C. Anteneodo and A.R. Plastino, *Maximum entropy approach to stretched exponential probability distributions*, J. Phys. A **32**, 1089 (1999).
 - [41] G. Kaniadakis, *Nonlinear kinetics underlying generalized statistics*, Physica A **296**, 405 (2001); G. Kaniadakis, *Statistical mechanics in the context of special relativity*, Phys. Rev. E **66**, 0561125 (2002); G. Kaniadakis, M. Lissia and A.M. Scarfone, *Deformed logarithms and entropies*, Physica A **340**, 41 (2004).
 - [42] M. Gell-Mann, *The Quark and the Jaguar - Adventures in the Simple and the Complex*, (W.H. Freeman, New York, 1994). *Plectics* is the denomination suggested by M. Gell-Mann for the emerging science of the so called complex systems.
 - [43] E.G.D. Cohen, *Boltzmann and Einstein: Statistics and dynamics – An unsolved problem*, Boltzmann Award Lecture at STATPHYS 2004, *22nd IUPAP International Conference on Statistical Physics* (Bangalore, 4 to 9 July 2004), Pramana (2004), in press.