

Exact dynamics of a reaction-diffusion model with spatially alternating rates

M. M. Obidia, B. Schmittmann and R. K. P. Zia¹¹Center for Stochastic Processes in Science and Engineering,
Department of Physics, Virginia Tech, Blacksburg, VA, 24061-0435, USA
(Dated: February 24, 2005)

We present the exact solution for the full dynamics of a nonequilibrium spin chain and its dual reaction-diffusion model, for arbitrary initial conditions. The spin chain is driven out of equilibrium by coupling alternating spins to two thermal baths at different temperatures. In the reaction-diffusion model, this translates into spatially alternating rates for particle creation and annihilation, and even negative "temperatures" have a perfectly natural interpretation. Observables of interest include the magnetization, the particle density, and all correlation functions for both models. Two generic types of time-dependence are found: if both temperatures are positive, the magnetization, density and correlation functions decay exponentially to their steady-state values. In contrast, if one of the temperatures is negative, damped oscillations are observed in all quantities. They can be traced to a subtle competition of pair creation and annihilation on the two sublattices. We comment on the limitations of mean-field theory and propose an experimental realization of our model in certain conjugated polymers and linear chain compounds.

PACS numbers: 02.50.-r, 75.10.-b, 05.50.+q, 05.70.Ln

I. INTRODUCTION

Nonequilibrium many-body systems abound in the physical and life sciences and have recently received much attention (see e.g. [1, 2, 3] and references therein). Despite these efforts, a comprehensive theoretical framework is still lacking: As yet, there is no equivalent of Gibbs ensemble theory for nonequilibrium systems. As a consequence, in contrast to equilibrium statistical mechanics, macroscopic observables cannot be computed without explicit reference to the imposed dynamics, generally described by a master equation, and most progress in the field is made by studying paradigmatic models [2]. In this context, exact solutions of simple models are scarce, but very precious, since they can serve as testing grounds for approximate and/or numerical schemes and shed light on general properties of whole classes of related models. Not surprisingly, nontrivial solutions are almost entirely restricted to one dimension (1D; see e.g. [2, 3]), and have focused on completely uniform lattices with site-independent rates. Clearly, however, one would like to take into account more complex situations, e.g., those with spatially varying coupling constants or rates. Arguably, one of the simplest generalizations beyond a completely uniform system is one with alternating rates. In the following, we consider a 1D kinetic Ising chain (KISC), coupled to two alternating temperatures and endowed with Glauber-like dynamics. Our analysis of this model provides a full description of its dual counterpart, namely a reaction-diffusion system (RDS), characterized by spatially alternating annihilation and creation rates. Members of these two classes { i.e., kinetic Ising and reaction-diffusion models } are prototypical nonequilibrium

systems which have been thoroughly studied on homogeneous lattices [2, 3, 4, 5, 6, 7]. Yet, they still offer surprises and novel behaviors, when non-trivial spatial rates are investigated.

Our model was first introduced by Racz and Zia [8] who recognized that (stationary) two-point correlation functions are easily found exactly, even though spins on alternating sites are coupled to different temperatures. Schmittmann and Schuster subsequently realized that all stationary correlation functions are exactly calculable [9]. While this information is equivalent to the full stationary solution, its representation as $\exp(-H_e)$ is non-trivial, involving a proliferation of longer-ranged multi-spin couplings [10]. Finally, we recently reported the exact solution for all dynamic correlation functions, starting from a very simple initial condition, i.e., zero magnetization and vanishing correlations [11].

In this article, we complete these earlier studies by demonstrating how competing site-dependent rates may dramatically affect the dynamics by giving rise to an oscillatory approach toward the nonequilibrium steady state. We use a generating functional approach to obtain the complete solution for all correlation functions with arbitrary initial conditions. We focus specifically on the dynamic magnetization and the spin-spin correlations and explore their long-time behavior. We will also consider the dynamics of domain walls in the spin chain which can be mapped onto a reaction-diffusion system. Interpreting our results in the language of particle annihilation and creation, negative "temperatures" acquire a natural physical meaning, leading to unexpected oscillatory dynamics. From a more technical point of view, we are able to obtain a complete solution for two nontrivial nonequilibrium many-body systems which provides some insight into the solvability of two whole classes of related models.

The mapping to a reaction-diffusion system is of inter-

Electronic address: m.m.obidia, schmittm, rkpzia @ vt.edu

est for two reasons. On the theoretical side, the equations for densities and correlation functions in the RDS form an infinite hierarchy whose solution is not at all apparent until one recognizes the equivalent spin chain model. Also, from an experimental perspective, it is well known that diffusion-limited reactions with annihilation and creation of pairs of particles are good models for the photogrowth properties of excited states (solitons/antisoliton pairs) in certain conjugated polymers and linear chain compounds [12, 13, 14]. We propose that spatially alternating creation/annihilation rates in these systems (especially in MX chain compounds) can be generated with the help of a laser with spatially modulated power output.

This article is organized as follows: In the next section we introduce the kinetic spin chain and its dual reaction-diffusion system. Section III presents the complete solution of the spin chain. Some technical details are relegated to two Appendices. In Section IV, we map the two-temperature spin chain onto a reaction-diffusion system with alternating rates, whose density and correlation functions are computed. We analyze the conditions under which damped oscillations characterize the approach to the steady state, and we compare our exact results to a simple mean-field description. Section V is devoted to a brief discussion of the solvability of related models, with Section VI reserved for our conclusions.

II. THE MODELS

We consider two closely related nonequilibrium many-particle systems on a one-dimensional lattice: (i) a kinetic Ising spin chain (KISC) endowed with a generalized Glauber-like dynamics; and (ii) a reaction-diffusion system (RDS), with spatially periodic pair annihilation and creation rates. For convenience, we restrict ourselves to a periodic lattice (a ring) with an even number of sites and study the thermodynamic limit. We expect our exact results to be valid for the general cases of odd number of sites and/or arbitrary boundary conditions, apart from the usual caveats.

Since the RDS follows from the spin chain via a duality relationship, we focus mainly on the detailed description of model (i). A spin variable, $s_j = \pm 1$, denotes the value of the spin at site j , with $j = 1; 2; \dots; L$, and L an even integer. Nearest-neighbor spins interact according to the usual Ising Hamiltonian: $H = -J \sum_j s_j s_{j+1}$, where $J > 0$ ($J < 0$) is the (anti-)ferromagnetic exchange coupling. Our model is endowed with a nonequilibrium generalization of the usual Glauber [5] dynamics: spins on even and odd sites experience different temperatures, T_e and T_o , which implies the violation of detailed balance [8, 9, 10]. To be specific, a configuration $s_1; s_2; \dots; s_L$ evolves into a new one by random sequential spin flips: A spin s_j flips to $-s_j$ with rate

$$w_j(s_j \rightarrow -s_j) = w_j(s_j \neq -s_j)$$

$$= \frac{1}{2} - \frac{j}{4} (s_{j-1} + s_{j+1}) \quad (1)$$

where $s_{2i} = s_e = \tanh(2J/k_B T_e)$ and $s_{2i+1} = s_o = \tanh(2J/k_B T_o)$, on even ($j = 2i$) and odd ($j = 2i+1$) sites. The time-dependent probability distribution $P(s; t)$ obeys the master equation:

$$\begin{aligned} \frac{\partial}{\partial t} P(s; t) = \\ = \sum_j w_j(s_j \rightarrow -s_j) P(s_j \rightarrow -s_j; t) - w_j(s_j) P(s; t) \end{aligned} \quad (2)$$

where the state s_j differs from s only by the spin flip of s_j . Our main goal in this work is to compute the time-dependent distribution $P(s; t)$. To do so, we compute all correlation functions $h_{j_1} \dots h_{j_n}$ $s_{j_1} \dots s_{j_n} P(s; t)$ and invoke the following relationship [5]:

$$\begin{aligned} 2^L P(s; t) = 1 + \sum_{i_1} h_{i_1} + \sum_{j_1 > j_2} h_{j_1} h_{j_2} + \\ + \sum_{j_1 > j_2 > j_3} h_{j_1} h_{j_2} h_{j_3} + \dots \end{aligned} \quad (3)$$

This expression illustrates that the knowledge of all equal-time correlation functions is equivalent to the complete knowledge of the distribution function $P(s; t)$. Recently, this implication was exploited for the steady state [9], and for the time-dependent situation yet restricted to a particularly simple initial condition [11].

The spin-flip dynamics of this Ising chain can be expressed in terms of the creation, annihilation and diffusion of domain walls, i.e., pairs of spins with opposite sign. For example, flipping s_j in the local configuration $s_{j-1} = s_j = s_{j+1} = +1$ creates two domain walls: $s_{j-1} = s_j$ and $s_j = s_{j+1}$, located on the bonds $(j-1; j)$ and $(j; j+1)$. Similarly, flipping s_j in the local configuration $s_{j-1} = s_j = s_{j+1} = -1$ has the effect of moving the domain wall on bond $(j; j+1)$ by one lattice constant to the left, corresponding to domain wall diffusion. By identifying a domain wall with a "particle", A , our spin-flip dynamics can be recast as a reaction-diffusion model, and the two examples translate into $;; \rightarrow AA$ and $AA \rightarrow ;;$, respectively. The mapping from the KISC into its dual RDS is described in detail in Table 1.

Clearly, the presence of alternating temperatures T_e , T_o in spin language translates into alternating pair annihilation and creation rates $(1 - s_e)/2$ in the RDS. We can see easily that letting T_e or T_o vanish simply prohibits pair creation entirely at even or odd sites. Remarkably, we can derive an additional, and possibly rather unexpected, benefit from this mapping: Assigning negative values for the temperatures T_e and/or T_o may appear artificial in the KISC, but is perfectly natural in the RDS: For example, $T_e < 0$ simply corresponds to a creation rate $(1 - s_e)/2 > 1/2$ which is easily implemented in a simulation. In other words, the RDS version is physically meaningful, and readily accessible, on a much wider parameter space.

$\langle n_{\uparrow} :: n_{\uparrow} i_t \rangle_{fng} = \langle n_{\uparrow} :: n_{\uparrow} \rangle \hat{P}(fng; t)$. To simplify notation, we continue to denote averages by $\langle \cdot \rangle$ for both spins and occupation variables, even though they are controlled by different statistical weights, $P(fg; t)$ and $\hat{P}(fng; t)$, respectively. In each case, it should be perfectly clear from the context which distribution is relevant. The dynamics of our model is characterized by symmetric diffusion of particles (with rate $1=2$) and pair annihilation/creation of particles with spatially alternating rates ($1 - j = 2$). In this case, the two particles are created on the (dual lattice) sites \uparrow and $\uparrow + 1$, by flipping a spin on the (original lattice) site j . Since j can be positive or negative, subject only to $1 - j = 1$ for all j , two very distinct behaviors emerge: (i) when both e and o are positive (corresponding to positive "temperatures" in the spin model), the annihilation process always occurs with a larger rate than the creation process, irrespective of whether j is even or odd; (ii) when, e.g., o is negative and e positive, the system displays a mild site-dependent frustration: at even sites j (i.e., \uparrow even and $\uparrow + 1$ odd) annihilation is more likely than creation, whereas the situation is reversed on odd sites (where \uparrow odd and $\uparrow + 1$ even). As we will see shortly, this gives rise to oscillatory dynamics.

Before diving into the details, some further remarks on physical realizations of this model are in order. When the rates are uniform ($e = o$), it is well known that such an RDS describes the dynamics of photo-excited solitons in conjugated polymers or linear chain compounds. MX chain compounds, $[Pt(en)_2][Pt(en)Cl_2]Y_4$, where Y stands for ClO_4 or BF_4 and (en) for ethylenediamine, are of particular experimental interest [12, 13]. In these compounds, photogenerated solitons are so long-lived that they can be experimentally studied. Irradiation with continuous wave (non-pulsed) blue light generates soliton-antisoliton pairs which can diffuse apart or annihilate. Their static and dynamic properties are in quantitative agreement with theoretical models [4, 18]. Since creation, annihilation, and hopping rates can be controlled by tuning the laser power, we believe that spatially alternating rates such as ours will be generated if an MX chain compound is exposed to a spatially modulated light intensity.

Returning to our model, our goal in this section is first, to derive all correlation functions from our exact solution of the KISC. We will also comment on the validity of a simple mean-field theory which is widely used for the homogeneous ($e = o$) case [18, 19]. Further, we show that particle hops in the RDS develop a peculiar directional preference in the steady state, even though there is no explicit bias in the rates, boundary or initial conditions. Finally, we illustrate how oscillatory behaviors may result from a competition of the underlying processes.

A. Density of particles in the RDS

The observable of most immediate interest is the average density of particles, $\langle \uparrow(t) \rangle$, in the RDS. Its equation of motion can be derived easily from the associated master equation, resulting in:

$$\begin{aligned} 2 \frac{d}{dt} \langle \uparrow(t) \rangle &= (2 - j - j + 1) + (j + 1 - \uparrow + 1(t)) \\ &+ j \uparrow + 1(t) - (4 - j - j + 1) \langle \uparrow(t) \rangle \\ &2 [j \langle n_{\uparrow} n_{\uparrow+1} i_t \rangle + j + 1 \langle n_{\uparrow} n_{\uparrow+1} i_t \rangle] \end{aligned} \quad (13)$$

It is worthwhile noting that this equation is the first member of an infinite hierarchy, connecting lower-order correlations to higher-order ones. In general, such hierarchies cannot be solved directly, without recourse to crude approximations. Here, the mapping to the spin chain develops its full power, allowing us to compute all correlation functions for the RDS.

The mapping from spins to particles implies that $\langle \uparrow(t) \rangle = \langle n_{\uparrow} i_t \rangle = \frac{1}{2} [1 - \langle h_{j+1} j i_t \rangle]$, so that we can just turn to Eqn. (11) to read off the answer. To express it fully in RDS language, we also need to translate the initial correlations, $\langle h_k \cdot i_0 \rangle$. For $k < \infty$ and any t (including $t = 0$), we may write $\langle h_k \cdot i_t \rangle = \langle h_{k-k+1} k+1 k+2 \dots \cdot 1 \cdot i_t \rangle = \langle h(1 - 2n_{k+1})(1 - 2n_{k+2}) \dots (1 - 2n_{\infty}) i_t \rangle$ [18, 20] whence we obtain, for arbitrary initial condition:

$$\begin{aligned} \langle \uparrow(t) \rangle &= \frac{1}{2} [1 - \langle c_{j,j+1}(t) g \rangle] \\ &= \frac{1}{2} \sum_{k < \infty} \langle h(1 - 2n_{k+1})(1 - 2n_{k+2}) \dots (1 - 2n_{\infty}) i_0 \rangle \\ &\quad [M_{k,j+1}(t) M_{j,j}(t) - M_{k,j}(t) M_{j,j+1}(t)] \end{aligned} \quad (14)$$

Since the "propagators" $M_{i,j}(t)$ decay exponentially as $t \rightarrow \infty$, the steady-state density is independent of initial conditions and spatially uniform:

$$\langle \uparrow \rangle = \langle \downarrow \rangle = \frac{1}{2} [1 - \langle \prod_{e,o} \dots \rangle] \quad (15)$$

In Appendix B, we explicitly evaluate Eqn. (14) for a generic but simple initial condition, characterized by a uniform, uncorrelated initial distribution of particles, with density $\langle \uparrow \rangle = \langle \downarrow \rangle = 0$. For simplicity, we discuss only its long-time limit here, for $\langle \uparrow \rangle = \langle \downarrow \rangle = 1=2$. We observe two distinct kinds of behaviors:

(i) When $e, o > 0$, the stationary density of particles is approached exponentially fast [except when $e = o = 1$, see (B15)], with inverse relaxation-time $2(1 - \dots)$, and a subdominant power-law prefactor $t^{-3=2}$:

$$\begin{aligned} \langle \uparrow(t) \rangle &= \frac{1}{2} [1 - \dots] \\ &= \frac{1}{2} [1 + \frac{t^{3=2} e^{-2(1 - \dots)t}}{2^{3=2} (1 - \dots)}] \end{aligned} \quad (16)$$

This long-time behavior is very similar to that found in the usual ($e = o \in [1, \infty)$) pair diffusion, annihilation, and creation process AA \rightleftharpoons ; [4, 18].

(ii) For $\rho_e < 0$, we observe a competition between the different processes. For example, when $\rho_e < 0$ and $0 < \rho_e < 1$, the annihilation (creation) reaction dominates on even (odd) sites. As a result, the stationary density is reached exponentially fast with damped oscillations:

$$\rho(t) = \frac{1}{2} \left(1 - \frac{1}{j} \right) e^{-\frac{Z}{2} t} + \frac{1}{j} e^{-\frac{Z}{2} t} \cos\left(\frac{2\pi}{j} t\right) \quad (17)$$

For initial densities other than $\rho_e = 1/2$, as shown in Appendix B, only the amplitude, or the subdominant power-law prefactor, of the expressions (16, 17) change. Since they depend on all parameters of the model, including the initial density, the dynamics is manifestly nonuniversal.

B. Two-point correlation functions of the RDS

A deeper understanding of the time-dependent spatial structures of our RDS is provided by the m -point correlation functions, $\langle n_{\uparrow} \dots n_{\uparrow} \rangle_t$ of such a model. These are related to the correlation functions of the dual spin chain, via $\langle n_{\uparrow} \dots n_{\uparrow} \rangle_t = 2^{-m} \langle \sigma_{j_1-1} \dots \sigma_{j_m-1} \rangle_t$, and are therefore exactly known. It is interesting to note that the m -point correlation function for the RDS is a superposition of all $2n$ -point correlation functions for the spin chain, with $n = 1; 2; \dots; m$. In the following, we focus on the most directly observable correlation, namely, the two-point function. To avoid unnecessary technical complications which add little insight, we specially consider a system that is initially homogeneously half-filled with A particles, without any initial correlations: $\rho_{\uparrow}(0) = 1/2$ and $\langle n_{\uparrow}(0) n_{\hat{k}}(0) \rangle = 1/4$, for $\uparrow \neq \hat{k}$. Such an initial configuration corresponds, in the KISC picture, to a system with initially neither magnetization nor correlations. In this case, as we showed in [11], the generating function takes a rather simple bilinear form which simplifies the spin-spin correlations.

With this initial condition, both the spin chain and the RDS are translationally invariant, modulo period 2. As a result, the two-point correlations $C_{\hat{k}\uparrow}^{(\hat{k});(\uparrow)}(t) = \langle n_{\uparrow} n_{\hat{k}} \rangle_t$ between two sites \uparrow and \hat{k} (with $\hat{k} > \uparrow$) depend only on the distance $\hat{k} - \uparrow$ and the parity (\hat{k}) ; $(\uparrow) \bmod 2$ of the two sites. We therefore need to distinguish four distinct correlation functions: $C_{\hat{k}\uparrow}^{ee}(t)$, $C_{\hat{k}\uparrow}^{eo}(t)$, $C_{\hat{k}\uparrow}^{oe}(t)$, and $C_{\hat{k}\uparrow}^{oo}(t)$. By virtue of our mapping to the KISC, these are determined by the 2- and 4-point spin correlations as explained in Section III C [from Eqns. (9) and (10)]. Exploiting translational invariance, the two-point correlations for the RDS, for $\hat{k} > \uparrow$, then follow as:

$$\langle n_{\uparrow} n_{\hat{k}} \rangle_t = \frac{1}{4} \left(1 - C_1^{eo}(t) \right)^2 \quad \text{if } \hat{k} - \uparrow \text{ is even}$$

$$+ \frac{1}{4} \left(1 + C_1^{eo}(t) \right)^2 \quad \text{if } \hat{k} - \uparrow \text{ is odd} \quad (18)$$

Now we are ready to discuss our results. First of all, we consider a special case, namely, nearest-neighbor correlations. If $\hat{k} = \uparrow + 1$, Eqn. (18) reduces to

$$\langle n_{\uparrow} n_{\uparrow+1} \rangle_t = \begin{cases} \frac{1}{4} (1 - C_1^{eo}(t))^2 & \text{if } \uparrow \text{ is even} \\ \frac{1}{4} (1 + C_1^{eo}(t))^2 & \text{if } \uparrow \text{ is odd} \end{cases} \quad (19)$$

Again, we should emphasize that the quantities $C_n^{eo}(t)$, $C_n^{ee}(t)$ and $C_n^{oo}(t)$ which appear in this section are the spin correlations for a particular initial condition (cf. Eqn. (9)), in contrast to the more general correlations computed in Appendix B.

It is interesting to note that, generically, $C_1^{eo}(t) \neq C_1^{oe}(t)$. Of course, after a little thought this becomes less surprising, since $\langle n_{\uparrow} n_{\uparrow+1} \rangle_t$ involves the 4-spin correlation $\langle \sigma_{j-1} \sigma_j \sigma_{j+1} \sigma_{j+2} \rangle_t = \langle \sigma_{j-1} \sigma_{j+1} \sigma_j \sigma_{j+2} \rangle_t$. So, if \uparrow is odd (even), both $j-1$ and $j+1$ are even (odd), leading to a contribution of $C_2^{ee}(t)$ vs. $C_2^{oo}(t)$, respectively.

For the general case, when \hat{k} and \uparrow are not nearest neighbors, this difference between $C_{\hat{k}\uparrow}^{eo}(t)$ and $C_{\hat{k}\uparrow}^{oe}(t)$ does not persist. If \hat{k} is even and \uparrow is odd, we find:

$$C_{\hat{k}\uparrow}^{eo}(t) = \frac{1}{4} \left(1 - C_1^{eo}(t) \right)^2 \left[C_{\hat{k}-\uparrow}^{eo}(t) \right]^2 + \frac{1}{4} C_{\hat{k}-\uparrow+1}^{ee}(t) C_{\hat{k}-\uparrow-1}^{oo}(t) \quad (20)$$

and for \hat{k} odd and \uparrow even, one obtains

$$C_{\hat{k}\uparrow}^{oe}(t) = \frac{1}{4} \left(1 - C_1^{eo}(t) \right)^2 \left[C_{\hat{k}-\uparrow}^{eo}(t) \right]^2 + \frac{1}{4} C_{\hat{k}-\uparrow+1}^{oo}(t) C_{\hat{k}-\uparrow-1}^{ee}(t) \quad (21)$$

Thanks to the simple relation between even-even and odd-odd spin correlations, Eqn. (9), the two right-hand sides are now identical.

A similar line of reasoning shows that $C_{\hat{k}\uparrow}^{ee}(t) = C_{\hat{k}\uparrow}^{oo}(t)$ for arbitrary separation $\hat{k} - \uparrow$. Invoking the two-spin correlations again, we may write

$$C_{\hat{k}\uparrow}^{ee}(t) = C_{\hat{k}\uparrow}^{oo}(t) = \frac{1}{4} \left(1 - C_1^{eo}(t) \right)^2 \left[C_{\hat{k}-\uparrow}^{eo}(t) C_{\hat{k}-\uparrow}^{ee}(t) + C_{\hat{k}-\uparrow+1}^{eo}(t) C_{\hat{k}-\uparrow-1}^{eo}(t) \right] \quad (22)$$

In the following, we discuss the consequences of these results. We first consider the steady state. Recalling our previous analysis of the spin correlations, Eqn. (7), the stationary limit of the density-density correlations becomes very simple: Provided $\hat{k} - \uparrow > 1$, we find $C_{\hat{k}\uparrow}^{(\hat{k});(\uparrow)}(1) = \frac{1}{4} (1 - C_1^{eo}(1))^2 = \frac{1}{2} (1 - C_1^{eo}(1))^2$. In other words, the two-point correlations of non-nearest-neighbor sites factorize into one-point functions, independent of parity. This kind of mean-field-like behavior

is typical of free fermion systems [4, 18]. However, the nonequilibrium nature of this model still in poses its signature. Turning to the nearest-neighbor correlations, we find that this simple factorization no longer holds { except in the special case where $\epsilon_e = \epsilon_o$. More specifically, we find

$$C_1^{oe}(l) = \frac{1}{2}(l) - \frac{1}{4} \left(\frac{2}{\epsilon_e} - \frac{2}{\epsilon_o} \right) \frac{1}{4} \quad (23)$$

and

$$C_1^{eo}(l) = \frac{1}{2}(l) + \frac{1}{4} \left(\frac{2}{\epsilon_e} - \frac{2}{\epsilon_o} \right) \frac{1}{4} \quad (24)$$

Considering, e.g., $0 < \epsilon_o < \epsilon_e$, we find that $C_1^{eo}(l)$ is enhanced over the mean-field result while $C_1^{oe}(l)$ is suppressed. This can be understood easily: Since $\epsilon_o < \epsilon_e$ implies $T_e < T_o$, energetically costly spin flips occur more frequently on odd sites j , creating a particle pair on the nearest-neighbor dual sites $(j+1; j)$. Clearly, these sites form an $(e;o)$ pair. Moreover, the rate for pair annihilation is lower on $(e;o)$ sites. Hence, particle pairs are more likely to reside on $(e;o)$ than on $(o;e)$ sites. This also implies that $(e;o)$ sites act as net particle sources, while $(o;e)$ sites function as sinks [10]. Not surprisingly, therefore, we find $C_1^{eo}(l) > C_1^{oe}(l)$. By virtue of this reasoning, it is also immediately apparent that this difference can only persist for nearest-neighbor correlations. The same argument holds for $\epsilon_o < 0 < \epsilon_e$.

A direct consequence of $C_1^{oe}(t) \neq C_1^{eo}(t)$ is the presence of a peculiar directional preference in the RDS. If we consider a particle on site j , we can ask for the average rate, $R_j(t)$, with which it will jump to the left (i.e., to site $j-1$) vs to the right, defined as $R_j(t) = \frac{1}{2} \langle \dot{n}_j(t) - \dot{n}_{j+1}(t) \rangle$. Here, the first (second) term is the average rate for a particle on site j to jump to site $j+1$ ($j-1$). In our case, one might expect this difference to vanish since neither bulk rates nor boundaries impose a directional bias. Moreover, to avoid a potential bias at $t=0$, we start from a translationally invariant initial condition with $\langle n_j(0) \rangle = 1/2$. Yet, since $R_j(t) / C_1^{oe}(t)$, it is manifestly nonzero. Explicitly, we find:

$$R_j(t) = \begin{cases} \frac{1}{8} \left(\frac{2}{\epsilon_e} - \frac{2}{\epsilon_o} \right) C_2^{ee}(t), & j \text{ even} \\ \frac{1}{8} \left(\frac{2}{\epsilon_o} - \frac{2}{\epsilon_e} \right) C_2^{ee}(t), & j \text{ odd} \end{cases} \quad (25)$$

which even persists in the steady state:

$$R_j(l) = \begin{cases} \left(\frac{2}{\epsilon_e} - \frac{2}{\epsilon_o} \right) \frac{1}{4} \frac{1}{4}, & j \text{ even} \\ \left(\frac{2}{\epsilon_o} - \frac{2}{\epsilon_e} \right) \frac{1}{4} \frac{1}{4}, & j \text{ odd} \end{cases}$$

Specifically, for $\epsilon_o < \epsilon_e$, particles on an even (odd) site jump preferentially to the right (left). Of course, this directional preference vanishes as soon as $\epsilon_e = \epsilon_o$. Moreover, even when it is nonzero, it does not generate a mass current. Counting the net flow of particles between sites j and $j+1$, the natural definition of such a current is $J_j(t) = \frac{1}{2} \langle \dot{n}_j(t) - \dot{n}_{j+1}(t) \rangle$. Clearly,

this expression reduces to a density difference which vanishes for all times provided the initial condition is homogeneous. For inhomogeneous initial condition, $J_j(t)$ exhibits nonzero transients for finite times but decays as t^{-1} .

Let us conclude this section with a few brief remarks about the validity of the mean-field approximation for this system. We already noted that it does not predict the nearest-neighbor correlations correctly, except in the special case $\epsilon_e = \epsilon_o$. We now show that it also generically misses the stationary density.

We begin by recalling Eqn. (13). Seeking a translationally invariant (modulo 2) solution with $n_{2j}(t) = \epsilon_e(t)$, $n_{2j+1}(t) = \epsilon_o(t)$ for all j , the mean-field approximation corresponds to truncating two-point functions: $\langle n_j(t) n_{j+1}(t) \rangle \approx \epsilon_e(t) \epsilon_o(t)$. Starting from a uniform initial density $\langle n_j(0) \rangle = \epsilon_e(t) = \epsilon_o(t) = \rho$, with

$$\rho_{MF}(t) = \frac{\rho [\langle n_j(0) \rangle - \rho] + \rho [\langle n_{j+1}(0) \rangle - \rho]}{\langle n_j(0) \rangle + \langle n_{j+1}(0) \rangle} e^{-\frac{\rho}{4} \frac{t}{\epsilon_e + \epsilon_o}} + \frac{\rho}{\langle n_j(0) \rangle + \langle n_{j+1}(0) \rangle} e^{-\frac{\rho}{4} \frac{t}{\epsilon_e + \epsilon_o}}; \quad (26)$$

where

$$\rho_{MF} = \frac{1}{2} \left(1 - \frac{2}{\epsilon_e + \epsilon_o} \right) \frac{\rho}{4} \frac{t}{\epsilon_e + \epsilon_o}$$

The stationary limit is clearly $\rho_{MF}(l) = \rho$ which differs from our exact result, Eqn. (15), except if $\epsilon_e = \epsilon_o$. In other words, the remarkable accuracy [18, 19] of the mean-field approximation for the stationary state of the uniform system ($\epsilon_e = \epsilon_o$) appears to be an "accident" due to the fact that when rates are uniform the steady state is a product measure. We also note that the exact relaxation time to the steady state, $\tau_{\text{exact}} = [2 \frac{\rho}{\epsilon_e \epsilon_o}]^{-1}$, does not coincide with the mean-field prediction, $\tau_{MF} = [4 \frac{\rho}{(\epsilon_e + \epsilon_o)^2}]^{-1}$. For such dynamic quantities, the exact and the approximate results differ for any choice of ϵ_e and ϵ_o . In particular, the mean-field theory always predicts an exponential decay to the steady state, completely missing the possibility of oscillatory behavior.

V. SOLVABILITY AND RELATIONSHIP WITH FREE FERMION SYSTEMS

The crucial ingredient for the solvability of the KISC is the quadratic spin dependence of its Glauber-like kinetics. Thanks to this simple form, the hierarchy of equations for the correlation functions is closed: to solve the equations for the N -spin correlation functions, one needs to know only m -point correlations with $m \leq N$.

In RDS language, the dynamics of the particles can be rewritten as a free fermion model, by defining a suitable quadratic (but non-Hermitian) "stochastic Hamiltonian". Following standard methods [3, 4, 18, 21], we

lation on odd sites while the relation is reversed on the even sites. Hence, a given initial particle density may first decrease, due to annihilation processes and then recover, as the available empty sites are (partially) filled again by the strong creation process, and so on, until the stationary density is reached.

Remarkably, even in the absence of any bias in the rates, boundary or initial conditions, particles still "know" the difference between right and left: For, e.g., $\rho_o < \rho_e$, particles on even (odd) sites jump preferentially to the right (left). Even though this directional preference does not lead to a systematic particle current, it is still somewhat surprising. However, once we recall that particles are most often created (annihilated) on pairs of neighboring sites, with the odd site on the left (right), we recognize that the directional preference is simply a response to this density gradient.

Since exact solutions, especially of a full nonequilibrium dynamics are rare, we hope that our model can serve as a testing ground for various generalizations or approximations. The features reported here (exponential decays, damped oscillations, and directional preference) should be generic for a whole class of genuine out-of-equilibrium models. Moreover, they should be experimentally observable in MX chain compounds exposed to spatially modulated laser light.

Acknowledgments: It is a pleasure to acknowledge fruitful discussions with I. Georgiev, H. Hilhorst, J.R. Hein, and U.C. Tauber. MM acknowledges the support of Swiss NSF Fellowship No. 81EL-68473. This work was also partially supported by US NSF DMR-0088451, 0308548 and 0414122.

APPENDIX A: THE DERIVATION OF THE GENERATING FUNCTION

In this appendix we provide some details for the derivation of the generating function (10), which is one of the key results of this work. We follow Aïev's work and notation [15]. Aïev established that the generating functions of a very general class of disordered Glauber-Ising spin chains, including our case, can formally be expressed in terms of two functions $M_{j;k}(t)$ and two additional quantities $W_{j;k}(t)$, which depend in a very involved fashion on $M_{k;j}(t)$. Below, we will see that these quantities are closely related to physical observables, namely, the magnetization and the two-point correlations. Here, we follow Aïev by noting that the Laplace transform of $\hat{M}_{j;k}$ is the inverse of an $L \times L$ band matrix $(s+1)\mathbb{1} - \frac{1}{2}U$. For our case, the entries of this matrix can be taken from the rates and read explicitly (L is even):

$$\begin{aligned} (s+1)\mathbb{1} - \frac{1}{2}U_{2j-1;k} &= (s+1)_{2j-1;k} - \frac{o}{2}(_{2j-1;k-1} + _{2j-1;k+1}); \quad (1 < j \leq L/2) \\ (s+1)\mathbb{1} - \frac{1}{2}U_{2j;k} &= (s+1)_{2j;k} - \frac{e}{2}(_{2j;k-1} + _{2j;k+1}) \quad (1 \leq j < L/2) \\ (s+1)\mathbb{1} - \frac{1}{2}U_{1;k} &= (s+1)_{1;k} - \frac{o}{2}(_{2;k} - _{k;L}) \\ (s+1)\mathbb{1} - \frac{1}{2}U_{L;k} &= (s+1)_{L;k} - \frac{o}{2}(_{L-1;k} - _{1;k}) \end{aligned} \quad (A1)$$

Given (A1), it is easy to evaluate the inverse of $[(s+1)\mathbb{1} - \frac{1}{2}U]_{j;k}$:

$$\hat{M}_{k;j} = \frac{1}{L} \sum_{j=1}^L \frac{X_k^j}{k} \frac{e^{i(k-j)n}}{s+1 - \cos n}; \quad (A2)$$

where $\frac{+}{n} = \frac{(2n-1)}{L}$ and $\frac{-}{n} = \frac{2n}{L}$, with $n = 1; 2; \dots; L$. In the thermodynamic limit $L \rightarrow \infty$, the two quantities $\hat{M}_{j;k}^+$ and $\hat{M}_{j;k}^-$ coincide whence we simply have

$$\hat{M}_{k;j}^+ = \hat{M}_{k;j}^- = \sum_{j=0}^L \frac{Z_j}{k} \frac{d}{2} \frac{e^{i(k-j)}}{s+1 - \cos}: \quad (A3)$$

Taking the inverse Laplace transform of (A3), we recover Eqn. (4) for the propagator.

Since the $W_{j;k}(t)$ can be expressed in terms of the $M_{j;k}(t)$, we may immediately conclude that $W_{j;k}(t) = W_{j;k}(t)$ as $L \rightarrow \infty$. According to Aïev, $W_{j;k}(t)$ is simply the two-point correlation function, $h_{j-k;j_0}$, for a particular initial condition, namely, $m_j(0) = 0$ and $h_{j-k;j_0} = 0$. For our case, these correlations were given in Eqn. (5).

For readers familiar with A l i e v's work [15], these remarks fill in the gaps between A l i e v's formal and general analysis and the special case we are interested in here. It follows that the generating function of our K I S C admits the compact and explicit representation of Eqn. (10) which encodes the complete dynamics of the system.

APPENDIX B: TRANSLATIONALLY-INVARIANT INITIAL CONDITIONS: THE SPIN-SPIN CORRELATION FUNCTIONS AND THE PARTICLE DENSITY

In Section III B, we have derived an exact expression, Eqn. (11), for the spin-spin correlation functions of our K I S C, valid for arbitrary initial conditions. Here, we impose a natural restriction, namely, translational invariance, on the initial conditions. Thanks to the symmetry, Eqn. (11) simplifies considerably, as we will show now.

As we already pointed out in [11], for translationally invariant initial conditions, we only need to consider the correlations between spins at two even sites, two odd sites, and one even, one odd site. We denote these by $c_{2n}^{ee}(t)$, $c_{2n}^{eo}(t)$, $c_{2n}^{oe}(t)$, $c_{2n}^{oo}(t)$. Of course, there is no need to study $n < 0$ cases. For the special case of zero initial magnetization and correlations, these correlations are already known [11] and are given by Eqn. (9). Here, we seek their form in a more general case, starting from a homogeneous initial condition.

Let us recall from [11] that in the translationally-invariant case the quantities $a_{2n}(t) = \frac{1}{2} [c_{2n}^{oo}(t) + c_{2n}^{ee}(t)]$; $a_{2n-1}(t) = c_{2n-1}^{eo}(t) = c_{2n-1}^{oe}(t)$, obey the following simple equation: $\frac{d}{dt}a_j = 2a_j + [a_{j+1} + a_{j-1}]$; $j > 0$ with the initial condition $a_0(t) = 0$ and $(e + o) = 2$. The equations of motion of the K I S C [11] also give the following relationships among the correlators: $c_{2n}^{oo}(t) = \frac{e}{o}c_{2n}^{ee}(t) + c_{2n}^{oo}(0) - \frac{e}{o}c_{2n}^{ee}(0)e^{-2t}$:

The explicit expressions for the correlators follow from (11), or by the methods of images directly from $a_n(t) = a_n(1) + e^{-2t} \sum_{m=0}^{\infty} [a_m(0) - a_m(1)] f_{I_n-m}(2t) - I_{n+m}(2t)g$ where $a_n(1) = \delta_n^k$ and $n \geq 0$. From the definitions of a_n , we immediately infer:

$$\begin{aligned} c_{2n}^{ee}(t) &= \frac{e}{o}c_{2n}^{oo}(t) - \frac{e}{o}c_{2n}^{oo}(0) - c_{2n}^{ee}(0)e^{-2t} \\ &= \frac{a_{2n}(1)}{o} + \frac{e^{-2t}}{o} \sum_{m=0}^{\infty} [a_{2m}(0) - a_{2m}(1)] [I_{2(n-m)}(2t) - I_{2(n+m)}(2t)] \\ &\quad + \frac{e^{-2t}}{o} \sum_{m>0} [c_{2m-1}^{eo}(0) - c_{2m-1}^{eo}(1)] [I_{2(n-m)+1}(2t) - I_{2(n+m)-1}(2t)] \\ &\quad - \frac{e^{-2t}}{2o} (e c_{2n}^{oo}(0) - o c_{2n}^{ee}(0)) : \end{aligned} \quad (B1)$$

Following the same steps for $c_{2n-1}^{eo} = c_{2n-1}^{oe}$, we obtain, for $(n > 0)$:

$$\begin{aligned} c_{2n-1}^{eo}(t) &= c_{2n-1}^{eo}(1) + e^{-2t} \sum_{m>0} [c_{2m-1}^{eo}(0) - c_{2m-1}^{eo}(1)] [I_{2(n-m)}(2t) - I_{2(n+m-1)}(2t)] \\ &\quad + \frac{e^{-2t}}{o} \sum_{m>0} [a_{2m}(0) - a_{2m}(1)] [I_{2(n-m)-1}(2t) - I_{2(n+m)-1}(2t)] : \end{aligned} \quad (B2)$$

The expressions (B1)-(B2) illustrate that the time-dependence of the spin-spin correlation function depends non-trivially on the initial condition, and we may therefore anticipate non-universal behavior. Of course, when $e = o$, the expressions (B1), (B2) coincide with those obtained by Glauber [5].

An interesting situation occurs when, say, o is negative while $0 < e < 1$, so that $\frac{e}{o} = ij$. Then, we have $I_{2n}(2ijj) = (-1)^n J_{2n}(2j)$ and $I_{2n-1}(2ijj) = (-1)^n J_{2n-1}(2j)$, where $J_n(x) = \int_0^x \cos(x \sin q - nq) dq$ is the Bessel function of first kind [16]. Further, when $\frac{e}{o} = ij$, the expressions (B1), (B2) become:

$$\begin{aligned} c_{2n}^{ee}(t) &= \frac{e}{o}c_{2n}^{oo}(t) - \frac{e}{o}c_{2n}^{oo}(0) - c_{2n}^{ee}(0)e^{-2t} \\ &= \frac{a_{2n}(1)}{j \circ j} + \frac{e^{-2t}}{j \circ j} \sum_{m=0}^{\infty} [a_{2m}(0) - a_{2m}(1)] (-1)^{n+m} [J_{2(n-m)}(2j) - J_{2(n+m)}(2j)] \end{aligned}$$

$$+ \frac{j}{j} \frac{j}{j} \frac{2t}{j} \sum_{m>0}^X [c_{2m-1}^{eo}(0) - c_{2m-1}^{eo}(1)] (1)^{n+m} J_{2(n-m)+1}(2j) + J_{2(n+m)-1}(2j) \frac{e^{2t}}{2} (e^{c_{2n}^{eo}(0)} - e^{c_{2n}^{eo}(0)}) : \quad (B 3)$$

$$c_{2n-1}^{eo}(t) = c_{2n-1}^{eo}(1) + e^{2t} \sum_{m>0}^X [c_{2m-1}^{eo}(0) - c_{2m-1}^{eo}(1)] (1)^{n+m} J_{2(n-m)}(2j) + J_{2(n+m)-1}(2j) \frac{e^{2t}}{j} \sum_{m>0}^X [a_{2m}(0) - a_{2m}(1)] (1)^{n+m} J_{2(n+m)-1}(2j) - J_{2(n-m)-1}(2j) : \quad (B 4)$$

In the long-time limit, these expressions exhibit a damped oscillatory approach to the stationary state.

We now turn to the equivalent RDS. As we have seen in Section IV A, the density of particles is related to the nearest-neighbor spin correlations, c_1^{eo} , according to $\rho(t) = \frac{1}{2}(1 - c_1^{eo}(t))$. Here, our goal is to determine the long-time behavior of this density for a homogeneous (but otherwise arbitrary) initial concentration of particles $\rho(0)$. In this respect, the expressions (B 2) and (B 4) are not very practical as they involve infinite sums of Bessel functions. At this point, for further convenience, it is useful to introduce four auxiliary functions defined as follows (with $0 \leq \rho \leq 1$):

$$F_1(\rho; t) = \sum_{m>0}^X c_{2m-1}^{eo} e^{2t} [I_{2(m-1)}(2t) - I_{2m}(2t)] = 2(1+\rho^2) \int_0^Z \frac{dq}{1+\rho^4+2\rho^2\cos 2q} \frac{\sin^2 q}{2} \quad (B 5)$$

$$F_2(\rho; t) = \sum_{m>0}^X c_{2m}^{eo} e^{2t} [I_{2m-1}(2t) - I_{2m+1}(2t)] = 2\rho^2 \int_0^Z \frac{dq}{1+\rho^4+2\rho^2\cos 2q} \frac{\sin 2q \sin q}{2} \quad (B 6)$$

$$G_1(\rho; t) = \sum_{m>0}^X c_{2m-1}^{eo} (1)^m e^{2t} [J_{2(1-m)}(2j) + J_{2m}(2j)] = 2(1+\rho^2) \int_0^Z \frac{dq}{1+\rho^4+2\rho^2\cos 2q} e^{2t} \cos(2j) \sin q \frac{\cos^2 q}{2} \quad (B 7)$$

$$G_2(\rho; t) = \sum_{m>0}^X c_{2m}^{eo} (1)^m e^{2t} [J_{2m+1}(2j) - J_{1-2m}(2j)] = 2\rho^2 \int_0^Z \frac{dq}{1+\rho^4+2\rho^2\cos 2q} e^{2t} \sin(2j) \sin q \frac{\sin 2q \cos q}{2} \quad (B 8)$$

To establish these expressions, we have invoked the integral representation of the Bessel functions [16] and the properties of geometric series. With these functions and the help of Eqn. (14), the density of particles in the RDS model can now be recast in compact form. Two cases emerge naturally:

When $\rho > 0$:

$$\rho(t) = \frac{1}{2} [F_1(\rho; t) + F_2(\rho; t)] = \frac{1}{2} [F_1(1-\rho^2(0); t) + F_2(1-\rho^2(0); t)] \quad (B 9)$$

When $\rho < 0$:

$$\rho(t) = \frac{1}{2j} [G_2(1-\rho^2(0); t) - G_1(\rho; t)] = \frac{1}{2} [G_2(\rho; t) + G_1(1-\rho^2(0); t)] \quad (B 10)$$

The case $\rho = 0$ is special and gives rise to a purely exponential time-dependence:

$$\rho(t) = \frac{1}{2} + \frac{2}{4} \rho(0) e^{-2t} \quad (B 11)$$

We now proceed with the analysis of the long-time behavior of these expressions. A gain, we first consider the case where $\rho > 0$ and then $\rho < 0$.

When $\rho > 0$, the main contribution to the long-time behavior arises from the small q contribution in the expression of the functions F_1 and F_2 . Therefore, one may expand the integrand of F_1 and F_2 in Eqn. (B 9). It is also essential to pay due attention to the initial condition:

1. When $0 < \rho_0 < 1$ and $0 < \rho(0) < 1$ (also $\rho(0) \notin (1)$), we obtain

$$(t) \quad (1)' \frac{1}{4} \frac{!}{(1!^2)} \frac{1 + - f1 \quad 2 \quad (0)g + 2 \quad (0)^2}{(0)(1 \quad (0))} \frac{e^{2(1)t}}{t^{\frac{!}{t}}} : \quad (\text{B } 12)$$

2. When $0 < \rho_0 < 1$ and $\rho(0) = 0$, we find

$$(t) \quad (1)' \quad 1 + - \frac{!}{(1!^2)} \frac{1}{t} \frac{e^{2(1)t}}{4^{\frac{!}{t}}} : \quad (\text{B } 13)$$

3. When $0 < \rho_0 < 1$ with $\rho(0) = 1$, we have

$$(t) \quad (1)' \quad 1 - + \frac{!}{(1!^2)} \frac{1}{t} \frac{e^{2(1)t}}{4^{\frac{!}{t}}} : \quad (\text{B } 14)$$

These results show that for $\rho_0 > 0$, the density generically approaches its stationary value as $/ t^{3=2} e^{2(1)t}$, with some nontrivial amplitude. Only if the lattice is initially completely empty/occupied by particles, the long-time behavior is modified to $/ t^{1=2} e^{2(1)t}$ (provided $- \notin 1$).

4. The case where $\rho_0 = \rho = 1$ is critical and we can check from (B 9) that one recovers the previously known results [4, 18]:

$$(t) \quad (1)' \frac{!}{2^{\frac{!}{t}}} : \quad (\text{B } 15)$$

In this case, it is well known [4, 18] that the density of particles approaches the steady state algebraically slowly ($/ t^{1=2}$). For $\rho = 1$ (only pair annihilation), the stationary value is $(1) = 0$; in contrast, we find $(1) = 1$ for $\rho = 1$ (only pair creation). We emphasize that for such a critical dynamics neither the dynamical exponent nor the amplitude of (B 15) depend on the initial condition.

When $\rho_0 < 0$, it is difficult to directly analyze the long-time behavior of the oscillating function G_1 and G_2 ; instead, we seek upper and lower bounds for $\rho \notin 1$. We observe that the denominator of the integrand in the expressions for G_1 and G_2 can be bounded as follows: $(1 - \rho^2)^2 = (1 + \rho^4 + 2\rho^2 \cos 2q) = (1 + \rho^2)^2$. Therefore, we obtain for the auxiliary functions G_1 and G_2 :

$$\frac{e^{2t} J_1(2j)}{1 + \rho^2 \frac{e^{2t} J_1(2j)}{2j}} \quad G_1(j; t) \quad \frac{(1 + \rho^2) e^{2t} J_1(2j)}{(1 - \rho^2)^2 \frac{e^{2t} J_1(2j)}{2j}} \quad (\text{B } 16)$$

$$2 \frac{e^{2t} J_2(2j)}{1 + \rho^2 \frac{e^{2t} J_2(2j)}{2j}} \quad G_2(j; t) \quad 2 \frac{e^{2t} J_2(2j)}{1 - \rho^2 \frac{e^{2t} J_2(2j)}{2j}} : \quad (\text{B } 17)$$

If $\rho = 1$, one has the exact expressions: $G_1(1; t) = G_1(-1; t) = e^{2t} J_0(2j)$; $G_2(1; t) = G_2(-1; t) = \frac{e^{2t}}{2j} J_1(2j)$. At long times and for finite n , $e^{2t} J_n(2j) \sim \frac{e^{2t}}{2j} \cos 2j \sim \frac{1}{4} (2n + 1)$ and therefore the upper and lower bounds in Eqns. (B 16) and (B 17) display the same time-dependence. With the help of Eqn. (B 10), we thus deduce:

1. When $0 < \rho(0) < 1$ (and obviously $\rho(0) \notin (1)$),

$$(t) \quad (1)' \quad t^{3=2} e^{2t} \left[A \cos 2j + \frac{1}{4} + B \cos 2j - \frac{1}{4} \right] ; \quad (\text{B } 18)$$

where A and B are some amplitudes depending nontrivially on all the parameters of the system and on the initial density.

2. When $\rho(0) = 1$, we obtain an explicit expression for the long-time behavior of the density:

$$(t) \quad (1)' \quad \frac{e^{2t}}{2} J_0(2j) \sim \frac{e^{2t}}{2j} \cos 2j \sim \frac{1}{4} : \quad (\text{B } 19)$$

3. When $\rho(0) = 0$, we also have an explicit expression for the long-time behavior of the density:

$$(t) \quad (1)' \quad \frac{e^{2t}}{2} J_0(2j) \sim \frac{e^{2t}}{2j} \cos 2j + \frac{3}{4} : \quad (\text{B } 20)$$

These results show that, for $\rho_0 < 0$, the density displays oscillations which are damped by a factor t^{-2} , where $\beta = 3/2$ for generic initial densities $\rho(0)$, with two exceptions: we have $\beta = 1/2$ if the system is initially completely empty or occupied.

Of course, following the same approach one would be able to compute every n point correlation functions, for both, the KIS and RDS. While perfectly straightforward, these computations become rather tedious.

-
- [1] B. Schmittmann and R.K.P. Zia, Statistical mechanics of driven diffusive systems. In: Phase Transitions and Critical Phenomena, Vol. 17, ed. by C. Domb and J.L. Lebowitz (Academic Press, New York 1995); D. Mukamel, Phase transitions in non-equilibrium systems, In: Soft and Fragile Matter: Nonequilibrium Dynamics, Metastability and Flow, eds. M.E. Cates and M.R. Evans (IOP Publishing, Bristol, 2000).
- [2] Nonequilibrium Statistical Mechanics in One Dimension, edited by V. Privman (Cambridge University Press, Cambridge, 1997).
- [3] G.M. Schutz, Integrable stochastic many-body systems. In: Phase Transitions and Critical Phenomena, Vol. 19, ed. by C. Domb and J. Lebowitz (Academic Press, London, 2000).
- [4] A.A. Lushnikov, Sov. Phys. JETP 64, 811 (1986); M.D. Grynberg, T.J. Newman and R.B. Stinchcombe, Phys. Rev. E 50, 957, (1994); M.D. Grynberg and R.B. Stinchcombe, Phys. Rev. Lett. 74, 1242 (1995); M.D. Grynberg and R.B. Stinchcombe, Phys. Rev. E 52, 6013 (1995); M.D. Grynberg and R.B. Stinchcombe, Phys. Rev. Lett. 76, 851 (1996); P.-A. Bares and M.M. Obilia, Phys. Rev. E 59, 1996 (1999); P.-A. Bares and M.M. Obilia, Phys. Rev. Lett. 83, 5214 (1999); S.-C. Park, J.-M. Park and D. Kim, Phys. Rev. Lett. 85, 892 (2000); P.-A. Bares and M.M. Obilia, Phys. Rev. Lett. 85, 893 (2000).
- [5] R.J. Glauber, J. Math. Phys. 44, 294 (1963).
- [6] Z. Racz, Phys. Rev. Lett. 55, 1707 (1985).
- [7] F. Family and J.G. Amar, J. Stat. Phys. 55, 1235 (1991).
- [8] Z. Racz and R.K.P. Zia, Phys. Rev. E 49, 139 (1994).
- [9] B. Schmittmann and F. Schumacher, Phys. Rev. E 66, 046130 (2002).
- [10] F. Schumacher and B. Schmittmann, J. Phys. A 35, 2569 (2002).
- [11] M.M. Obilia, R.K.P. Zia and B. Schmittmann, J. Phys. A 37, L407 (2004).
- [12] N. Kuroda, M. Nishida, Y. Tabata, Y. Wakabayashi and K. Sasaki, Phys. Rev. B 61, 11217 (2000).
- [13] N. Kuroda, Y. Wakabayashi, M. Nishida, N. Wakabayashi, M. Yamashita, and N. Matsushita, Phys. Rev. Lett. 79, 2510 (1997); N. Kuroda, Y. Tabata, M. Nishida, and M. Yamashita Phys. Rev. B 59, 12973 (1999); Y. Tabata and N. Kuroda, Synth. Met. 101, 329 (1999); Y. Tabata and N. Kuroda, Phys. Rev. B 61, 3085 (2000); H. Tanaka, K. Manumoto, S. Kuroda and M. Yamashita, Synth. Met. 101, 135 (2003).
- [14] R. Kroon, H. Fleurent and R. Sprik, Phys. Rev. E 47, 2462 (1993); R. Kopeľman, C. S. Li, and R. Sprik, J. Lum. 45, 40 (1990).
- [15] M.A. Aliev, Physica A 277, 261 (2000).
- [16] M. Abramowitz and I. Stegun, Handbook of Mathematical Functions (Dover, NY, 1965).
- [17] C. Itzykson and J.-M. Drouot, Statistical field theory. Vol. 1 (Cambridge University Press, Cambridge 1989).
- [18] M.M. Obilia, Phys. Rev. E 65, 046127 (2002); M.M. Obilia and P.-A. Bares, Phys. Rev. E 63, 056112 (2001); M.M. Obilia, PhD Thesis EPFL N. 2552 (April 2002).
- [19] J.R.G. Mendoca and M.J. de Oliveira, J. Stat. Phys. 78, 1429 (1998).
- [20] J.E. Santos, J. Phys. A 30, 3249 (1997).
- [21] B.U. Felderhof, Rep. Math. Phys. 1, 215 (1971), and 2, 151 (1971).
- [22] N.N. Bogoliubov, Studies in Statistical Mechanics, edited by J. de Boer and E. Uhlenbeck (North-Holland Publishing, Amsterdam 1962), Vol. 1.