

Are Barabási-Albert networks self-averaging?

Dietrich Stauffer and Amnon Aharony*

Institute for Theoretical Physics, Cologne University D-50923 Köln, Euroland

* Visiting from School of Physics and Astronomy, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv, Tel Aviv 69978, Israel and from Department of Physics, Ben Gurion University, Beer Sheva 84105, Israel.

e-mail: aharony@post.tau.ac.il, stauffer@thp.uni-koeln.de

Abstract: Yes and no. The size of the largest neighborhood in a Barabási-Albert scale-free network has strong fluctuations of the order of the average value. The number of sites having exactly ten neighbors increases linearly in the network size while its relative fluctuations decrease towards zero if the number of sites in the network increases from 1000 to ten million.




Figure 1: 100 examples for the size k_1 of the largest neighborhood, with $10^3, 10^4, 10^5, 10^6, 10^7$ nodes (bottom to top) added to the initial $m = 3$ nodes.




Figure 2: Same simulations as for Fig.1, now showing the size k_2 of the second-largest neighborhood.

Self-averaging quantities have relative fluctuations which go to zero for system size going to infinity, like the magnetization away from the Curie point. Not self-averaging systems are found e.g. at second-order phase transitions, like the magnetization at the Curie point. Here we want to find out whether the scale-free networks of Barabási and Albert [1] are self-averaging. The number of sites having a fixed number of neighbors is found to be strongly self-averaging, confirming earlier results [2]. However, the fluctuations in the two largest neighborhoods are seen to be proportional to these neighborhoods, implying large variations among different realizations. Although the distribution of the largest neighborhood was discussed in [2], the consequences for self-averaging were not emphasized there.

To build these networks we start with m sites all connected to each other; we took $m = 3$. Then N more sites are added one-by-one and each new site selects randomly exactly m old sites as neighbors, such that the probability for an old site to be selected is proportional to the number of neighbors it has already. Our network is undirected, i.e. if node A selected node B as neighbor, then A is neighbor of B and B is neighbor of A. The network is called scale-free since the number of sites (nodes) having exactly k neighbors decays as $1/k^3$. The size k of the largest neighborhood is called k_1 , that




Figure 3: Same simulations as for Figs.1 and 2, now showing the number of nodes having exactly 10 neighbors.

of the second-largest neighborhood is k_2 , and the number of nodes having exactly k neighbors is $n(k)$.

We made 100 samples for $N = 10^3, 10^4, 10^5, 10^6$ and show in our semilogarithmic figures without further analysis quite clear results: The relative fluctuations of k_1 in Fig.1 and of k_2 in Fig.2 remain strong even if we increase the network size by four orders of magnitude. For $n(k = 10)$, on the other hand, we see in Fig.3 less and less relative fluctuations, as predicted in [3]. Fig.4 confirms the visual impression from Figs.1-3 by showing the relative fluctuations of the three discussed quantities; the ones for $n(10)$ and $n(100)$ decay as $1/\sqrt{N}$. The non-universal variation of the relative fluctuations for the largest neighbourhood with varying m (= number of neighbours selected by newly-added site) is shown in Fig.5.

We are not surprised about these results since they are similar to cluster statistics in percolation (and presumably also in Ising) models. The size of the largest cluster at the critical point fluctuates strongly even in the thermodynamic limit [4], whereas the numbers of clusters of a fixed finite size get the more accurate the larger the system is [5].

The German-Israeli Foundation supported this collaboration.




Figure 4: Same simulations as for Figs.1-3, now showing the relative fluctuations of k_1 , k_2 , $n(10)$ and $n(100)$.

References

- [1] R. Albert and A.L. Barabási, Rev. Mod. Phys. 74, 47 (2002).
- [2] P.L. Kaprivsky and S. Redner, J. Phys. A 35, 9517 (2002) and Phys. Rev. Lett. 89, 258703 (2002); A.A. Moreira, J.S. Andrade, Jr., and L.A.N. Amaral, Phys. Rev. Lett. 89, 268703 (2002).
- [3] S.N. Dorogovtsev and A.N. Samukhin, Phys. Rev. E 67, 037103 (2003); see also S. Itzkovitz, R. Milo, N. Kashtan, G. Ziv, and U. Alon, Phys. Rev. E 68, 02617 (2003).
- [4] D. Stauffer and A. Aharony, *Introduction to Percolation Theory*, Taylor and Francis, London 1994.
- [5] A. Coniglio, H.E. Stanley and D. Stauffer, J. Phys. A 12, L 323 (1979); D. Tiggemann, Int. J. Mod. Phys. C 13, 777 (2002).




Figure 5: Log-log plot of relative fluctuations of k_1 versus m ; the straight line has slope -0.6 .