Sem jon Stepanow

Martin-Luther-Universitat Halle, Fachbereich Physik, D-06099 Halle, Germany

PACS num bers: 05.40.-a, 36.20.-r, 61.41.+ e

E-m ail: stepanow@physik.uni-halle.de

A bstract. Using the exact computation of a large number of moments of the distribution function of the end-to-end distance G(r;N) of the worm-like chain, we have established the analytical form of the coe cients in Taylor expansions of the moments for short chain lengths N. The knowledge of these coe cients enabled us to resum mate the moment expansion of G(r;N) by taking into account consecutively the deviations of the moments from their sti rod limit. W ithin this procedure we have derived the short-chain expansion for G(r;N), the scattering function, and the extension-force relation, which take into account the deviations of the moments from their sti rod limit is of the moments from the store of the moments from the extension for G(r;N).

## 1. Introduction

Polym ers with contour length L much larger than the persistence length  $l_p$ , which is the correlation length for the tangent-tangent correlation function along the polym er and is quantitative measure of the polym er sti ness, are exible and are described by using the tools of quantum mechanics and quantum eld theory []-[6]. If the chain length decreases, the chain sti ness becomes an important factor. M any polymer molecules have internal sti ness and cannot be modelled by the model of exible polymers developed by Edwards [1].

The standard coarse-graining model of a worm-like polymer was proposed by K ratky and Porod [7]. The essential ingredients of this model is the penalty for bending energy, and the local inextensibility. The latter make the treatment of the model much more di cult. There is a substantial amount of studies of the K ratky-Porod model in the last half century [8]-[18] (and the references therein). In recent years there is an increasing interest in theoretical description of semi i exible polymers [9]-[29] (and the references therein). A reason for this interest is due to potential applications of semi i exible polymers in biology and in researches on sem i-crystalline polymers.

In this article we present results of the study of the behaviour of the worm -like chain for short lengths. The consideration is based on our recent work [24], [29], where the Fourier-Laplace transform G (k;p) of the end-to-end distribution function G (r;N ) (p is the Laplace conjugate to N) was represented as the matrix element of the in nite order matrix  $P'(k;p) = (I + ikDM)^{1}D$  with matrices D and M related to the spectrum of the quantum rigid rotator. A truncation of P (k;p) by matrix of order n gives the end-to-end distribution function G (k;p) as a rational function being an in nite series in powers of  $k^2$ , i.e. it contains all m om ents of the end-to-end distribution function, and describes the rst 2n 2 m om ents exactly. In context of eigenstates of the quantum rigid rotator, the truncation at order n takes into account the eigenstates with quantum number of the angular momentum up to the value l = n 1. The moment expansion of G (k;p) can be represented as double series in powers of  $k^2$  and 1=p. A lternatively, one can expand G (k; N) in double series in powers of  $k^2$  and N. A nalyzing the series of known m om ents (we have analytically calculated the st 50 m om ents), we have established that the coe cients in subseries in powers of k<sup>2</sup> have a simple analytical structure, which enables one to perform a partial resummation of the moment expansion of G (k;p) or G (k;N) (see below). This resummation procedure results in an expansion of G (r;N) around the sti rod limit, where the end-to-end distribution function is given by the expression,  $G_r(r; N) = 1 = (4 N^2)$  (N r), so that because  $G_r(r; N)$  is a distribution, so derived short-chain expansion is not a Taylor expansion but rather an expansion in the space of distributions. The know ledge of G (k; N) enables one to compute directly the scattering function S (k;N) and the extension-force relation R (f). In this paper we present the results of the calculation of terms of the short-chain expansions of G (r; N), S (k; N), and R (f) by taking into account the deviations of the moments from their sti rod behaviour to the seven order in chain length N . The procedure can be extended to

higher orders. The short-chain expansions besides the interest in their own can be used for comparisons with approximative treatments, and also in studies of the behaviour of short sem i exible polymers.

The present paper is organized as follows. Section 2 introduces to the description of the worm-like chain using the form alism of quantum rigid rotator. Section 3 explains the idea of the derivation of the short-chain expansion, and presents the short-chain expansions for the distribution function of the end-to-end distance, scattering function, and extension-force relation.

# 2. The form alism

The Fourier transform of the distribution function of the end-to-end polymer distance of the continuous K ratky-P orod m odel [7] G (k;L) =  $d^{3}R \exp(ik(R R_{0}))G(R R_{0};L)$  is expressed by the path integral as follows

$$G(k;L) = Dt(s) Y (t(s)^{2} 1) \exp(ik \int_{0}^{Z_{L}} dst(s) \frac{1}{2} \int_{0}^{Z_{L}} ds(\frac{dt(s)}{ds})^{2}; \qquad (1)$$

where  $l_p$  is the persistence length, and t (s) = dr (s)=ds is the tangent vector at the point s along the contour of the polymer. The product over s in Eq.(1) takes into account that the polymer chain is locally inextensible. In the following the arc length of the polymer L will be measured in units of  $l_p$  and will be denoted by N . We now will consider the G reen's function P (;';N; <sub>0</sub>;' <sub>0</sub>;0) associated with Eq.(1). The dimensional equation for P is

$$\frac{@}{@N}P(;';N;_{0};'_{0};0) = \frac{1}{2}r^{2};'P + U()P = (N)() (_{0}); (2)$$

where U (kt ) = ikt is the potential energy of the rigid rotator in an external eld ik, where k is measured in units of  $l_p^{1}$ . The end-to-end distribution function is obtained from P (;N; <sub>0</sub>;0) as follows

$$G(k;N) = \frac{1}{4} \overset{2}{d} \overset{2}{d} 0 P(;N;_{0};0):$$
(3)

The di erential equation  $\ell$ ) can be rew ritten as an integral equation as follows  $Z_{\rm ev} = Z_{\rm ev}$ 

$$P(;N;_{0};0) = P_{0}(;N;_{0};0) \qquad ^{N} ds \quad d^{0}P_{0}(;N;_{0};s)U(kt_{0})P(_{0}^{0};s;_{0};0);(4)$$

where the bare G reen's function  $P_0$  (;'; N;  $_0$ ; ') reads

$$P_{0}(;';N;_{0};'_{0};0) = \sum_{lm}^{X} \exp(\frac{l(l+1)N}{2})Y_{lm}(;')Y_{lm}(_{0};'_{0});$$
(5)

with  $Y_{lm}$  (;') being the spherical harmonics, and l and m are the quantum numbers of the angular momentum. Due to the convolution character of the expression (4) with respect to the integration over the contour length (P<sub>0</sub>(;N; <sup>0</sup>;s) depends on the di erence N s), the Laplace transform of P(;N; <sub>0</sub>;0) in Eq.(4) with respect to N

perm its to get rid of integrations over the contour length. Thus, in the following we will consider the Laplace transform of G (k; N) with respect to N.

It was shown in [24], [29] that the solution of Eq.(4) results in the following expression of the Fourier-Laplace transform of the end-to-end distribution function as the matrix element of an in nite order square matrix

$$G (k;p) = 0 j P^{s} (k;p) j 0$$
(6)

with

$$\mathcal{P}^{s}(\mathbf{k};\mathbf{p}) = (\mathbf{I} + \mathbf{i}\mathbf{k}\mathbf{D}\mathbf{M}^{s})^{\perp}\mathbf{D};$$
 (7)

where the square matrices M  $^{\rm s}$  and D  $\,$  are de  $\,$  ned by

$$M_{1;1^{0}}^{s} = w_{1,1;1^{0}+1} + w_{1+1,1^{1}+1;1^{0}};$$
(8)

with 
$$w_1 = \frac{p}{l^2 = (4l^2 - 1)}$$
, and  
 $D_{1;l^0} = \frac{1}{\frac{1}{2}l(l+1) + p}$  1;l^0; (9)

respectively. The superscript s speci es that the quantities  $P^{s}$  and  $M^{s}$  are square m atrices. The quantity h0 jP<sup>s</sup> (k;p) j0i denotes the (1;1) m atrix element of the in nite order square m atrix  $P^{s}$ . Since the (1;1) m atrix element corresponds to the expectation value of the quantum rigid rotator in the ground state with the quantum number l=0, we prefer to use the above notation. Sum m ations over the m agnetic quantum number in the intermediate states in the expression of G (k;p) obtained by solving Eq.(4) can be eliminated [29], so that sum m ations in the intermediate states occur only over the eigenvalues of the angularm on enturn l=0;1;... This is the reason why the calculation of G (k;p) reduces to the computation of the matrix element of an in nite order square m atrix.

The quantity  $\mathbb{P}^{s}(k;p)$  plays the key role in the theory similar to the bare propagator in common quantum  $\mathrm{eld}_{D}$  theories. The end-to-end distribution function G (k;p) is simply the matrix element  $0 \ \mathbb{P}^{s}(k;p) \ \mathbb{D}$ , the scattering function of the polymer is the inverse Laplace transform of G (k;p)=p<sup>2</sup> multiplied by 2=N, the partition function of the stretched polymer is Z (f;N) = G (k = if=k\_{\mathrm{B}} T;N) etc [29].

#### 3. The behaviour of short worm -like polymer

#### 3.1. The idea of the short-chain expansion

To derive the short-chain expansion for the K ratky-P orod chain we insert the expansion of D given by Eq.(9) according to

$$\frac{1}{\frac{1}{2}l(l+1)+p} = \frac{1}{p} \sum_{m=0}^{X^{L}} \frac{(1)^{m} (l(l+1))^{m}}{2^{m}} \frac{1}{p^{m}}$$

into the moment expansion of G (k;p)

$$G(k;p) = \frac{1}{p} \sum_{m=0}^{M^{d}} (1)^{m} \quad 0 \neq (D M^{s})^{2m} \neq 0 \quad (k^{2})^{m};$$
(10)

and obtain as a result G (k;p) as double series in powers of  $(k=p)^2$  and 1=p as follows

$$G(k;p) = \sum_{s=1}^{X^{i}} \frac{1}{p^{s}} \sum_{n=0}^{X^{i}} C_{n}^{s} \frac{k^{2}}{p^{2}}$$
(11)

The analysis of 50 exactly computed m om ents of G (r;p), which are proportional to the coe cients h0 j (D M <sup>s</sup>)<sup>2m</sup> j0i in Eq.(10), shows that the coe cients  $C_n^s$  are polynom ials in powers of n of the order 2s 3, i.e. they have the form  $C_n^s = a_1n + a_2n^2 + \dots + a_{2s 3}n^{2s 3}$  (s 2). We determ ine the coe cients  $a_i$  at given s using 2s 3 known terms of the m om ent expansion. Further we test the correctness of so obtained  $C_n^s$  from the comparison with the remaining known terms in the m om ent expansion, which were not used to determ ine  $a_i$ . The exact know ledge of 50 m om ents enables one to determ ine  $C_n^s$  for s 11. The coe cients  $C_m^s$  we have obtained in this way are

$$C_{n}^{1} = \frac{1}{2n + 1};$$

$$C_{n}^{2} = \frac{n}{3};$$

$$C_{n}^{3} = \frac{1}{90}n(n + 1)(14n + 1);$$

$$C_{n}^{4} = \frac{1}{1890}n(n + 1)(2n + 3)(62n^{2} + 3n - 2);$$

$$C_{n}^{5} = \frac{1}{37800}n(n + 1)(n + 2)(2n + 3)(508n^{3} - 84n^{2} - 19n + 15);$$

$$C_{n}^{6} = \frac{1}{3742200}n(n + 1)(n + 2)(2n + 3)(2n + 5)(10220n^{4} - 6236n^{3} + 1597n^{2} + 737n - 372);$$
:::

We have also computed the coe cients  $C_n^7$  and  $C_n^8$ , but do not write them here to save space. The determination of higher coe cients demands the knowledge of more moments of the end-to-end distribution function and can be obtained in a similar way. A fler the determination of the coe cients  $C_n^s$  we have checked the equivalence of (11) with (10) to the corresponding order. Unfortunately, we did not succeed in deriving the expression for the coe cients  $C_n^s$  for arbitrary s.

W ith known coe cients  $C_n^s$  one can sum the series over n in (11). Restricting ourselves to the leading order  $s_{max} = 1$  ( $s_{max}$  is the number of terms in the sum over s in (11)) we obtain G (k;N)

:

$$G_{r}(k;N) = \frac{\sin (kN)}{kN};$$

and the distribution function of the end-to-end distance

$$G_{r}(r; N) = \frac{1}{4 N^{2}} (N r)$$

in the sti rod lim it. Taking into account the next term in the sum over s in Eq.(1) gives the correction to the sti rod lim it of the end-to-end distribution function. Thus, the resum m ation of the moment expansion of G (k;p) according to (11) yields the short-chain expansion for K ratky-P orod chain.

The above derivation does not allow to make claims on the convergence of the short-chain expansion. The comparison of results of computations of quantities under consideration (for example the scattering function or deformation-force relation) for dimensionary gives a criterion determining the quality of the short-chain expansion.

Notice that alternatively one can st carry out the inverse Laplace transform of (10), and then expand the moments in Taylor series in powers of N. As a result one will arrive at double series similar to (11) with 1=p replaced by N. This makes more clear the meaning of the resummation procedure. To leading order ( $s_{max} = 1$ ) one replaces the moments by its sti rod behaviour,  $hR^{2n}i = N^{2n}$ . In the next to leading order ( $s_{max} = 2$ ) one takes into account the next-order corrections to the sti rod behaviour of all moments, and so on.

Due to the fact that the scattering function of the sem i exible polymer S (k;N) is the inverse Laplace transform of G (k;p)= $p^2$  multiplied with the factor 2=N, the shortchain expansion of G (k;p) enables one to get the short-chain expansion of the scattering function. In fact, Herm and and Ullman [8] derived the sti rod limit of the scattering function using the sti rod limit of the moments.

In following subsections we will consider separately the short-chain expansion of the end-to-end distribution function, the scattering function, and the extension-force relation, which also can be obtained from G(k;N).

#### 3.2. D istribution function of the end-to-end distance

The series over n in (11) for known coe cients  $C_n^s$  can be easily expressed through the derivatives of the geom etric series. Carrying out the inverse Laplace transform over p gives the short-chain expansion of G (k;N). We have computed G (k;N) by taking into account eight terms in the sum over s in (11) i.e.  $s_{max} = 8$ . To save place we present below the result for  $s_{max} = 5$ 

$$G_{5}(k;N) = G_{r}(k;N) + \frac{1}{6} \frac{\sin(kN)}{k} \frac{1}{6}N \cos(kN)$$

$$+ \frac{1}{60} \frac{\sin(kN)N}{k} \frac{7}{360} k \sin(kN)N^{3} \frac{1}{60}N^{2}\cos(kN)$$

$$+ \frac{1}{630} \frac{\sin(kN)N^{2}}{k} + \frac{1}{5040} k \sin(kN)N^{4} \frac{1}{630}\cos(kN)N^{3} + \frac{31}{15120}\cos(kN)N^{5}k^{2}$$

$$+ \frac{1}{5040} \frac{\sin(kN)N^{3}}{k} \frac{1}{5600} k \sin(kN)N^{5} + \frac{127}{604800} k^{3}\sin(kN)N^{7}$$

$$\frac{1}{5040}\cos(kN)N^{4} \frac{53}{151200}\cos(kN)N^{6}k^{2}; \qquad (12)$$

The subscript at G and at quantities below is  $s_{max}$ . Since  $G_s(k;N)$  tends to one for  $k \mid 0$ , the end-to-end distribution function is normalized. It is easily to see that the

term s in (12) can be represented as derivatives of sin (kN) = k w ith respect to N i.e. of N G<sub>r</sub>(k;N). The short-chain expansion of the distribution function of the end-to-end distance is then immediately obtained from (12) as

$$G_{5}(\mathbf{r};\mathbf{N}) = \frac{1}{4r^{2}} (\mathbf{N} + \mathbf{r}) + \frac{1}{12r} (\mathbf{N} + \mathbf{r}) + \frac{1}{24} (\mathbf{N} + \mathbf{r}) + \frac{1}{1008} r^{2} (\mathbf{N} + \mathbf{r}) + \frac{1}{1008} r$$

where <sup>(k)</sup> (x) denotes the kth derivative of the D irac's delta-function. Note that G (r;N) is a distribution function with respect to r, while the contour length N is a parameter. This should be taken into account in using Eq.(13) to compute di erent mean values, for example the moments of the end-to-end distribution function of a short chain. Sim ilar to the corresponding expansion of the function  $_{a}(x) = 1 = (2 \ a^{2})^{1=2} \exp(x^{2}=2a^{2})$  in powers of a, the short-chain expansion of G (r;N) cannot be interpreted as a Taylor series but is rather a generalized Taylor expansion in space of distributions. The expansions (12-13) can be used to test approximative expressions of the end-to-end distribution function. We did not succeed so far to convert the short-chain expansion of the distribution function function function to a close expression.

## 3.3. The scattering function

The scattering function of a semi exible polymer is de ned by

$$S(q;N) = \frac{2}{N} \int_{0}^{Z_{N}} ds_{2} ds_{1} hexp(iq(r(s_{2}) r(s_{1})))i;$$
(14)

Expressing  $r(s_2)$   $r(s_1)$  in (14) through the tangent vectors,  $r(s_2)$   $r(s_1) = \frac{\kappa_{s_2}}{s_1} dst(s)$ , and representing the average in (14) using the form alism of the quantum rigid rotator yields that the scattering function of the sem i exible polymer S (q;N) is the inverse Laplace transform of G (q;p)=p<sup>2</sup> multiplied with the factor 2=N [24],[29]. Thus, the short-chain expansion of G (k;p) enables one to derive in a straightforward way the short-chain expansion of the scattering function. Taking into account the rst four terms in the sum over s in (11) results in the following expression for the scattering function

$$S_{4}(x; N) = N = S_{r}(x) = N + \frac{2N}{3x^{2}} + \frac{N \sin x}{x^{3}} + \frac{1}{3} \frac{N \cos x}{x^{2}} + \frac{7}{180} \frac{N^{2} \sin x}{x} + \frac{6N^{2}}{5x^{4}} + \frac{13}{5x^{4}} \frac{N^{2} \sin x}{x^{3}} + \frac{4}{15} \frac{N^{2} \cos x}{x^{2}} + \frac{6N^{2} \cos x}{5x^{4}} + \frac{32}{63} \frac{N^{3}}{x^{4}} + \frac{307}{7560} \frac{N^{3} \sin x}{x} + \frac{125}{126} \frac{N^{3} \sin x}{x^{3}} + 3 \frac{N^{3} \sin x}{x^{5}} + \frac{31}{126} \frac{N^{3} \cos x}{x^{2}} + \frac{157}{63} \frac{N^{3} \cos x}{x^{4}} - \frac{31}{7560} N^{3} \cos x;$$
  
where x = cN, and

$$S_r(x) = N = 2 \frac{\cos(x)}{x^2} - \frac{2}{x^2} + 2 \frac{\sin(x)}{x}$$

is the scattering function of a sti rod. The plot of the scattering function multiplied by q for di erent values of  $s_{max}$  is shown in Fig.1. The accuracy of the computations is



F igure 1. The plot of qS (q;N ) for chain length N = 1:5. Dots: sti rod; continuous:  $s_{m ax} = 5$ ; dashes:  $s_{m ax} = 8$ .

determ ined by the values of x = qN, where curves corresponding to dimensional error  $s_{max}$  begin to diverge. The Fig.1 shows that for N = 1.5 the continuous and dashed curves begin to diverge for x = 9.

#### 3.4. The extension-force relation

The partition function of a semi exible polymer with one end xed and force fapplied to the another end

$$Z (f; N) = \exp\left(\frac{f}{k_{\rm B}T} \int_{0}^{L_{\rm N}} dst(s)\right)$$
(15)

can be expressed through the distribution function of the end-to-end distance as follows

Z (f;N) = G (k = 
$$i\frac{f}{k_{\rm B}T}$$
;N): (16)

U sing the de nition of the free energy  $F = k_{\rm B} T \ln Z$  (f;N) the extension-force relation can be expressed through the partition function as

$$R = \frac{\theta F}{\theta f} = k_B T \frac{\theta \ln Z (f; N)}{\theta f}$$
(17)

Thus, the short-chain expansion of the extension-force relation is directly obtained from that of the Fourier transform of the distribution function of the end-to-end distance (12). Taking into account the  $\operatorname{rst}_{\Re ax} = 4$  terms in the series in (11) results in the following expression for the extension-force relation

$$\frac{R}{N} = \frac{1}{x} + \coth x \quad \frac{1}{6}N x \quad \frac{1}{6}N \cosh x + \frac{1}{6}N x \coth^{2} x + \frac{1}{90}N^{2} \coth x + \frac{1}{20}N^{2} \cosh x + \frac{1}{20}N^{2} \cosh x \quad \frac{7}{180}N^{2} x \coth^{2} x + \frac{1}{36}N^{2} x^{2} \coth^{3} x + \frac{1}{1512}N^{3} \cosh x + \frac{11}{1512}N^{3} x^{2} \coth x \quad \frac{19}{2520}xN^{3} \quad \frac{1}{120}N^{3} x^{2} \coth^{3} x$$

$$+\frac{1}{840}N^{3}x^{3} - \frac{11}{1890}N^{3}x^{3} \operatorname{coth}^{2}x + \frac{1}{216}N^{3}x^{3} \operatorname{coth}^{4}x + \frac{11}{2520}N^{3}x \operatorname{coth}^{2}x;$$
(18)

where x = fN. The st two terms on the right-hand side of 18 gives the extensionforce relation for sti rod. The log-log plot of 1 R=N for di erent  $s_{ax}$  and for sti rod is shown in Fig.2. The slope for the sti rod is 1, while the slope for the worm-like



Figure 2. Extension-force relation for chain length N = 1:1. Dots: sti rod; continuous:  $s_{m ax} = 4$ ; dashes:  $s_{m ax} = 6$ .

chain approaches the value 1=2, which is the asymptotic result for nite worm-like chain at large forces [30]. Fig 2 shows that the convergence of the short-chain expansion for the extension-force relation is worse than that of the scattering function.

### 4. Conclusions

To conclude, using the exact computation of a large number of moments of the endto-end distribution function G (r;N) of the worm-like chain, which are obtained from the representation of the distribution function as a matrix element of the in nite order matrix [24],[29], we have established the analytical form of the coe cients in Taylor expansions of the moments for short N. The knowledge of these coe cients enabled us to resum mate the moment expansion of G (r;N) by taking into account consecutively the deviations of the moments from their sti rod limit. W ithin this procedure we have derived the short-chain expansion for the distribution function of the end-to-end polymer distance, the scattering function, and the extension-force relation, by taking into account the deviations of the moments from their sti rod limit to the seventh order in N. The procedure can be extended to higher orders. The short-chain expansion could be useful in studies of the behaviour of short polymers, where the deviation from sti rod is small.

#### A cknow ledgm ents

The support from the Deutsche Forschungsgemeinschaft (SFB 418) is gratefully acknowledged.

### References

- [1] S.F.Edwards, Proc.Phys.Soc.85, 613 (1965).
- [2] P.G.de Gennes, Rep. Prog. Phys. 32, 187 (1969).
- [3] P.G. de Gennes, Scaling Concepts in Polymer Physics (Cornell University Press, Ithaca, 1979).
- [4] M. Doi and S.F. Edwards, The Theory of Polymer Dynamics (Clarendon Press, Oxford, 1986).
- [5] J. des C loizeaux and G. Jannink, Polymers in solution, Their modeling, and structure (Oxford University Press, 1990).
- [6] L.Schafer, Excluded Volume E ects in Polymer Solutions (Springer-Verlag, 1999).
- [7] O.Kratky, and G.Porod, Recl. Trav. Chim. Pays-Bas 68, 1106 (1949).
- [8] J.J.Herm ans and R.Ullm an, Physica 18, 951 (1952).
- [9] H.S.Daniels, Proc. Roy. Soc. Edinburgh, 63A, 29 (1952).
- [10] S.Heine, O.K ratky, and J.Roppert, Makrom ol Chem. 56, 150 (1962).
- [11] N.Saito, K.Takahashi, and Y.Yunoki, J.Phys. Soc. Jpn. 22, 219 (1967).
- [12] K.F.Freed, Adv.Chem.Phys.22,1 (1972).
- [13] M.Fixm an and J.J.Kovac, J.Chem. Phys. 58, 1564 (1973).
- [14] J. des C loizeaux, M acrom olecules 6, 403 (1973).
- [15] T.Yoshizakiand H.Yam akawa, Macrom olecules, 13, 1518 (1980).
- [16] M.Wamer, J.M.F.Gunn, and A.B.Baum gartner, J.Phys. A 18, 3007 (1985).
- [17] A.Kholodenko, Ann. Phys. 202, 186 (1990); J.Chem. Phys. 96, 700 (1991).
- [18] H.Yamakawa, Helical W orm like Chains in Polymer Solutions, (Springer, Berlin, 1997).
- [19] J.W ilhelm and E.Frey, Phys.Rev.Lett. 77, 2581 (1996).
- [20] G.Gompper and T.W. Burkhardt, Phys. Rev. A. 40, R6124 (1989); Macrom olecules 31, 2679 (1998).
- [21] E.Frey, K.Kroy, J.W ilhelm, and E.Sackmann, in DynamicalNetworks in Physics and Biology, edited by G.Forgacs and D.Beysens (Springer, Berlin, 1998).
- [22] J.Sam ueland S.Sinha, Phys. Rev. E. 66, 050801 (2002).
- [23] A.D har and D.C haudhuri, Phys. Rev. Lett. 89, 065502 (2002).
- [24] S. Stepanow and G. M. Schutz, Europhys. Lett. 60, 546 (2002).
- [25] R.G.W inkler, J.Chem. Phys. 118, 2919 (2003).
- [26] A.J.Spakow itz and Z-G.W ang, M acrom olecules 37, 5814 (2004).
- [27] G.A.Carriand M.Marucho, J.Chem. Phys. 121, 6064 (2004).
- [28] H. K leinert, Path Integrals in Quantum Mechanics, Statistics, Polymer Physics, and Financial Markets (W orld Scienti c Publishing Co., Singapore, 2004). Chapter IV.
- [29] S.Stepanow, Eur. Phys. J.B 39, 499 (2004).
- [30] J.F.M arko and E.D.Siggia, M acrom olecules 28, 8759 (1995).