

Interacting fermions in two dimensions: beyond the perturbation theory

Suhas Gangadharaiah¹, Dmitrii L. Maslov^{1,2}, Andrey V. Chubukov³, and Leonid I. Glazman⁴¹Department of Physics, University of Florida, P. O. Box 118440, Gainesville, FL 32611-8440²Abdus Salam International Center for Theoretical Physics, 11 Strada Costiera, 34014 Trieste, Italy³Department of Physics, University of Maryland, College Park, MD 20742-4111⁴Theoretical Physics Institute, University of Minnesota, Minneapolis, MN 55455

(Dated: April 14, 2024)

We consider a system of 2D fermions with short-range interaction. A straightforward perturbation theory is shown to be ill-defined even for an infinitesimally weak interaction, as the perturbative series for the self-energy diverges near the mass shell. We show that the divergences result from the interaction of fermions with the zero-sound collective mode. By re-summing the most divergent diagrams, we obtain a closed form of the self-energy near the mass shell. The spectral function exhibits a threshold feature at the onset of the emission of the zero-sound waves. We also show that the interaction with the zero sound does not affect a non-analytic, T^2 -part of the specific heat.

PACS numbers: 71.10.Ay, 71.10.Pm

The Landau Fermi-liquid (FL) theory states that the low-energy properties of an interacting fermion system are similar to those of an ideal Fermi gas [1]. A necessary (but not sufficient) condition for the validity of this theory is that various characteristic properties (thermodynamic parameters, quasi-particle lifetime, etc.) can be expressed via regular perturbative expansions in the interaction. Back in the 50s, it was shown to be the case for a model of 3D fermions with both short- and long-range (Coulomb) repulsion. Soon thereafter, it was realized that the perturbation theory is singular in 1D and, as a result, the FL is destroyed. The case of $D = 2$ had been the subject of an active and fairly recent discussion, with a number of proposals, most notably by Anderson [2], for the breakdown of the FL in 2D. Although the prevailing opinion currently is that the FL is stable in 2D for sufficiently short-range (including Coulomb) interactions, a full description of the FL in 2D is still lacking.

One of the problems in describing a 2D interacting system is that, in a contrast to the 3D case, a naive perturbative expansion in the interaction is singular. Namely, for a linearized single-particle spectrum, the imaginary part of the self-energy diverges on the mass shell. The divergence is logarithmic to second order in the interaction [3, 4, 5, 6] but, as we will show in this paper, it is amplified to a power-law, starting from the third order. This mass-shell singularity in 2D is a weaker form of the "infrared catastrophe" in 1D. There, an on-shell fermion can emit an infinite number of soft bosons (quanta of charge- and spin-density fluctuations [7]), which gives rise to a power-law divergence of the self-energy on the mass-shell. In 2D, this mechanism is weakened but not completely eliminated (the "memory" about the 1D infrared catastrophe is erased completely only for $D > 2$). At a first glance, the breakdown of the perturbation theory confirms the conjecture that the FL is destroyed in 2D [2]. However, all it really means is that in order to obtain physically meaningful results the perturbation the-

ory must be re-summed even for an arbitrarily weak interaction [3, 5].

In this Letter, we report the results of an asymptotically exact re-summation of the divergent perturbation theory for $D = 2$. In addition to obtaining a closed and divergence-free form of the self-energy to all orders in the interaction, this procedure also allows one to identify the nature of the singularities in a perturbation theory as originating from the interaction between the fermions and the zero-sound (ZS) collective mode. At any finite order of the perturbation theory, the collective mode coincides with the upper edge of the particle-hole (PH) continuum. In 2D, this degeneracy is strong enough to generate the divergences in the self-energy. Once perturbations are summed up to all orders, the zero-sound mode splits off from the continuum, and the power-law divergences disappear. The remaining log-divergences are eliminated by restoring the finite curvature of the spectrum near the Fermi surface [5, 6]. Thus the FL survives. However, the resulting self-energy contains several non-perturbative features that signal a deviation from the standard FL behavior. First, the imaginary part of a non-perturbative, ZS contribution to the self-energy, $\text{Im} \Sigma_{\text{ZS}}(\omega; k)$, is a non-monotonic function of the "distance" to the mass-shell, $\omega - \epsilon_k$, and exhibits an anomaly at the threshold for emission of ZS waves. This anomaly gives rise to a non-Lorentzian shape of the spectral function. Second, as the theory is regularized by the difference in the ZS and Fermi velocities, which is small for a weak interaction, $\text{Im} \Sigma_{\text{ZS}}$ is strongly enhanced near the mass shell. On the mass shell, $\text{Re} \Sigma_{\text{ZS}}$ is of the same $\{U^2 \ln |j| \}$ order, as the perturbative contribution. It has been argued that the non-analytic, perturbative self-energy $\text{Re} \Sigma / U^2 \ln |j|$ gives rise to a non-analytic, T^2 -term in the specific heat $C(T)$ [6, 8, 9]. Therefore, it becomes an issue whether the non-perturbative part of the self-energy also contributes to the T^2 -term in $C(T)$. We show that this is not the case, as the thermodynamic




FIG. 3: Scaling functions $F_I(x)$ and $F_R(x)$

imaginary parts of

$$F(x) = (1 + \frac{3}{2}x)^{\frac{1}{2}} \frac{1}{1+x} + \frac{3}{2}x^2 \ln \frac{1 + \frac{1}{2}x}{1-x}; \quad (4)$$

correspondingly. Plots of $F_{R,I}$ are presented in Fig. 3. Notice that $F_I(x < 0) = 0$ and $F_R(x > 1) = 0$:

For large x , i.e., far away from the mass shell, $F_I(x) / 1 = \frac{1}{2}x$, and $\text{Im } \Sigma_S / u_0^2 = \frac{1}{2}x^2$. This behavior reproduces the result of the perturbation theory. Expanding further in $1=x$, one obtains higher powers of $1=x$. However, we see from (4) that the emerging singularity at $x=1$ is actually cut off at $x=1$, i.e., at $x=1$. For small x , $F_I(x) = \frac{3}{2}x^2 = 8$, and $\text{Im } \Sigma_S / u_0^2 = \frac{3}{2}x^2$ for $x=0$. At $x=0$, $\text{Im } \Sigma_S$ vanishes. Vanishing of $\text{Im } \Sigma_S$ on the mass shell is a Cherenkov-type effect: because the zero-sound velocity $c > v_F$, an on-shell fermion cannot emit a ZS boson. The x^2 -behavior of $\text{Im } \Sigma_S$ in between $x=0$ and $x=1$ tracks an increase of the phase space available for the emission of ZS bosons. We emphasize that as $x \rightarrow 0$, the crossover at $x=1$ could have not been obtained within the perturbation theory. Near $x=1$, the derivative of $\text{Im } \Sigma_S(x)$ diverges: $\text{Im } \Sigma_S(x) / 1 = \frac{3}{2}x^2$. The physical meaning of scale

can be understood by noticing that a Cherenkov-type constraint restricts the frequencies of emitted bosons to the range $0 < \omega < (v_F - v)$ (for $v > 0$): This constraint is relevant for $x < 1$; for larger x , emission of bosons with frequencies in the entire interval, allowed by the Pauli principle ($0 < \omega < v_F$) is possible.

The real part of the self-energy varies slowly near the mass shell as $F_R(x-1) = 1+x$. Right on the mass shell, $\text{Re } \Sigma_S / u_0^2 = 1$. Notice that $\text{Re } \Sigma_S$ is of order u_0^2 , although it is obtained by summing up terms of order u_0^3 and higher. This enhancement is due to a non-perturbative cut-off of the mass-shell divergences in the perturbation theory.

The remaining contribution to the forward-scattering part of the self-energy comes from the PH continuum. It contains a logarithmic divergence to second order and also power-law divergences to third and higher orders, as

within the perturbation theory the interaction with the continuum is indistinguishable from the interaction with the zero sound. The logarithmic divergence in the self-energy is cut off by a finite curvature of the spectrum on a scale $\omega^2 = W$, where $W = E_F$ is the bandwidth. The power-law divergences in the PH contribution are cut at the same scale as in the Σ_S , this time because an increase of $U(0)$ near the boundary of the PH continuum reduces the effective interaction for $\omega < W$. However, contrary to the zero-sound contribution, Σ_{PH} is smooth at $x=1$ as no Cherenkov-type condition is involved. Near the mass shell, Σ_{PH} reduces to

$$\begin{aligned} \text{Re } \Sigma_{PH} &= \frac{u_0^2 j! j}{4E_F} \\ \text{Im } \Sigma_{PH} &= \frac{u_0^2 j!^2}{4E_F} \ln \frac{u_0^2 j}{j! (j-E_F)} \quad \text{for } j! j < E_F; \\ &\quad \frac{j! j}{j! (j-E_F)} \quad \text{for } j! j > E_F; \end{aligned}$$

where $j! = u_0^2 E_F$.

The sum $\Sigma_S + \Sigma_{PH}$ is the total contribution to the self-energy from forward scattering. Another contribution to the self-energy comes from the processes in which the fermions move initially in almost opposite directions, and then either scatter by small angle (Fig. 2b) or backscatter (Fig. 2c). We refer to both these processes as "backscattering". The backscattering part of the self-energy, Σ_B , is regular on the mass shell and, for weak interaction, needs to be evaluated only to second order [6]. Taking the result for Σ_B from Ref. [6] and adding it up with the results for Σ_S and Σ_{PH} , we obtain for the total self-energy on the mass shell

$$\begin{aligned} \text{Re } \Sigma(0) &= \frac{j! j}{8E_F} u_0^2 (u_0^2 + u_{2k_F}^2 - u_0 u_{2k_F}) \\ &= \frac{j! j}{8E_F} u_{2k_F} (u_0 + u_{2k_F}); \end{aligned} \quad (5a)$$

$$\text{Im } \Sigma(0) = \frac{u_0^2 j!^2}{2E_F} \ln \frac{E_F}{j! j} - \frac{j! j}{u_{\text{max}}^2 E_F}; \quad (5b)$$

where $u_{2k_F} = m U(2k_F) = 2$ and $u_{\text{max}} = m a v_F$; $u_{2k_F} g$: The scaling function $\Sigma(x)$ approaches the constant values of $1(1=2) + u_{2k_F}(u_{2k_F} - u_0) = (2u_0)$, for $x \rightarrow 1$ and $x \rightarrow 0$; correspondingly.

The first term in the first line of Eq. (5a) is the non-perturbative contribution from forward scattering, whereas the term in square brackets is the backscattering contribution. We see that near the mass shell both contributions are of the same order, i.e., the interaction with the zero-sound collective mode modifies significantly the perturbative result for the self-energy. For contact interaction ($u_0 = u_{2k_F}$), the zero-sound contribution cancels out with the backscattering one, so that the net $\text{Re } \Sigma$ vanishes at the mass shell. At the same time, the effect of the interaction with ZS on the on-shell $\text{Im } \Sigma$ is rather benign: all one has is a smooth crossover function interpolating between two limiting values of the prefactor in




FIG. 4: A log-plot of the spectral function $A(!;k)$ as a function of $x = 2(! - k_F)/u$ for fixed $L_1 = 2$ and $u = 0.1$. A kink at $x = 1$ is a signature of the interaction with the zero-sound mode.

a familiar $!^2 \ln |j|$ dependence. This is the consequence of the fact that the non-perturbative, ZS contribution to Im vanishes on the mass shell.

At the same time, away from the mass shell, the full self-energy and the spectral function $A(!;k) = (-1) \text{Im} G(!;k)$ are affected by the non-monotonic variation of $\text{Im} \Sigma_{\text{ZS}}$ with $!$ near a threshold for emission of ZS bosons. We consider a setup, when $!$ is fixed and the spectral function is measured as a function of k . This is equivalent to varying $x = (k_F - k)/u$ at fixed $!$. Near the mass shell, the spectral function consists of two terms $S_A(x) = A_{\text{qp}}(x) + A_{\text{ZS}}(x)$, where

$$A_{\text{qp}}(x) = \frac{1}{2u^2 E_F} \frac{L_1}{x^2 + 2} \quad (6)$$

is a quasiparticle contribution and

$$A_{\text{ZS}} = \frac{F_I(x)}{2u^2 E_F} \frac{x^2}{(x^2 + 2)^2} \quad (7)$$

is a ZS contribution. We introduced $L_1 = \ln E_F = u^2 |j|$ and $u = |j| \sqrt{L_1} = (2 E_F)$, and assumed for simplicity that $u_0 = u_{k_F} = u$. The ZS contribution $A_{\text{ZS}}(x)$ has a sharp maximum at $x = 1$, where function $F_I(x)$ in Fig. 3 has a peak. This maximum gives rise to a kink in total $A(x)$ (see Fig. 4). A similar consideration shows that the kink is present also for the Coulomb interaction. This kink should be detectable in photoemission experiments on layered materials and in momentum-conserving tunneling between parallel layers of 2D electron gases [12].

Finally, we consider an issue whether non-perturbative effects, discussed above, are relevant for the specific heat of two-dimensional fermions. We argue that they are not. The simplest way to see this is to use the definition, $C(T) = T \partial^2 \ln Z / \partial T^2$, and expand the thermodynamic potential in powers of U . The thermodynamic potential can be evaluated in the Matsubara

formalism, in which the polarization operator $\Pi_m = (m=2) [1 - j_m] / [(v_F Q)^2 + \frac{2}{m}]$ is regular at any m and Q . As a result, the Matsubara series for $C(T)$ converges for a weak, short-range interaction, and there is no need for re-summation of the perturbation theory for $C(T)$. Evaluating $C(T)$ to second-order in U yields

$$C(T) = u_0^2 + u_{2k_F}^2 - u_0 u_{2k_F} \frac{3m}{E_F} \frac{(3) T^2}{E_F} : \quad (8)$$

This result coincides with $C(T)$ obtained in Ref. [6] by finding the perturbative, backscattering part of the self-energy first and then using the relation between $C(T)$ and Π . The agreement between the two results shows that non-perturbative, forward-scattering self-energy does not affect $C(T)$.

For completeness, we also verified explicitly that there is no contribution to $C(T)$ from forward scattering. To this end, one can use the relation between $C(T)$ and the Green's function G which, to first order in $(! - k_F)^{-1} (!;k)$, reads

$$C(T) = \frac{m T}{\partial T} \frac{\partial}{\partial T} \frac{1}{T} \sum_k \frac{Z_1}{d_k} \frac{Z_1}{d_k} \frac{\partial n_0}{\partial !} \quad (9)$$

$$+ \frac{1}{(! - k_F) \text{Re } R(!;k)} \frac{1}{P} \frac{1}{!} \frac{1}{k} \text{Im } R(!;k) :$$

Here $n_0(!)$ is the Fermi function. It is crucial that both $\text{Re } R(!;k)$ and $\text{Im } R(!;k)$ are present in Eq. (9). The forward scattering self-energy, $\Sigma_F = \Sigma_{\text{ZS}} + \Sigma_{\text{PH}}$, depends on k due to the presence of a non-perturbative scale u , and therefore $\text{Im } \Sigma_F$ yields a finite contribution to $C(T)$. Substituting the results for Σ_{ZS} and Σ_{PH} into Eq. (9), we find that the forward scattering contributions to the specific heat from $\text{Re } \Sigma_F$ and $\text{Im } \Sigma_F$ cancel each other; hence, there is no contribution to $C(T)$ from forward scattering.

We acknowledge stimulating discussions with I. A. Akiner, B. A. Itskuler, A. A. Andreev, W. Metzner, A. M. Illis, and C. Pepin. The research has been supported by NSF DMR-0240238 and Condensed Matter Theory Center at UMD (A. V. Ch.), NSF DMR-0308377 (D. L. M.), and NSF DMR-0237296 (L. I. G.).

[*] Permanent address.

[1] A. A. Abrikosov, L. P. Gorkov, and I. E. Dzyaloshinski, Methods of quantum field theory in statistical physics, (Over Publications, New York, 1963).

[2] P. W. Anderson, Phys. Rev. Lett. 65, 2306 (1990).

[3] C. Castellani, C. Di Castro, and W. Metzner, Phys. Rev. Lett. 72, 316 (1994).

[4] H. Fukuyama and M. Ogata, J. Phys. Soc. Jpn. 63, 3923 (1995).

[5] C. Halboth and W. Metzner, Phys. Rev. B 57, 8873 (1998).

- [6] a) A.V.Chubukov and D.L.Maslov, Phys.Rev.B 68, 155113 (2003); b) *ibid.* 69, 121102 (2004).
- [7] Yu.A.Bychkov, L.P.Gor'kov, and I.E.Dzyaloshinskii, Zh.Eksp. Teor. Fiz. 50, 738 (1966) [Sov. Phys. JETP 23, 489].
- [8] D.Coey and K.S.Bedell, Phys. Rev. Lett. 71, 1043 (1993).
- [9] V.M.Galitski and S.DasSarma, Phys. Rev. B 70, 035111 (2004).
- [10] A.V.Chubukov, D.L.Maslov, S.Gangadharaiah, and L.I.Glazman, cond-mat/0412283.
- [11] In principle, there is no guarantee that Eq. (9) works for non-linear terms in $C(T)$. However, it can be verified by a direct comparison with the general Luttinger-Ward formula that the T^2 -term is correctly described by (9) if one uses the T -dependent self-energy rather than the zero-temperature one.
- [12] see S.Q.Murphy, J.P.Eisenstein, L.N.Pfeifer, and K.W.West, Phys. Rev. B 52, 14825-14828 (1995) and references therein.