

Effect of Long-Range Interaction on the Critical Behavior of Three-Dimensional Disordered Systems

S.V. Belin

Omsk State University, 55-a, pr. Mira, Omsk, Russia, 644077 nn

(Dated: March 22, 2024)

A field-theoretical description of the behavior of a disordered Ising system with long-range interaction is presented. The description is performed in the two-loop approximation in three dimensions using the Padé-Borel resummation technique. The renormalization group equations are analyzed, and the fixed points determining the critical behavior of the system are found. It is shown that the effect of frozen structural defects on a system with long-range interaction may cause a change in its critical behavior or smearing of the phase transition.

PACS numbers: 64.60.-i

The effect of long-range interaction, which at long distances is described by the power law r^{-D} , was studied analytically in terms of the ϵ -expansion [13] and numerically by the Monte Carlo method [46] for two- and one-dimensional systems. It was found that the effect of long-range interaction on the critical behavior of Ising systems is considerable for $D < 2$. The study [7] carried out for a three-dimensional space in the two-loop approximation confirmed the prediction of the ϵ -expansion for a homogeneous system with long-range interaction.

According to the results of [8, 9], the introduction of frozen impurities into a system leads to a change in its critical behavior. In this connection, it is of interest to determine the effect of frozen structural defects on the critical behavior of a system with long-range interaction.

This paper describes the critical behavior of a three-dimensional disordered system with a long-range interaction for different values of the parameter ϵ .

The Hamiltonian of a system with long-range interaction has the form

$$H_0 = \frac{1}{2} \sum_{\mathbf{r}} \sum_{\mathbf{r}'} d^D(\mathbf{r} - \mathbf{r}') S_{\mathbf{r}} S_{\mathbf{r}'} + \frac{1}{2} \sum_{\mathbf{r}} \sum_{\mathbf{r}'} d^D(\mathbf{r} - \mathbf{r}') S_{\mathbf{r}} S_{\mathbf{r}'} + u_0 \sum_{\mathbf{r}} d^D(\mathbf{r}) S_{\mathbf{r}} S_{\mathbf{r}} S_{\mathbf{r}} S_{\mathbf{r}} \quad (1)$$

where $S_{\mathbf{r}}$ denotes the fluctuations of the order parameter, D is the space dimension, $d^D(\mathbf{r}) = \int_{-\infty}^{\infty} T_c \delta(T_c - \mathbf{r})$, T_c is the critical temperature, u_0 is a positive constant, and $d^D(\mathbf{r})$ is the random field of impurities of the random temperature type.

The critical behavior strongly depends on the parameter ϵ , which determines the rate of interaction decrease with distance. According to [1], the effect of long-range interaction on the system behavior is considerable for $0 < \epsilon < 2$, while, for $\epsilon > 2$, the critical behavior is equivalent to that of a system with short-range interaction. Therefore, only the case of $0 < \epsilon < 2$ is considered below.

For a low impurity concentration, the distribution of the random field q can be considered as Gaussian and described by the function

$$P[q] = A \exp\left[-\frac{1}{2} \sum_{\mathbf{q}} d^D(\mathbf{q}) q^2\right] \quad (2)$$

where A is the normalization factor and $d^D(\mathbf{q})$ is a positive constant proportional to the concentration of frozen structural defects.

Applying the replica procedure for averaging over random fields caused by the frozen structural defects, we obtain the effective Hamiltonian of the system

$$H_R = \frac{1}{2} \sum_{\mathbf{r}} \sum_{\mathbf{r}'} d^D(\mathbf{r} - \mathbf{r}') \sum_{b=1}^m S_{\mathbf{r}}^b S_{\mathbf{r}'}^b + \frac{1}{2} \sum_{b,c=1}^m \sum_{\mathbf{r}} \sum_{\mathbf{r}'} d^D(\mathbf{r} - \mathbf{r}') (S_{\mathbf{r}}^b S_{\mathbf{r}'}^c) (S_{\mathbf{r}}^c S_{\mathbf{r}'}^b) + u_0 \sum_{b=1}^m \sum_{\mathbf{r}} d^D(\mathbf{r}) S_{\mathbf{r}}^b S_{\mathbf{r}}^b S_{\mathbf{r}}^b S_{\mathbf{r}}^b \quad (3)$$

The properties of the initial system can be obtained in the limit of the number of replicas (transformations) $m \rightarrow 0$.

Applying the standard renormalization group procedure based on the Feynman diagram technique [10] with the propagator $G(\mathbf{k}) = 1/(\mathbf{k}^2 + \mathbf{k}^2)$, we arrive at the expressions for the functions χ , χ' , χ'' , and χ''' , which de-

Electronic address: belin@univer.omsk.su

term in the differential renormalization group equation:

$$\begin{aligned}
 u &= (2-D)u + 1 - 36uJ_0 + 24J_0 \\
 &+ 1728(2J_1 - J_0^2) \frac{2}{9}G u^2 \\
 &+ 2304(2J_1 - J_0^2) \frac{1}{6}G u + 672(2J_1 - J_0^2) \frac{2}{3}G^2; \\
 &= (2-D) - 1 - 24uJ_0 + 16J_0 \\
 &+ 576(2J_1 - J_0^2) \frac{2}{3}G_1 u^2 \\
 &+ 1152(2J_1 - J_0^2) \frac{1}{3}G u + 352(2J_1 - J_0^2) \frac{1}{22}G^2; \\
 t &= (2-D) - 12uJ_0 + 4J_0 + 288(2J_1 - J_0^2) \frac{1}{3}G u^2 \\
 &+ 288(2J_1 - J_0^2) \frac{2}{3}G u + 32(2J_1 - J_0^2) \frac{1}{2}G^2; \\
 \nu &= (2-D)64G(3u^2 - 3u + 2); \\
 J_1 &= \frac{d^D q d^D p}{(1 + \frac{q^2}{Z})^2 (1 + \frac{p^2}{Z})}; \\
 J_0 &= \frac{d^D q}{(1 + \frac{q^2}{Z})^2}; \\
 G &= \frac{\int \frac{d^D q d^D p}{(1 + \frac{q^2}{Z})^2 (1 + \frac{p^2}{Z})}}{\int \frac{d^D q d^D p}{(1 + \frac{q^2}{Z})^2 (1 + \frac{p^2}{Z}) + 2\frac{q^2 p^2}{Z}}};
 \end{aligned}$$

Let us redetermine the effective interaction vertex:

$$v_1 = u - \bar{J}; \quad v_2 = -\bar{J}; \quad (5)$$

As a result, we obtain the following expressions for the functions ν , ν , and t :

$$\begin{aligned}
 \nu &= (2-D) - 1 - 36v_1 + 24v_2 \\
 &+ 1728(2\bar{J}_1 - 1) \frac{2}{9}G v_1^2 \\
 &+ 2304(2\bar{J}_1 - 1) \frac{1}{6}G v_1 v_2 + 672(2\bar{J}_1 - 1) \frac{2}{3}G v_2^2; \\
 \nu &= (2-D) - 1 - 24v_1 + 16v_2 \\
 &+ 576(2\bar{J}_1 - 1) \frac{2}{3}G v_1^2 \\
 &+ 1152(2\bar{J}_1 - 1) \frac{1}{3}G v_1 v_2 + 352(2\bar{J}_1 - 1) \frac{1}{22}G v_2^2; \\
 t &= (2-D) - 12v_1 + 4v_2 + 288(2\bar{J}_1 - 1) \frac{1}{3}G v_1^2 \\
 &+ 288(2\bar{J}_1 - 1) \frac{2}{3}G v_1 v_2 + 32(2\bar{J}_1 - 1) \frac{1}{2}G v_2^2; \\
 \nu &= (2-D)64G(3v_1^2 - 3v_1 v_2 + v_2^2);
 \end{aligned}$$

Such a redetermination makes sense for $D=2$. In this case, J_0 , J_1 and G are divergent functions. Introducing

the cutoff parameter and considering the ratios $\bar{J}=J_0^2$ and $G=J_0^2$ we obtain finite expressions in the limit $\Lambda \rightarrow \infty$.

The values of the integrals were determined numerically. For the case $a=D=2$, a sequence of the values of $\bar{J}_1=J_0^2$ and $G=J_0^2$ was constructed for different and approximated to infinity.

The critical behavior is completely determined by the stable fixed points of the renormalization group transformation. These points can be found by equating the β -functions to zero:

$$\beta_1(v_1; v_2) = 0; \quad \beta_2(v_1; v_2) = 0; \quad (7)$$

The stability requirement for a fixed point reduces to the condition that the eigenvalues b_i of the matrix

$$B_{ij} = \frac{\partial \beta_i(v_1; v_2)}{\partial v_j} \quad (8)$$

lie in the right half-plane of the complex plane.

The critical exponent characterizing the growth of the correlation radius in the vicinity of the critical point (R_c, J^*, T_c) is obtained from the relation

$$= \frac{1}{\nu} (1 + \nu)^{-1};$$

The Fisher exponent describing the behavior of the correlation function in the vicinity of the critical point in the wave-vector space $(G \sim k^{2+\nu})$ is determined from the scaling function $\nu = 2 + \nu(v_1; v_2)$. The values of other critical exponents can be determined from the scaling relations.

It is well known that the perturbation series expansions are asymptotic, and the interaction vertices of the order parameter fluctuations in the fluctuation region are sufficiently large, so that Eqs. (7) applies. Therefore, to extract the desired physical information from the expressions derived above, the Padé-Borel method generalized to the many-parameter case was used. The corresponding direct and inverse Borel transformations have the form

$$\begin{aligned}
 f(v_1; v_2) &= \int_0^\infty dt \sum_{i_1, i_2} C_{i_1, i_2} v_1^{i_1} v_2^{i_2} = \int_0^\infty dt e^{-t} F(v_1 t; v_2 t); \\
 F(v_1; v_2) &= \sum_{i_1, i_2} \frac{C_{i_1, i_2}}{(i_1 + i_2)!} v_1^{i_1} v_2^{i_2};
 \end{aligned} \quad (9)$$

For the analytic continuation of the Borel transform of a function, a series expansion in powers of the auxiliary variable is introduced:

$$F^*(v_1; v_2; \lambda) = \sum_{k=0}^{\infty} \sum_{i_1, i_2} \frac{C_{i_1, i_2}}{k!} v_1^{i_1} v_2^{i_2} \lambda^{i_1 + i_2 + k}; \quad (10)$$

and the $[L/M]$ Padé approximation is applied to this series at the point $\lambda = 1$. This approach was proposed and tested in [11] for describing the critical behavior of systems characterized by several interaction vertices of the order parameter fluctuations. The property (revealed in

	v_1	v_2	b_1	b_2
1.5	-0.395432	0.951135	84.530	59.517
1.6	-0.227628	0.594810	45.302	32.575
1.7	-0.045234	0.274890	13.235	3.915
1.8	0.064189	0.046878	0.626	0.626
1.9	0.066557	0.040818	0.559	0.559

[11-14]) that the system retains its symmetry when using the Padé approximants in the variable v is essential in the description of multivortex models.

The table shows the stable fixed points of the renormalization group transformation and the eigenvalues of the stability matrix at a fixed point for the parameter values $1.5 < v < 1.9$. One can see that stable fixed points exist in the physical region ($v_1, v_2 > 0$) only when the long-range interaction parameter is a 1.8 . The cal-

culations showed that, for any $v < 1.8$, the stable points of a three-dimensional impurity system are characterized by a negative value of the vertex.

The calculation of the critical exponents provided the following values:

$$\begin{aligned} \nu &= 1.9; & \eta &= 0.706786; & \beta &= 0.1344048 \quad (11) \\ \nu &= 1.8; & \eta &= 0.732789; & \beta &= 0.250978 \end{aligned}$$

Thus, for a three-dimensional Ising system with long-range interaction, the introduction of frozen structural defects leads to a change in its critical behavior if the long-range interaction parameter is 1.8 and to smearing the phase transition if the parameter is < 1.8 .

The work is supported by Russian Foundation for Basic Research N 04-02-16002.

-
- [1] M. E. Fisher, S.-K. Ma, and B. G. Nickel, Phys. Rev. Lett. 29, 917 (1972).
[2] J. Honkonen, J. Phys. A 23, 825 (1990).
[3] E. Luijten and H. M. ebtingfeld, Phys. Rev. Lett. 86, 5305 (2001).
[4] E. Bayong and H. T. Diep, Phys. Rev. B 59, 11 919 (1999).
[5] E. Luijten, Phys. Rev. E 60, 7558 (1999).
[6] E. Luijten and H. W. J. Blote, Phys. Rev. B 56, 8945 (1997).
[7] S. V. Belin, Pisma Zh. Eksp. Teor. Fiz. 77, 118 (2003) [JETP Lett. 77, 112 (2003)].
[8] I. O. Mayer, A. I. Sokolov, and B. N. Shalaev, Ferroelectrics 95, 93 (1989).
[9] I. O. Mayer, J. Phys. A 22, 2815 (1989).
[10] J. Zinn-Justin, Quantum Field Theory and Critical Phenomena (Clarendon Press, Oxford, 1989).
[11] S. A. Antonenko and A. I. Sokolov, Phys. Rev. B 49, 15901 (1994).
[12] K. B. Vamashev and A. I. Sokolov, Fiz. Tverd. Tela (St. Petersburg) 38, 3665 (1996) [Phys. Solid State 38, 1996 (1996)].
[13] A. I. Sokolov, K. B. Vamashev, and A. I. Mudrov, Int. J. Mod. Phys. B 12, 1365 (1998).
[14] A. I. Sokolov and K. B. Vamashev, Phys. Rev. B 59, 8363 (1999).