

Effect of Elastic Deformations on the Critical Behavior of Disordered Systems with Long-Range Interactions

S.V. Belin

Omsk State University, 55-a, pr. Mira, Omsk, Russia, 644077 nn

(Dated: April 14, 2024)

A field-theoretic approach is applied to describe behavior of three-dimensional, weakly disordered, elastically isotropic, compressible systems with long-range interactions at various values of a long-range interaction parameter. Renormalization-group equations are analyzed in the two-loop approximation by using the PadBorel summation technique. The fixed points corresponding to critical and tricritical behavior of the systems are determined. Elastic deformations are shown to changes in critical and tricritical behavior of disordered compressible systems with long-range interactions. The critical exponents characterizing a system in the critical and tricritical regions are determined.

PACS numbers: 64.60.-i

The effect of a long-range interaction described by a power law r^{-D} was studied analytically in the framework of the ϵ -expansion [13] and numerically by Monte Carlo methods [46] for two- and one-dimensional systems. It was shown that effects due to long-range interaction are essential for the critical behavior of Ising systems when $D < 2$. The two-loop approximation applied in [7] directly in the three-dimensional space corroborates the predictions of the ϵ -expansion for homogeneous systems with long-range interactions.

In structural phase transitions without piezoelectric effect in the paraphase, elastic deformations play the role of a secondary order parameter whose fluctuations are not critical in most cases [8, 9]. Since the main contribution to striction effects in the critical region is due to the distance dependence of the exchange integral, only isotropic elastic systems are discussed below.

It was shown in [10, 11] that coupling between order-parameter fluctuations and elastic deformations can be responsible both for change in critical behavior and for emergence of tricritical and tetracritical points. The introduction of frozen point impurities into a system both changes the critical behavior and eliminates multicritical points [12]. The analysis presented in [13] showed that frozen structural defects introduced into spin systems with long-range interactions increased the value of a long-range parameter corresponding to the transition to mean-field critical behavior. The effect of elastic deformations on homogeneous systems with long-range interactions also changes the critical behavior [14]. Therefore, it would be interesting to analyze the combined effect of structural defects and elastic deformations on systems with long-range interactions.

In this work, the critical behavior of disordered compressible systems with long-range interactions in the three-dimensional space is described for various values of the long-range parameter.

The Hamiltonian of the disordered Ising model including elastic deformations and long-range interactions can be written as

$$\begin{aligned}
 H_0 = & \frac{1}{2} \sum_{\mathbf{q}} d^D q (t_0 + q) S_{\mathbf{q}} S_{-\mathbf{q}} + \frac{1}{2} \sum_{\mathbf{q}} d^D q_{\mathbf{q}} S_{\mathbf{q}} S_{-\mathbf{q}} \\
 & + u_0 \sum_{\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3} d^D q_{\mathbf{q}_1} S_{\mathbf{q}_1} S_{\mathbf{q}_2} S_{\mathbf{q}_3} \\
 & + a_3 \sum_{\mathbf{q}_1, \mathbf{q}_2} d^D q_{\mathbf{q}_1} S_{\mathbf{q}_1} S_{\mathbf{q}_2} \\
 & + \frac{a_3^{(0)}}{Y_0} \sum_{\mathbf{q}} d^D q S_{\mathbf{q}} S_{-\mathbf{q}} + \frac{1}{2} a_1 \sum_{\mathbf{q}} d^D q Y_{\mathbf{q}} Y_{-\mathbf{q}} \\
 & + \frac{1}{2} \frac{a_1^{(0)}}{Y_0^2} \sum_{\mathbf{q}} d^D q h_{\mathbf{q}} Y_{\mathbf{q}} + \frac{h_0}{Y_0} Y_0
 \end{aligned} \tag{1}$$

where $S_{\mathbf{q}}$ is an order parameter, u_0 is a positive constant, $t_0 = T - T_c$, T_c is the phase-transition temperature, $d^D q$ is the long-range parameter, $q_{\mathbf{q}}$ is a random impurity field (e.g., random temperature), a_1 and a_2 are the elastic constants of a crystal, and a_3 is the quadratic striction constant. The coupling between impurities and the non-fluctuating order parameter $y(\mathbf{x}) = \sum_{\mathbf{q}} u_{\mathbf{q}}(\mathbf{x})$, where $u_{\mathbf{q}}$ is the strain tensor, is specified by the random field $h_{\mathbf{q}}$ thermodynamically conjugate to $u_{\mathbf{q}}(\mathbf{x})$. In Eq. (1), integration is performed in the terms depending on non-fluctuating variables that are not coupled to the $S_{\mathbf{q}}$, and the Y_0 terms (describing homogeneous deformations) are separated. It was shown in [8] that this separation is necessary because inhomogeneous deformations $y_{\mathbf{q}}$ are responsible for exchange of acoustic phonons and for long-range effects that are absent under homogeneous deformations. When the impurity concentration is low, random fields $q_{\mathbf{q}}$, $h_{\mathbf{q}}$ and h_0 can be treated as

Electronic address: belin@univer.omsk.su

Gaussian and specified by the function

$$P[\mathbf{h}; \mathbf{h}_0] = A \exp \left[\frac{1}{8b_1} \sum_q d^D q \frac{1}{8b_2} \sum_q h_q^2 d^D q \frac{1}{8b_3} \sum_q h_0 d^D q \frac{1}{4b_4} \sum_q q h_q d^D q \frac{1}{4b_5} \sum_q q h_0 d^D q \right]; \quad (2)$$

where A is a normalization factor and b_i denotes positive constants proportional to the concentration of frozen structural defects.

Using the replica procedure to average over the random fields generated by frozen structural defects, we obtain an effective Hamiltonian of the system:

$$\begin{aligned} H_R = & \frac{1}{2} \sum_q d^D q (v_0 + q) \sum_{a=1}^{X^n} S_q^a S_q^a \\ & + \frac{v_0}{2} \sum_{a,b=1}^{X^n} \sum_q d^D q (S_{q1}^a S_{q2}^a) (S_{q3}^b S_{q1}^b S_{q2}^b S_{q3}^b) \\ & + u_0 \sum_{a=1}^{X^n} \sum_q d^D q S_{q1}^a S_{q2}^a S_{q3}^a S_{q1}^a S_{q2}^a S_{q3}^a \\ & + g_0 \sum_{a=1}^{X^n} \sum_q d^D q y_{q1}^a S_{q2}^a S_{q1}^a \\ & + \frac{g_0^{(0)}}{2} \sum_{a=1}^{X^n} y_0^a \sum_q d^D q S_q^a S_q^a + \frac{1}{2} \sum_q d^D q y_q y_q + \frac{1}{2} y_0^2 \end{aligned} \quad (3)$$

where the positive constants $v_0, g_0, g_0^{(0)}$, and y_0 can be expressed in terms of a_i and b_i . The properties of the original system can be obtained in the limit $m \rightarrow 0$, where m is the number of replicas (images).

Define an effective Hamiltonian depending only on the strongly fluctuating order parameter S_q by the relation

$$\exp \int d^D y H[\mathbf{S}; \mathbf{y}] = B \int d^D y \exp \int d^D y H_R[\mathbf{S}; \mathbf{y}] \quad (4)$$

If an experiment is carried out at constant volume, then y_0 is a constant and integration in Eq. (4) is performed over inhomogeneous deformations, whereas homogeneous deformations do not contribute to the effective Hamiltonian. At constant pressure, the Hamiltonian is modified by adding the term P , volume is represented in terms of the strain tensor components as

$$= v_0 \left[1 + \sum_{i=1}^X u_i + \sum_{i,j=1}^X u_i u_j + O(u^3) \right] \quad (5)$$

and integration over homogeneous deformations is also performed in Eq. (4). It was noted in [15] that the quadratic terms in Eq. (5) can be important at high pressures and for crystals with largestriction effects. The

result is

$$\begin{aligned} H = & \frac{1}{2} \sum_q d^D q (v_0 + q) \sum_{a=1}^{X^n} S_q^a S_q^a \\ & + (u_0 - \frac{z_0}{2}) \sum_{a=1}^{X^n} \sum_q d^D q f_{q1} g_{q1}^a S_{q1}^a S_{q2}^a S_{q3}^a S_{q1}^a S_{q2}^a S_{q3}^a \\ & - \frac{1}{2} \sum_{a,b=1}^{X^n} \sum_q d^D q f_{q1} g_{q1}^a (S_{q1}^a S_{q2}^a) (S_{q3}^b S_{q1}^b S_{q2}^b S_{q3}^b) \\ & + \frac{1}{2} (z_0 - w_0) \sum_{a=1}^{X^n} \sum_q d^D q f_{q1} g_{q1}^a (S_{q1}^a S_{q1}^a) (S_{q2}^a S_{q2}^a); \\ & z_0 = g_0^2; \quad w_0 = g_0^{(0)2} = 0; \end{aligned} \quad (6)$$

The effective coupling constant $v_0 = u_0 - z_0/2$, which arises in the Hamiltonian owing to thestriction effects determined by g_0 , can have positive and negative values. Therefore, this Hamiltonian describes both first and second-order phase transitions. When $v_0 = 0$, the system exhibits tricritical behavior. Furthermore, the effective interaction determined by the difference $z_0 - w_0$ in Eq. (6) can also change the order of the phase transition. This representation of the effective Hamiltonian entails the existence of a higher order critical point where tricritical curves intersect when $v_0 = 0$ and $z_0 = w_0$ [16]. Note that Hamiltonian (6) is isomorphic to the Hamiltonian of the disordered Ising model with long-range interactions under the tricritical condition $z_0 = w_0$.

Using the standard renormalization-group procedure based on the Feynman diagram technique [17, 18] with the propagator $G(\mathbf{k}) = 1/(\mathbf{k}^2 + \mathbf{k}^2)$, we derive the following expressions for the functions v, z, w, y , and t specifying the differential renormalization-group equation:

$$\begin{aligned} v = & (2 - D) v_0 - 36 v J_0 + 24 J_0 \\ & + 1728 (2J_1 - J_0^2) \frac{2}{9} G v^2 \\ & - 2304 (2J_1 - J_0^2) \frac{1}{6} G v \\ & + 672 (2J_1 - J_0^2) \frac{2}{3} G^2 v^3; \\ w = & (2 - D) w_0 - 24 v J_0 + 16 J_0 \\ & + 576 (2J_1 - J_0^2) \frac{2}{3} G_1 v^2 \\ & - 1152 (2J_1 - J_0^2) \frac{1}{3} G v + 352 (2J_1 - J_0^2) \frac{1}{22} G^2 v^3; \end{aligned}$$

$$\begin{aligned}
z &= (2-D)z_1 - 24vJ_0 - 2zJ_0 + 8J_0 \\
&+ 576(2J_1 - J_0^2) \frac{2}{3}G v^2 \\
&+ 120(2J_1 - J_0^2) \frac{8}{5}G v + 96(2J_1 - J_0^2) \frac{2}{3}G^2 v^{\frac{1}{2}}; \\
w &= (2-D)w_1 - 24vJ_0 + 8J_0 - 4zJ_0 + 2wJ_0 \\
&+ 576(2J_1 - J_0^2) \frac{2}{3}G v^2 \\
&+ 120(2J_1 - J_0^2) \frac{8}{5}G v + 96(2J_1 - J_0^2) \frac{2}{3}G^2 v^{\frac{1}{2}}; \\
t &= (2-D) - 12vJ_0 + 4J_0 - 2zJ_0 + 2wJ_0 \\
&+ 288(2J_1 - J_0^2) \frac{1}{3}G v^2 \\
&+ 288(2J_1 - J_0^2) \frac{2}{3}G v + 32(2J_1 - J_0^2) \frac{1}{2}G^2 v^{\frac{1}{2}}; \\
r &= (2-D)64G(3v^2 - 3v + 1); \\
J_1 &= \frac{d^D q d^D p}{(1 + \frac{q^2}{J_0})^2 (1 + \frac{p^2}{J_0}) (1 + \frac{q^2 + p^2}{J_0} + 2\frac{qp}{J_0})}; \\
J_0 &= \frac{d^D q}{(1 + \frac{q^2}{J_0})^2}; \\
G &= \frac{d^D q d^D p}{(1 + \frac{q^2}{J_0}) (1 + \frac{p^2}{J_0} + k^2 + 2\frac{qp}{J_0}) (1 + \frac{q^2 + p^2}{J_0} + 2\frac{qp}{J_0})};
\end{aligned} \quad (7)$$

In terms of the new effective interaction vertexes

$$v_1 = v - \bar{g}; \quad v_2 = -\bar{g} \quad v_3 = z - \bar{g}; \quad v_4 = w - \bar{g}; \quad (8)$$

the functions z , r , and t are expressed as

$$\begin{aligned}
z &= (2-D)z_1 - 36v_1 + 24v_2 \\
&+ 1728(2\bar{g}_1 - 1) \frac{2}{9}\bar{g} v_1^2 - 2304(2\bar{g}_1 - 1) \frac{1}{6}\bar{g} v_1 v_2 \\
&+ 672(2\bar{g}_1 - 1) \frac{2}{3}\bar{g} v_2^2; \\
w &= (2-D)w_1 - 24v_1 + 8v_2 \\
&+ 576(2\bar{g}_1 - 1) \frac{2}{3}\bar{g} v_1^2 - 1152(2\bar{g}_1 - 1) \frac{1}{3}\bar{g} v_1 v_2 \\
&+ 352(2\bar{g}_1 - 1) \frac{1}{2}\bar{g} v_2^2; \\
t &= (2-D)t_1 - 24v_1 + 16v_2 - 2v_3 \\
&+ 576(2\bar{g}_1 - 1) \frac{2}{3}\bar{g} v_1^2 - 120(2\bar{g}_1 - 1) \frac{8}{5}\bar{g} v_1 v_2 \\
&+ 96(2\bar{g}_1 - 1) \frac{2}{3}\bar{g} v_2^2;
\end{aligned} \quad (9)$$

$$\begin{aligned}
t &= (2-D)t_1 - 24v_1 + 8v_2 - 4v_3 + 2v_4 \\
&+ 576(2\bar{g}_1 - 1) \frac{2}{3}\bar{g} v_1^2 - 120(2\bar{g}_1 - 1) \frac{8}{5}\bar{g} v_1 v_2 \\
&+ 96(2\bar{g}_1 - 1) \frac{2}{3}\bar{g} v_2^2; \\
r &= (2-D) - 12v_1 + 4v_2 - 2v_3 + 2v_4 \\
&+ 288(2\bar{g}_1 - 1) \frac{1}{3}\bar{g} v_1^2 - 192(2\bar{g}_1 - 1) \frac{2}{3}\bar{g} v_1 v_2 \\
&+ 32(2\bar{g}_1 - 1) \frac{1}{2}\bar{g} v_2^2; \\
r &= (2-D)64\bar{g}(3v_1^2 - 3v_1 v_2 + v_2^2);
\end{aligned}$$

This definition is meaningful for $D=2$. In this case, J_0 , J_1 and G are divergent functions. Introducing the cutoff parameter Λ , we obtain finite expressions for the ratios $J_1=J_0^2$ and $G=J_0^2$ as $\Lambda \rightarrow 1$.

The integrals are performed numerically. For $D=2$, a sequence of $J_1=J_0^2$, $G=J_0^2$ corresponding to various values of Λ is calculated and extrapolated to infinity.

It is well known that perturbation-theory series are asymptotic and expressions (9) cannot be applied directly since the interaction vertexes for fluctuations of order parameters in the fluctuation region are too large. For this reason, the required physical information was extracted from these expressions by applying the Padé-Borel method extended to the four-parameter case. The appropriate direct and inverse Borel transforms have the form

$$\begin{aligned}
f(v; z; w) &= \sum_{i_1, \dots, i_4}^P C_{i_1, \dots, i_4} v_1^{i_1} v_2^{i_2} v_3^{i_3} v_4^{i_4} \\
&= \int_0^R e^{-t} F(v_1 t; v_2 t; v_3 t; v_4 t) dt; \\
F(v; z; w) &= \sum_{i_1, \dots, i_4}^P \frac{C_{i_1, \dots, i_4}}{(i_1 + \dots + i_4)!} v_1^{i_1} v_2^{i_2} v_3^{i_3} v_4^{i_4};
\end{aligned}$$

To obtain an analytic continuation of the Borel transform of a function, a series in an auxiliary variable k is introduced:

$$F(v; z; w; k) = \sum_{k=0}^X \sum_{i_1, \dots, i_4}^X \frac{C_{i_1, \dots, i_4}}{k!} v_1^{i_1} v_2^{i_2} v_3^{i_3} v_4^{i_4} k^{i_1 + \dots + i_4}; \quad (10)$$

The Padé approximant $[L/M]$ is applied to this series at $k=1$. This procedure was proposed and tested in [19-22] for describing the critical behavior of a number of systems characterized by several interaction vertexes for order-parameter fluctuations. It was found in [19-22] that Padé approximants in the variable k preserve the symmetry of the system. This property is important for analysis of multivertex models. The approximants [2/1] are used to calculate the functions in the two-loop approximation.

Critical behavior is completely determined by the stable fixed points of the renormalization group transformation. These points can be found from the condition

$$v_i(v_1; v_2; v_3; v_4) = 0 \quad (i=1; 2; 3; 4); \quad (11)$$

The requirement of stability of a fixed point reduces to the condition that the eigenvalues b_i of the matrix

$$B_{i,j} = \frac{\partial^2 \Phi(v_1, v_2, v_3, v_4)}{\partial v_i \partial v_j} \quad (12)$$

are positive.

The exponent characterizing the growth of the correlation radius near the critical point (R_c, T_c) is determined by the relation

$$\nu = \frac{1}{d - \beta} (1 + \beta)^{-1} :$$

The Fisher exponent describing the behavior of the correlation function near the critical point in the wave-vector space $(G \propto k^{2+\nu})$ is determined by the scaling function $\nu = 2 + \nu(v_1, v_2, v_3, v_4)$. The remaining critical exponents can be determined from the scaling relations.

The table shows the stable fixed points of the renormalization group transformation and the fixed-point eigenvalues of the stability matrix for $\nu = 1.8$ and 1.9 . It was shown in [13] that stable fixed points exist in the physical region ($v_i > 0$) for disordered systems only when $\nu < 1.8$. When $\nu > 1.8$, the stable points of any three-dimensional impurity system are characterized by negative values of the vertex v_i .

An analysis of the critical points and their stability shows that the fixed critical points of disordered systems with long-range interactions (nos. 1 and 4) are unstable under elastic deformations. The critical behavior of disordered compressible systems with long-range interactions is described by their respective fixed points (nos. 2 and 5). The phase diagram of the substance can contain tricritical points specified by fixed points (nos. 3 and 6).

Calculations of the critical exponents for the stable fixed points (nos. 2 and 5) yield

$$\begin{aligned} \nu &= 1.9; & \beta &= 0.721; & \gamma &= 0.134 \\ \nu &= 1.8; & \beta &= 0.758; & \gamma &= 0.251 \end{aligned} \quad (13)$$

For the tricritical points (nos. 3 and 6), the critical exponents are

$$\begin{aligned} \nu &= 1.9; & \beta &= 0.686; & \gamma &= 0.134 \\ \nu &= 1.8; & \beta &= 0.721; & \gamma &= 0.251 \end{aligned} \quad (14)$$

Thus, elastic deformations change both critical and tricritical behavior of three-dimensional disordered Ising systems with long-range interactions.

The work is supported by Russian Foundation for Basic Research N 04-02-16002.

-
- [1] M. E. Fisher, S.-K. Ma, and B. G. Nickel, Phys. Rev. Lett. 29, 917 (1972).
 [2] J. Honkonen, J. Phys. A 23, 825 (1990).
 [3] E. Luijten and H. M. Eisingfeld, Phys. Rev. Lett. 86, 5305 (2001).
 [4] E. Bayong and H. T. Diep, Phys. Rev. B 59, 11 919 (1999).
 [5] E. Luijten, Phys. Rev. E 60, 7558 (1999).
 [6] E. Luijten and H. W. J. Bloote, Phys. Rev. B 56, 8945 (1997).
 [7] S. V. Belin, Pisma Zh. Eksp. Teor. Fiz. 77, 118 (2003) [JETP Lett. 77, 112 (2003)].
 [8] A. I. Larkin and S. A. Pikin, Zh. Eksp. Teor. Fiz. 56, 1664 (1969) [Sov. Phys. JETP 29, 891 (1969)].
 [9] D. J. Bergman and B. I. Halperin, Phys. Rev. B 13, 2145 (1976).
 [10] V. M. Laptev and A. V. Skryabin, Phys. Status Solidi B 91, K 143 (1997).
 [11] Y. N. Skryabin and A. V. Shchanov, Phys. Lett. A 234, 147 (1997).
 [12] S. V. Belin and V. V. Prudnikov, Fiz. Tverd. Tela (St. Petersburg) 43, 1299 (2001) [Phys. Solid State 43, 1353 (2001)].
 [13] S. V. Belin, Pisma Zh. Eksp. Teor. Fiz. 77, 509 (2003) [JETP Lett. 77, 434 (2003)].
 [14] S. V. Belin, Pisma Zh. Eksp. Teor. Fiz. 77, 659 (2003) [JETP Lett. 77, 556 (2003)].
 [15] M. A. de Moura, T. C. Lubensky, Y. Imry, and A. Aharony, Phys. Rev. B 13, 2176 (1976).
 [16] Y. Imry, Phys. Rev. Lett. 33, 1304 (1974).
 [17] D. Amit, Field Theory, the Renormalization Group, and Critical Phenomena (McGraw-Hill, New York, 1976).
 [18] J. Zinn-Justin, Quantum Field Theory and Critical Phenomena (Clarendon Press, Oxford, 1989).
 [19] S. A. Antonenko and A. I. Sokolov, Phys. Rev. B 49, 15 901 (1994).
 [20] K. B. Vamashev and A. I. Sokolov, Fiz. Tverd. Tela (St. Petersburg) 38, 3665 (1996) [Phys. Solid State 38, 1996 (1996)].
 [21] A. I. Sokolov, K. B. Vamashev, and A. I. Mudrov, Int. J. Mod. Phys. B 12, 1365 (1998).
 [22] A. I. Sokolov and K. B. Vamashev, Phys. Rev. B 59, 8363 (1999).

N	v_1	v_2	v_3	v_4	b_1	b_2	b_3	b_4
= 1.8								
1	0.064189	0.046878	0	0	0.626	0.626	-0.123	-0.123
2	0.064189	0.046878	0.066101	0	0.626	0.626	0.124	0.125
3	0.064189	0.046878	0.066101	0.066101	0.626	0.626	0.124	-0.124
= 1.9								
4	0.066557	0.040818	0	0	0.559	0.559	-0.118	-0.118
5	0.066557	0.040818	0.065716	0	0.559	0.559	0.119	0.119
6	0.066557	0.040818	0.065716	0.065716	0.559	0.559	0.119	-0.119