

1=f noise in experimental Sinai quantum billiards

E. Faleiro¹, U. Kuhf², R. A. Molina³, A. Relano⁴, J. Retamosa⁴, and H.-J. Stockmann²

¹ Departamento de Física Aplicada, E. U. I. T. Industrial,
Universidad Politécnica de Madrid,
E-28012 Madrid, Spain

² Fachbereich Physik, Philipps-Universität, Renthof 5,
D-35032 Marburg, Germany

³ Max-Planck-Institut für Physik Komplexer Systeme
Nothnitzer Str. 38, D-01187 Dresden, Germany

⁴ Departamento de Física Atomica, Molecular y Nuclear,
Universidad Complutense de Madrid,
E-28040 Madrid, Spain

It was recently conjectured that $1/f$ noise is a fundamental characteristic of spectral fluctuations in chaotic quantum systems. In this Letter we show that the level fluctuations of experimental realizations of the Sinai billiard exhibit $1/f$ noise, corroborating the conjecture. Assuming that the statistical properties of these systems are those of the Gaussian orthogonal ensemble (GOE), we compare the experimental results with the universal behavior predicted in the random matrix framework, and find an excellent agreement. The deviations from this behavior observed at low frequencies can be easily explained with the semiclassical periodic orbit theory. In conclusion, it is shown that the main features of the conjecture are affected neither by non-universal properties due to the underlying classical dynamics nor by the uncertainties of the experimental setup.

PACS numbers: 05.40.-a, 05.45.Tp, 05.45.Mt, 05.45.Pq

Quantum chaos is a well established subject since the pioneering work of Bohigas and collaborators [1] (for further review see, for example, [2, 3]). Recently, a different approach to quantum chaos has been proposed [4]. Considering the sequence of energy levels as a discrete time series, and the appropriate statistic related to the fluctuations of the excitation energy, it was conjectured that chaotic quantum systems are characterized by $1=f$ noise, whereas integrable ones exhibit $1=f^2$ noise. Since chaotic spectra behave like antipersistent time series, and regular ones as neither persistent nor antipersistent time series, it was possible to establish a relationship between spectral rigidity and antipersistence. This conjecture is supported by numerical calculations which involve atomic nuclei and the classical Random Matrix Ensembles (RME) [4]. Later, theoretical expressions were derived in Ref. [5] using random matrix theory. They explain, without free parameters, the universal behavior of the excitation energy fluctuations power spectrum. The theory gives excellent agreement with numerical calculations. It also reproduces the $1=f$ power law characteristic of chaotic systems and the $1=f^2$ law characteristic of integrable system through several orders of magnitude of the frequency interval.

All these results were obtained in the Random Matrix Theory (RMT) and Nuclear Shell Model frameworks. These models allow an accurate statistical analysis since very long sequences can be obtained with high precision; moreover, RMT is the usual theoretical model in studying quantum chaos, and the high excited states of the atomic nuclei follow RMT predictions almost perfectly [6, 7]. However, other quantum chaotic systems cannot provide such quality of data. Experiments usually provide short level sequences with some precision problems: if two resonances are separated by a distance smaller than the experimental line width, they are registered as one; in microwave billiard experiments, certain resonances are missing if the emitting antenna is positioned close to a nodal line. Quantum systems with a classical analogue can provide very long sequences, but they present some non-universal features that spoil the universal behavior of long-range correlations [8, 9], which reveal some dynamic properties of the classical analogue (the behavior of the shorter periodic orbits of the system). Therefore, analyzing whether experimental limitations and the existence of short periodic orbits modify our previous result becomes an important issue.

In this Letter we study the spectral fluctuations of experimentally modeled quantum Sinai billiards. We find that the excitation energy level fluctuations of these systems are in clear agreement with the theoretical expressions found in Ref. [5]. Therefore, they exhibit $1=f$ noise, in spite of small deviations due to experimental limitations and non-universal properties due to short periodic orbits. Moreover, we present a simple analysis of these anomalies using periodic orbit theory.

In the present work, quantum billiards have been substituted by suitably shaped microwave resonators, whose eigenfrequencies are measured. This kind of experiment uses the fact that the time-independent Schrödinger equation and the wave equation are mathematically equivalent. The data we analyze in this work has already been used in Refs. [10, 11], although in a different context.

To be specific, we have used a quarter Sinai billiard varying the values of the length l as depicted in Fig. 1. The




FIG. 1: Sketch of the quarter Sinai billiard with varying length printed in scale. The length of the billiard was varied from $l = 480 \text{ mm}$ to 460 mm in steps of 0.2 mm . The width $b = 200 \text{ mm}$ and the radius of the quarter circle $r = 70 \text{ mm}$.

experimental setup makes it possible to measure resonances in the frequency range 14.5 to 15.5 GHz , where there are about 120 eigenvalues for each value of the parameter l . We have considered about 100 of such sequences as realizations of a stochastic process representative of the billiard, as we have previously done in numerical experiments [4].

Prior to the statistical analysis, we have to remove the modulation induced by the mean level density in the spectra. This procedure is called unfolding; it consists in locally mapping the actual energy levels E_i into new dimensionless levels ϵ_i

$$\epsilon_i = \frac{E_i - E_0}{E_N - E_0} \quad ; \quad i = 1, 2, \dots, N \quad (1)$$

where N is the dimension of the spectrum and $\overline{N}(E)$ is a smooth approximation to the actual accumulated density. Quite generally, $\overline{N}(E)$ can be calculated with the aid of the following rule: in the limit of large E , the average density of eigenstates is proportional to the volume of the phase space with energy less than E [12]. For quantum billiards we can use an improved version of this result, which is also valid near of the ground-state [9]:

$$\overline{N}(E) = \overline{N}^{\text{Weyl}}(E) + \overline{N}^{\text{bb}}(E) \quad (2)$$

$\overline{N}^{\text{Weyl}}$ is the so called Weyl law; it includes a main term related to the volume of the phase space, and two correction terms related to the surface and the curvature of the boundary [13]. $\overline{N}^{\text{bb}}(E)$ is a further correction term which takes into account the smooth trend produced by a set of non-isolated marginally stable periodic orbits bouncing between the two straight sides of the billiard.

After performing the unfolding procedure, we can calculate the nearest neighbor level spacings defined by $s_i = \epsilon_{i+1} - \epsilon_i$; $i = 1, 2, \dots, N-1$; since the unfolded spectrum has a mean level density equal to 1 , their average value is $\overline{s} = 1$. To study the long-range correlation structure, we use the χ_n statistic [4, 14] defined by

$$\chi_n = \frac{1}{N} \sum_{i=1}^N (s_i - \overline{s})^n \quad ; \quad n = 2, 3, \dots, N-1 \quad (3)$$

This statistic represents the deviation of the excitation energy of the $(n+1)$ th unfolded level from its mean value n , and it is closely related to the spectrum fluctuations. Indeed, we can write $\chi_n = \overline{N}^n(E_{n+1})$ if we appropriately shift the ground state energy; thus, χ_n represents the cumulated level density fluctuations. If we consider the index n as the analogue of a discrete time [16], we can analyze χ_n with numerical techniques normally used in the study of complex systems. The most simple is the analysis of the power spectrum obtained from the discrete Fourier transform (DFT) [17] of the signal

$$b_k = \frac{1}{N} \sum_{n=0}^{N-1} \chi_n \exp \left(\frac{2\pi i k n}{N} \right) \quad ; \quad k = 0, 1, \dots, N-1 \quad (4)$$

as

$$S(|k|) = |b_k|^2 \quad (5)$$

where

$$\Delta_k = \frac{2\pi k}{N} \quad (6)$$

and Δ is the sampling interval of the signal.

Here we face the following problem. The experiment cannot provide all energy levels in the interval 14.5 and 15.5 GHz: there are a few missing eigenvalues due to nodal lines of the wavefunction at the position of the antenna. Nevertheless, due to the level dynamics measurement, one can detect where a certain level is missing. A comparison with eq. (2) shows that all levels are accounted for. Thus, we are able to arrange the known energy levels as $f_{p_1}; f_{p_2}; \dots; f_{p_M}$, where $M (N)$ represents the total number of levels whose energy is known, and $f_{p_1}; f_{p_2}; \dots; f_{p_M}$ are their positions in the spectrum. Therefore, the n_j function can only be calculated for certain whole numbers. Taking f_{p_1} as a fictitious ground state, we have

$$n_j = f_{p_{j+1}} - f_{p_1} \quad n_j; n_j = p_{j+1} - p_1; j \in [0; M-1]; \quad (7)$$

Consequently, n_j is not a standard time series, but an unevenly spaced time series.

There are some obvious ways to get an evenly sampled signal from the unevenly spaced p_j points. Interpolation is one possible way, but generally speaking, interpolation techniques perform poorly. Therefore, we have decided to proceed in a different way to analyze the experimental data. For an arbitrarily sampled time series the power spectrum can be estimated by the periodogram, $S(\Delta_k) = \frac{1}{M} |b(\Delta_k)|^2$, where $b(\Delta_k)$ is given by

$$b(\Delta_k) = \frac{1}{M} \sum_{j=0}^{M-1} n_j \exp(-i \Delta_k n_j) \quad (8)$$

Eq. (8) is defined for any frequency Δ_k . However, since the original signal is discrete, one should be able to choose a finite set of frequencies to evaluate the Fourier transform, and then using the values of eq. (8) at these frequencies, recover the original signal. For evenly spaced time series of length N eq. (6) provides a natural set of frequency values. This set has been built from a fundamental frequency $\Delta_1 = 2\pi/N$, which defines a sine wave of period equal to the whole interval length; the remaining frequencies are multiples of it. As commented above, the set $f_{p_k}(\Delta_k)$, where $k \in [0; N-1]$, is known as the Discrete Fourier transform of the signal.

In our case we have generated an appropriate set of frequencies as follows:

(i) Since we know the smooth part of the level density (even if there are some missing levels), we can perform the unfolding of the spectrum, leading to a constant average level density. Thus, we can set the fundamental sampling interval to $\Delta = 1$.

(ii) Given a window $N_a - N_b$, we select all the experimental billiards for which there is at least a pair of measured levels at positions p_a and p_b , satisfying that $N = p_b - p_a$.

(iii) We consider the following frequency set $\Delta_k = 2\pi k/N$ with $k = 1; 2; \dots$. This is probably the best choice because the fundamental frequency, $\Delta_1 = 2\pi/N$, corresponds to a sine wave of period equal to the whole energy interval $p_b - p_a$ (Remember that after the unfolding procedure $\langle s \rangle = 1$.)

Due to the definition of our periodogram and to the selected frequency set, we have that $S(\Delta_k + 2\pi) = S(\Delta_k - 2\pi) = S(\Delta_k)$. For this reason the power spectrum is defined in the basic interval $[0; \Delta_k]$. For evenly sampled series the frequency $\Delta_k = \Delta_k/2$ is known as the Nyquist frequency and a change of trend in the power spectrum can be observed near this upper limit.

(iv) Finally, the average power spectrum of the n_j signal is calculated using the selected billiards, and the experimental results are compared with the theoretical expression found in Ref. [5]. Assuming that $N \gg 1$, and that the statistical properties of the spectral fluctuations are those of the Gaussian orthogonal ensemble (GOE), the standard model for time-reversal invariant systems, the universal behavior of the n_j power spectrum is given by

$$\langle S(\Delta_k) \rangle = \frac{K}{\Delta_k^2} + \frac{1}{2} \frac{K}{(\Delta_k)^2} + \frac{1}{4 \sin^2 \frac{\Delta_k}{2}}; \quad \Delta_k \in (0; \Delta_k/2) \quad (9)$$

Here,

$$K_{GOE}(\Delta_k) = \frac{2}{\Delta_k} \frac{\log(2 + \Delta_k)}{\log \frac{2 + \Delta_k}{\Delta_k}}; \quad \Delta_k > 1 \quad (10)$$

is the spectral form factor of G O E , whose analytical expression is well known [14]. The constant shift reflects that ρ_n is not a continuous two-point measure (defined for every energy interval), but a discrete function (defined only for each energy level). For G O E $\rho_n = 1/2$, provided that the whole sequence of ρ_n values is included in the calculation of the power spectrum. In this case, we cannot evaluate ρ_n for all n , but only for some energy levels. This feature entails that the value of ρ_n should be affected by the number M of levels whose energy is known. When $\rho_n \rightarrow 0$, eq. (9) reduces to

$$\langle S(\omega_k) \rangle = \frac{1}{\omega_k}; \quad \omega_k \rightarrow 0; \quad (11)$$

showing that for small frequencies the excitation energy fluctuations exhibit $1/f$ noise in G O E -like systems.




FIG. 2: Averaged power spectra of the ρ_n function for the Sinai microwave billiards (filled circles) compared with the theoretical prediction (solid line) as given by eq. (9). The dashed straight line is the $1/f$ noise predicted by the approximated expression (11). Vertical arrows indicate the frequencies of the two shortest periodic orbits of the billiard.

In the forthcoming analysis we have chosen $N = 85$ since it is large enough to obtain significant results, and it is possible to find measured levels at positions p_a and p_b fulfilling $p_b - p_a = 85$ in almost every sequence of levels; in fact there are 97 sequences satisfying this condition. Using a log-log scale, Fig. 2 compares the average power spectrum of ρ_n and the theoretical predictions of eq. (9) with $\rho_n = 1/2$. This value was obtained by means of a least-squares fit to the numerical data. In spite of the experimental limitations, the agreement between the numerical and experimental results is very good. Moreover, it is clearly seen that the spectral fluctuations of these systems show a $1/f$ power law. The significant power damping observed at lower frequencies can be understood by a simple analysis of the contribution of the bouncing ball orbits to the power spectrum. If we consider $\langle l \rangle = 470$ mm and $\langle b \rangle = 200$ mm as the average length of the straight sides of the billiards, the frequencies corresponding to these orbits are $\omega = 0.09$ and $\omega = 0.21$. These values define approximately the interval where the power is significantly diminished; therefore, we can conclude that this phenomenon is due to the removal of the contribution of bouncing ball orbits to the level density. All this is consistent with the idea that long period classical orbits give rise to an universal behavior for integrable and ergodic systems regardless of the system specificities [18]. Only when the main contribution comes from the scarce short periodic orbits, deviations from the universal behavior might be observed.

In summary, we have studied the spectral fluctuations of quantum Sinai billiards modeled experimentally. We have considered the ρ_n statistic, that characterizes the fluctuations of the excitation energy, as a discrete time series. Using about one hundred different lengths of the straight sides of the billiard, the average power spectrum of ρ_n was calculated. Recall that performing an average is very important to reduce the fluctuations and clarify the secular trend of the ρ_n function. The main result of this analysis is that the spectral fluctuations of these systems exhibit a clear $1/f$ noise through almost the whole frequency domain. We have also compared the theoretical predictions of eq. (9) for G O E -like systems with these results, finding an excellent agreement except in the lower frequency region. Nevertheless, a simple study reveals that these deviations from the universal behavior predicted by RMT are due to the removal of the bouncing ball orbits during the unfolding procedure; therefore there is no contribution to the power spectrum in the low frequency domain. Last but not least, we can say that ρ_n statistic is not affected by the experimental limitations, therefore we can conclude that it is a quite strong statistic.

Prior to this work, we have found 1=f noise in energy spectrum fluctuations for random matrix ensembles and atomic nuclei; these results allowed us to conjecture that 1=f characterize spectral fluctuations of chaotic quantum systems. The new results obtained in this Letter corroborate this conjecture and give us a hint on the non-universal features that can appear in the power spectrum at lower frequencies due to the short periodic orbits.

This work is supported in part by Spanish Government grants BFM 2003-04147-C02 and FTN 2003-08337-C04-04. The experiments have been supported by the Deutsche Forschungsgemeinschaft via several grants.

-
- [1] O. Bohigas, M. J. Giannoni and C. Schmit, Phys. Rev. Lett. 52, 1 (1984).
 - [2] T. Guhr, A. Müller-Groeling and H. A. Weidenmüller, Phys. Rep. 299, 189 (1998).
 - [3] H. J. Stockmann, Quantum Chaos, (Cambridge University Press, Cambridge, 1999).
 - [4] A. Relano, J. M. G. Gómez, R. A. Molina, J. Retamosa, and E. Faleiro, Phys. Rev. Lett. 89, 244102 (2002).
 - [5] E. Faleiro, J. M. G. Gómez, L. Muñoz, R. A. Molina, A. Relano, and J. Retamosa, Phys. Rev. Lett. 93, 244101 (2004).
 - [6] R. U. Haq, A. Pandey, O. Bohigas, Phys. Rev. Lett. 48, 1086 (1982).
 - [7] Phys. Rev. C 69, 057302 (2004); and references therein.
 - [8] M. V. Berry, Proc. R. Soc. Lond. A 400, 229 (1985).
 - [9] H. D. Graf et al, Phys. Rev. Lett. 69, 1296 (1992).
 - [10] M. Barth, U. Kuhl, and H.-J. Stockmann, Phys. Rev. Lett. 82, 2026 (1999).
 - [11] M. Barth, U. Kuhl, and H.-J. Stockmann, Ann. Phys. (Leipzig) 8, 733 (1999).
 - [12] M. C. Gutzwiller, Chaos in Classical and Quantum Mechanics, Springer-Verlag, New York (1990).
 - [13] R. Balian and C. Bloch, Ann. Phys. 60, 441 (1970).
 - [14] M. L. Mehta, Random Matrices, (Academic Press, 1991)
 - [15] O. Bohigas, P. Leboeuf and M. J. Sanchez, Physica D 131, 186 (1999).
 - [16] The analogy of the dimensionless n index of ρ_n and a time leads to a quite confusing terminology, because there is also a physical time (the period of each classical periodic orbit) which plays a fundamental role in semiclassical theory. If we do not state explicitly the opposite, we shall use the word frequency to refer to the conjugate variable in the ρ_n Fourier transform.
 - [17] J. D. Scargle, Ap. J. Suppl. 451 (1981), Ap. J. 263, 835 (1982) and Ap. J. 343, 874 (1989).
 - [18] J. H. Hannay and A. M. Ozorio de Almeida, J. Phys. A 17, 3429 (1984).