

Bosonic Resonating Valence Bond wave function for doped Mott insulators

Z. Y. Weng

Center for Advanced Study, Tsinghua University, Beijing 100084, China

Y. Zhou

Max-Planck-Institut für Physik komplexer Systeme
Noethnitzer Str. 38, 01187 Dresden, Germany

V. N. Muthukumar

Department of Physics,
City College at the City University of New York,
New York, NY 10031
(Dated: today)

We propose a class of ground states for doped Mott insulators in the electron second quantization representation. They are obtained from a bosonic resonating valence bond (RVB) theory of the t - J model. At half filling, the ground state describes spin correlations of the $S=1/2$ Heisenberg model very accurately. Its spin degrees of freedom are characterized by RVB pairing of spins, the size of which decreases continuously as holes are doped into the system. Charge degrees of freedom emerge upon doping and are described by twisted holes in the RVB background. We show that the twisted holes exhibit an off-diagonal long range order (ODLRO) in the pseudogap ground state, which has a finite pairing amplitude, but is short of phase coherence. Unpaired spins in such a pseudogap ground state behave as free vortices, preventing superconducting phase coherence. The existence of nodal quasiparticles is also ensured by such a hidden ODLRO in the ground state, which is non Fermi liquid-like in the absence of superconducting phase coherence. Two distinct types of spin excitations can also be constructed. The superconducting instability of the pseudogap ground state is discussed and a d -wave superconducting ground state is obtained. This class of pseudogap and superconducting ground states unifies antiferromagnetism, pseudogap, superconductivity, and Mott physics into a new state of matter.

I. INTRODUCTION

An important question posed by the study of high temperature superconductors is whether the superconducting ground state of doped Mott insulators can be described by a BCS-like wave function. A d -wave BCS wave function and/or its derivatives have been widely used in the phenomenology of high temperature superconductors, partly because electron pairing, as evidenced by 2π flux quantization and similar experimental results, is observed in the cuprate superconductors. However, owing to the effect of strong on-site Coulomb repulsion between the electrons, superconductivity in these systems may be quite different. Whereas in a conventional BCS superconductor, screening and retardation effects serve to minimize the role of Coulomb interaction, the role of the latter becomes crucial in a doped Mott insulator where the charge degrees of freedom are partially frozen by the severity of the Coulomb interaction. Strong arguments have been presented in the literature that a high temperature superconductor evolves from a Mott insulator doped by holes, and consequently the ground state can be very different from a conventional BCS state¹.

A simple step towards incorporating Mott physics into a BCS description was suggested many years ago^{1,2}. This involves a projection (out of the Hilbert space) of doubly occupied sites in a d -wave BCS state³. In this approach, the charge degrees of freedom on a lattice site i , are partially frozen by a Gutzwiller operator $(1 - n_{i\uparrow}n_{i\downarrow})$. At half filling, only spin degrees of freedom survive, and the wave function describes a Mott insulating state. This class of wave functions has achieved some success in the phenomenology of the cuprate superconductors^{4,5,6}. A recent paper argues the case further⁷. However, the treatment of low energy spin degrees of freedom in this description is not satisfactory. Experimentally, strong antiferromagnetic (AF) correlations have been observed for the Cu spins, but projected d -wave BCS states cannot account correctly for such correlations. Since we opine that spin correlations are intimately related to superconductivity in a doped Mott insulator, we are motivated to seek an alternate description in which the spin degrees of freedom can be treated more accurately and systematically.

In this paper, we propose a new class of ground states based on a microscopic treatment of the t - J model. These ground state wave functions are different from the projected BCS wave functions, and possess the following properties: (i) the spin degrees of freedom can fully restore the AF long range order (AFLRO) as well as low-lying spin wave excitations at half filling, and evolve into a spin liquid with strong short-range AF correlations at low doping; (ii) the no double occupancy constraint is always satisfied for any doping, and is incorporated self-consistently instead

where $e^{i\hat{\phi}}$ is a nonlocal operator describing the phase string effect. We call a hole that is created by c_i a twisted hole, and as will be shown in this paper, such a twisted hole is a bosonic object whose kinetic energy is optimized in the (modified) RVB background. We shall obtain a ground state of the following form

$$|j_G\rangle = \sum_1 \prod_{i=1}^{N_h} Z_{\#}(1) c_{i\#}^{\dagger} \sum_1 \prod_{i=1}^{N_h} Z_{\#}(1) c_{i\#}^{\dagger} |\text{RVB}\rangle; \quad (2)$$

where $Z_{\#}(1)$ describes the wave function of the twisted hole c_i , and $N_h^{\#} + N_h^{\#} = N_h$ is the total number of holes. The modified RVB vacuum $|\text{RVB}\rangle$ has the same structure as $|j_0\rangle$ at half-filling, except that the RVB pairing amplitudes, W_{ij} 's, become complex and change continuously with doping, which can be determined self-consistently as we shall show.

The ground state (2) is one of the central results of this paper. It generalizes the wave function (1) written down by Liang et al., to the doped case. We will derive (2) from a microscopic treatment of the t - J model, based on a reformulation of the model in the so-called phase string representation.

The wave function (2) differs from the Gutzwiller projected BCS wave functions in a fundamental way. Spin correlations with various length scales emerge self-consistently as a function of doping in $|\text{RVB}\rangle$ and their interplay with the charge degrees of freedom is incorporated through the nonlocal phase string operator $e^{i\hat{\phi}}$ in c_i . We will find that the ground state (2) describes both an AF insulator (at zero doping) and a pseudogap phase with short range AF correlations (at finite doping) in which two distinct kinds of spin excitations emerge. We will identify a hidden ODLRO in this pseudogap ground state which ensures a finite pairing amplitude and nodal fermionic quasiparticles. Although superconducting phase coherence of this state is prevented by the presence of free spins which behave like vortices, there exists an inherent superconducting instability at finite doping, which can naturally give rise to d-wave superconductivity.

The outline of the paper follows. In Section II, we reformulate the t - J model using the phase string representation. This is an exact reformulation and allows us to sort out the effects of hole motion in an AF background discussed earlier. A singular gauge transformation automatically ensures that the single occupancy constraint is satisfied. A brief review of this formalism and important physical consequences will be discussed in Section II. In Section III, we obtain the ground state within a mean-field theory, and in Section IV we transform the ground state wave function back to the electron representation, thus completing the derivation of the wave function (2). A discussion of the properties predicted by (2) follows and finally a superconducting ground state is obtained. The final section is devoted to a general discussion about the region of validity (in the phase diagram) of this approach, as well as the questions that this study raises. In Appendix A, some detailed derivations of the mean-field theory in the phase string representation are presented. In Appendix B, we discuss briefly, an effective theory of spin and charge degrees of freedom that are coupled by the phase string. In Appendix C, we show that the ground state obeys time reversal and spin rotational symmetries.

II. PHASE STRING FORMULATION AND PHYSICAL IMPLICATIONS

In this section, we reformulate the t - J model using the phase string representation¹¹. This representation, like the slave-particle representations used in the literature^{2,13}, is an exact reformulation. However, the phase string formulation has a distinct advantage over these other schemes, in the following sense. The most singular effects encapsulated in the t - J model are sorted out by a unitary transformation, and all nontrivial phases present in the original model now occur as topological gauge fields. Without these gauge fields, the model becomes trivial, i.e., free of sign problems, which is the case in one dimension, or in the particular case of half-filling in two dimensions. The central issue in two dimensions is then how to handle these topological gauge fields at finite doping. It turns out that they are well controlled, at least for small doping.

A. Phase string representation

The most singular effect of the hole moving in an AF background is the phase string effect. This leads to a competition between hopping and superexchange processes. Since it is most singular at length scales of a lattice constant (as Z_2 signs), it is crucial to track this effect explicitly, before constructing any effective theory. As shown by Weng et al., this can be achieved starting from a slave-fermion representation and by performing a unitary transformation of the basis states¹¹. In this procedure, the electron operator is decomposed into

$$c_i = h_i^\dagger b_i e^{i\hat{\phi}_i} \quad (3)$$

where holon h_i^y and spinon b_i operators are both bosonic fields, satisfying the constraint

$$h_i^y h_i + \sum_l b_l^y b_l = 1 : \quad (4)$$

The phase factor $e^{i\hat{\phi}_i}$ in Eq. (3) is defined by

$$e^{i\hat{\phi}_i} = e^{i\phi_i^{string}} (\pm)^{\hat{N}_h} (-1)^{\hat{S}_i}; \quad (5)$$

where \hat{N}_h is the (total) holon number operator, the phase string operator $\phi_i^{string} = \frac{1}{2} \sum_l s_l - \frac{1}{2} \sum_l h_l$ is a nonlocal operator with

$$s_i = \sum_l \frac{1}{\mathcal{L}} \frac{1}{|z_l - z_i|} \sum_{l \in \mathcal{L}} n_l^b; \quad (6)$$

$$h_i = \sum_l \frac{1}{\mathcal{L}} \frac{1}{|z_l - z_i|} \sum_{l \in \mathcal{L}} n_l^h; \quad (7)$$

and

$$h_i = \sum_l \frac{1}{\mathcal{L}} \frac{1}{|z_l - z_i|} \sum_{l \in \mathcal{L}} n_l^h; \quad (8)$$

Here, n_l^b and n_l^h are spinon and holon number operators respectively, at site l ; and $z_l = x_l + iy_l$ is a complex coordinate on the lattice.

It is easily verified that the fermionic statistics of c_i is automatically ensured by $e^{i\phi_i^{string}}$; besides ensuring fermionic statistics, the string operator also incorporates the singular effects of hole motion in the RVB background, as will be discussed later. The factor $(\pm)^{\hat{N}_h}$ guarantees anticommutation relations between opposite spins. The factor $(-1)^{\hat{S}_i} = \pm 1$ in (5) is added for convenience, and incorporates the Marshall sign into the decomposition (3) of the electron in terms of the holon and spinon operators, which may be regarded as a bosonization scheme for the electron moving in an RVB background. The same bosonization decomposition can be also obtained from the slave-boson representation^{14,15}.

B. The t-J model in the phase string representation

Rewriting the t-J model using the electron decomposition (3), we get $H_{t-J} = H_t + H_J$; where the hopping term

$$H_t = \sum_{\langle ij \rangle} \sum_{\sigma} e^{iA_{ij}^s - i\phi_{ij}^0} h_i^y h_j e^{iA_{ji}^h} b_j^y b_i + \text{h.c.}; \quad (9)$$

and the superexchange term

$$H_J = \sum_{\langle ij \rangle} \frac{J}{2} \hat{S}_{ij}^x \hat{S}_{ij}^y : \quad (10)$$

The RVB pair operator \hat{S}_{ij}^s in H_J is defined by

$$\hat{S}_{ij}^s = \sum_{\sigma} e^{iA_{ij}^h} b_i^{\sigma} b_j^{\sigma} : \quad (11)$$

The unique feature in this formalism is the emergence of three phases (link fields): A_{ij}^s , ϕ_{ij}^0 , and A_{ij}^h ; defined on the nn links. Without these, there should be no nontrivial sign problem in H_{t-J} , because h and b are both bosonic fields. The matrix elements of H_{t-J} would then be real and negative-definite in the occupation number representation of h and b . Consequently, the ground state expanded in terms of these bosonic fields will have real and positive coefficients. Therefore, any nontrivial signs (phases) of the t-J model arise solely from these three link fields, defined by

$$A_{ij}^s = \frac{1}{2} \sum_l \frac{1}{|z_l - z_i|} \sum_{l \in \mathcal{L}} [s_l - s_j] \sum_{l \in \mathcal{L}} n_l^b; \quad (12)$$

$$A_{ij}^0 = \frac{1}{2} \sum_{\langle i,j \rangle} [A_i(l) - A_j(l)] ; \quad (13)$$

and

$$A_{ij}^h = \frac{1}{2} \sum_{\langle i,j \rangle} [A_i(l) - A_j(l)] n_l^h ; \quad (14)$$

It is easy to see that the Hamiltonian H_{t-J} is invariant under $U(1) \times U(1)$ gauge transformations:

$$h_i \rightarrow h_i e^{i\theta_i} ; \quad A_{ij}^s \rightarrow A_{ij}^s + (\theta_i - \theta_j) ; \quad (15)$$

and

$$b_i \rightarrow b_i e^{i\phi_i} ; \quad A_{ij}^h \rightarrow A_{ij}^h + (\phi_i - \phi_j) ; \quad (16)$$

Thus A_{ij}^s and A_{ij}^h are gauge fields, seen by holons and spinons respectively, as the latter carry their gauge charges according to (15) and (16). But they are not conventional $U(1)$ gauge fields, as they must satisfy the topological constraints: $\oint_C A_{ij}^s = \frac{1}{2} \sum_{\langle i,j \rangle \in C} n_{i\#}^s - n_{j\#}^s$ and $\oint_C A_{ij}^h = \frac{1}{2} \sum_{\langle i,j \rangle \in C} n_l^h$ according to (12) and (14), respectively. Here the subscript c denotes an arbitrary closed loop such that the fluxes enclosed, $\oint_C A_{ij}^s$ and $\oint_C A_{ij}^h$, are determined by the number of spinons and holons respectively, inside the loop c ; viz., A_{ij}^s and A_{ij}^h are not independent gauge fields with their own dynamics. Rather they are directly connected to matter fields as a pair of mutual Chern-Simons gauge fields¹⁶. The term "mutual" refers to the fact that A_{ij}^s describes quantized vortices attached to spinons, coupled to the holons in H_t . Conversely, A_{ij}^h describes quantized vortices bound to holons, coupled to spinons in H_J . The phase A_{ij}^0 describes a uniform and constant flux with a strength per plaquette: $\frac{1}{2} \sum_{\langle i,j \rangle} A_{ij}^0 = \frac{1}{2} \sum_{\langle i,j \rangle} 1$ according to (13).

In the case of the one dimensional model, $A_{ij}^s = A_{ij}^h = 0$: Thus there is no sign problem in the phase-string representation of the $t-J$ model. It implies that the Hamiltonian can be treated within a mean field framework, which indeed results in a correct Luttinger-liquid solution for the large- U Hubbard model¹¹. In two dimensions, these non-link phases can no longer be "gauged away". They compose a mutual gauge structure which completely determines the essential sign problem of the $t-J$ model. These gauge fields are generally well controlled in the regimes of our interest: A_{ij}^0 is a non-dynamic phase describing a constant flux per plaquette; A_{ij}^s is canceled when spinons are RVB paired at low temperature phases; A_{ij}^h remains weak at small doping - it vanishes at half filling, where there is no sign problem in the Hamiltonian. Therefore, these gauge fields in the phase string representation are presumably well tractable in low doping and low temperature regimes. This is one of the main advantages of the phase string representation over other approaches.

Finally, it is important to note that the phase string representation is defined in a Hilbert space where the total S^z is an eigen operator. The total numbers of \uparrow and \downarrow spinons are conserved individually, such that the topological gauge field A_{ij}^s behaves smoothly as defined in (12), depicting circulating vortices bound to spinons. (Non-conserved S^z would result in non-conserved \uparrow and \downarrow vortices in A_{ij}^s). Different S^z states are connected by the spin flip operators, defined in the phase-string representation as

$$S_i^\pm = \left(\frac{1}{2} \right)^{\frac{1}{2}} e^{i\theta_i} b_{i\#}^\pm ; \quad (17)$$

(a factor $(-1)^{N_h}$ has been dropped for simplicity) and $S_i = (S_i^+)^{\dagger}$, and $S_i^z = \frac{1}{2} \sum_{\langle i,j \rangle} b_i^\dagger b_j$. These definitions follow from (3). The nonlocal phase θ_i in (17) plays a crucial role in restoring the spin rotational symmetry.

C. No double occupancy constraint

Since the $t-J$ model is a projective Hamiltonian, the no double occupancy constraint is a central issue. It imposes a severe restriction on the Hilbert space

$$\sum_i c_i^\dagger c_i = 1 ; \quad (18)$$

which ensures that the system is a Mott insulator at half filling and a doped Mott insulator away from half filling. In conventional slave particle decompositions, the constraint reduces to a single occupancy constraint like (4), and

is still difficult to treat. Typically, the constraint is relaxed at the level of mean field theory, and restoring its effect beyond the mean field theory (say, by a gauge theory) is a very challenging problem. In contrast, the phase string representation outlined in the previous subsection provides a new way to handle the constraint.

A wavefunction ψ_e in the electron c-operator representation is defined in a quantum state as

$$\psi_e = \sum_{f_{i_u} g f_{j_d} g}^X e(f_{i_u} g; f_{j_d} g) c_{i_1}^Y c_{i_2}^Y \dots c_{j_M}^Y c_{j_1 \#}^Y c_{j_2 \#}^Y \dots c_{j_N}^Y c_{\#}^Y \psi_i; \quad (19)$$

where the $\#$ spin electron sites, $f_{i_u} g = i_u; \dots_M$; and the $\#$ spin sites, $f_{j_d} g = j_d; \dots_{N_e} j_d$; obey the no double occupancy constraint in the restricted Hilbert space of the tight-J Hamiltonian. On the other hand, in the phase string representation, the wave function $\psi_b(i_1; \dots_M; j_1; \dots_{N_e} j_1; \dots_{N_h} j_1; \dots_{N_h})$ is defined as

$$\psi_b = \sum_{f_{i_u} g f_{j_d} g}^X b(f_{i_u} g; f_{j_d} g; f_{l_h} g) \bar{\psi}(i_u g; f_{j_d} g; f_{l_h} g); \quad (20)$$

where the indices $f_{l_h} g = l_h; \dots_{N_h}$ denote the empty sites, that are not independent from $f_{i_u} g$ and $f_{j_d} g$ under the constraint. Here the spinon-holon basis in the phase string representation, $\bar{\psi}(i_u g; f_{j_d} g; f_{l_h} g)$ is given by

$$\bar{\psi}(i_u g; f_{j_d} g; f_{l_h} g) = (-1)^{N_A} b_{i_1}^Y b_{i_2}^Y \dots b_{j_M}^Y c_{j_1 \#}^Y c_{j_2 \#}^Y \dots c_{j_N}^Y b_{\#}^Y h_{i_1} h_{i_2} \dots h_{j_N} h_{\#} \psi_i; \quad (21)$$

where the vacuum ψ_i is chosen to be filled by holons. The sign factor $(-1)^{N_A}$ in (21) can be identified with the Marshall sign, and N_A denotes the total number of $\#$ spins in sublattice A. Here and henceforth, we will always use i to specify an $\#$ spin, j a $\#$ spin, and l a holon, where the subscripts u , d , and h label the sequences of the $\#$ spins, $\#$ spins, and holons, respectively.

Now, the decomposition (3) of the electron operator c_i , relates the wave functions written in the two representations:

$$c_{i_1}^Y c_{i_2}^Y \dots c_{j_M}^Y c_{j_1 \#}^Y c_{j_2 \#}^Y \dots c_{j_N}^Y c_{\#}^Y \psi_i = K^{-1} \bar{\psi}(i_u g; f_{j_d} g; f_{l_h} g); \quad (22)$$

where

$$\begin{aligned} K &= \prod_{u,d} \frac{z_{i_u}}{z_{j_d}} \prod_{u < u^0} \frac{z_{i_u}}{z_{j_d}} \prod_{d < d^0} \frac{z_{j_d}}{z_{j_d^0}} \prod_{u,h} \frac{z_{i_u}}{z_{j_d}} \prod_{h < h^0} \frac{z_{j_d}}{z_{j_d^0}} \prod_{u,h} \frac{z_{i_u}}{z_{j_d}} \prod_{d,h} \frac{z_{j_d}}{z_{j_d^0}} \dots \\ &= C^{-1} \prod_{u,d} (z_{i_u} z_{j_d}) \prod_{u < u^0} (z_{i_u} z_{j_d^0}) \prod_{d < d^0} (z_{j_d} z_{j_d^0}) \prod_{h < h^0} z_{j_d} z_{j_d^0} \prod_{u,h} (z_{i_u} z_{j_d}) \prod_{d,h} z_{j_d} z_{j_d^0} : \quad (23) \end{aligned}$$

The coefficient C is given by

$$C = \prod_{u,d} \frac{z_{i_u}}{z_{j_d}} \prod_{u < u^0} \frac{z_{i_u}}{z_{j_d}} \prod_{d < d^0} \frac{z_{j_d}}{z_{j_d^0}} \prod_{h < h^0} \frac{z_{j_d}}{z_{j_d^0}} \prod_{u,h} \frac{z_{i_u}}{z_{j_d}} \prod_{d,h} \frac{z_{j_d}}{z_{j_d^0}} \dots$$

in which k and m run through all lattice sites, such that C is a constant.

Correspondingly, the wave functions in the electron and phase string representations are related by

$$\psi_e(i_1; \dots_M; j_1; \dots_{N_e} j_1) = K \psi_b(i_1; \dots_M; j_1; \dots_{N_e} j_1; l_1; \dots_{N_h} l_1); \quad (24)$$

which holds generally for the tight-J model.

Note that the above expression for K has been obtained strictly under the no double occupancy constraint. But if the constraint is relaxed in (23), while still treating C as a constant, then one finds $K = 0$ for double occupancies when any two particles (spinons and/or holons) occupy the same site; viz., K defined in (23) automatically enforces the constraint (4) through the Jastrow-like factors. Therefore, as far as ψ_e is concerned, the no double occupancy constraint in ψ_b or $\psi_b = C$ is no longer important, since K in (24) naturally plays the role of a projection operator.

Clearly, an exact ψ_b would satisfy the single occupancy constraint (4). But, the point is that an approximate ψ_b determined without the constraint does not affect ψ_e , owing to K . It means that in the phase string representation the constraint (4) is indeed "unimportant", which may be understood in the following way. In the phase string representation, the effect of K in the original ψ_e , is transformed into the topological gauge fields A_{ij}^s and A_{ij}^h ; in the Hamiltonians, (9) and (10), which describe spinons and holons as mutual vortices, as perceived by each other

(cf. the discussion in the previous subsection). This clearly implies a mutual repulsion between two species, since a spinon cannot stay at the center of its vortex (which is a holon), and vice versa. Thus the constraint that a holon and a spinon cannot occupy the same site is now reflected in the interactions present in the new Hamiltonian, and the condition (4) is not needed as an extra condition to enforce. Note that the constraint (4) also requires the hard core conditions among the holons or spinons themselves. But since both holon and spinon fields are bosonic fields, local hard core exclusions usually do not involve the sign change of the wave function. Hence, in the phase string representation, the local constraint (4) is neither crucial nor singular, as far as low energy physics is concerned.

D. Phase string effect: A singular doping effect

In this subsection, we point out that the other important effect, i.e., the phase string effect, is also explicitly built into this representation via K .

Let us first rewrite K (23) in a more compact form,

$$K = J G ; \quad (25)$$

where

$$J = \prod_{u < u^0}^Y (z_{i_u} / z_{i_{u^0}}) \prod_{d < d^0}^Y (z_{j_d} / z_{j_{d^0}}) \prod_{u d}^Y (z_{i_u} / z_{j_d}) \prod_{h < h^0}^Y z_{i_h} / z_{j_h} \prod_{u h}^Y z_{i_u} / z_{j_h} \prod_{d h}^Y z_{j_d} / z_{i_h} ; \quad (26)$$

and

$$G = C^{-1} \prod_{u h}^Y \frac{z_{i_u} / z_{j_h}}{z_{j_u} / z_{i_h}} ; \quad (27)$$

It is easily seen that the Jastrow-like factors in J enforce the single occupancy constraint discussed previously: J vanishes if two spins (or holes) occupy the same site, or if a hole and a spin occupy the same site. The factor J also explicitly captures the fermionic statistics of electrons.

Now let us focus on the additional factor G in K , which is asymmetric with regard to " and # spins: it only involves an " spin complex coordinate z_{i_u} and a holon coordinate z_{j_h} . Suppose that the holon is taken along a closed loop shown in Fig. 1. The K factor will generally acquire a nontrivial phase through G . In the following we examine this.

For a spinon inside or outside the loop, there will be no net contribution to G as the acquired phase is either 2π or 0 under such an operation. So any nontrivial contribution is solely from spinons occupying lattice sites on the holon's loop-path. At each step of hopping, the holon under consideration will have to switch positions with a spinon occupying a nn site on the loop, obeying the no double occupancy constraint. Then G picks up a minus sign from the Jastrow factor in G every time when the holon and an " spinon are exchanged. Thus, if one changes the holon's coordinate continuously through the loop without encountering other holons, G will acquire a sequence of signs,

$$G \rightarrow (-1)^{N_{\text{loop}}} G ; \quad (28)$$

where N_{loop} denotes the number of " spinons exchanged with the holon during the above thought experiment. (Note that the spinons displaced by holon hopping during the operation should all be restored to the original configuration by pure spin flips in terms of the superexchange H_J , and this does not generate any further signs in G).

Therefore, the electron wave function ψ_e acquires a sign factor (which can also be thought of as a Berry phase) through G :

$$(-1)^{N_{\text{loop}}} = \prod_{\text{loop}}^Y (z_{i_m} / z_{j_m}) = \prod_{\text{loop}}^Y z_{i_m} / z_{j_m} ; \quad (29)$$

This factor (for the closed loop) is just the phase string shown in Fig. 1, where i_m denotes the index of spins on the loop exchanged with a holon, under the thought experiment of moving the holon around a closed path (note that a closed loop on a bipartite lattice always involves an even number of nn links). Equation (29) shows that the singular hopping effect is explicitly incorporated in ψ_e through K , in the phase string formulation of the $t-J$ model.

To conclude this section, the two singular effects in the $t-J$ model, viz., the single occupancy constraint and the phase string effect, are explicitly represented by the factor K , (23). Consequently the ground state wave function in the phase string representation, ψ_b ; is presumably no longer singular, justifying a mean field theory.

III. CONSTRUCTION OF THE GROUND STATE IN PHASE STRING REPRESENTATION

A mean field theory (bosonic RVB theory) based on the phase string representation of the t - J model was constructed in an earlier paper¹⁷, and we will follow a similar route here in dealing with the spin degrees of freedom. However, an important distinction in this work is the way the charge degrees of freedom are incorporated, in order to better facilitate and optimize the kinetic energy of the holes, as well as to construct a state in which the no double occupancy constraint is strictly enforced. As shown in Sec. IV, this construction directly results in a pseudogap ground state characterized by a finite pairing amplitude and the emergence of nodal quasiparticles. The ground state has an inherent superconducting instability which occurs, when phase coherence is realized.

A. Spin degrees of freedom and bosonic RVB order parameter

Let us begin by considering the superexchange term H_J , which is expressed in terms of an RVB pair operator \hat{s}_{ij}^s in (10). Note that \hat{s}_{ij}^s is invariant under the gauge transformation (16). It is then natural to define the bosonic RVB order parameter by

$$s = \frac{D}{E} \frac{\hat{s}_{ij}^s}{nn} \quad (30)$$

for nn sites, i and j . This RVB order parameter is different from the RVB order parameter introduced by Anderson and collaborators^{1,2}: it involves the charge degrees of freedom through the gauge field, A_{ij}^h , and is gauge invariant under (16). Note that $b_i^y b_j^y \rightarrow 0$ in this theory, and s is the sole order parameter. At half filling, $A_{ij}^h = 0$ in \hat{s}_{ij}^s , and s reduces to the well known Schwinger-boson mean field order parameter $\langle b_i b_j \rangle$ ¹³ [in our definition b_i differs from the conventional Schwinger boson representation by a sign factor $(-1)^i$]. The mean field solution describes both long and short range AF correlations of the Heisenberg model quite well, as will be discussed below. Thus, the bosonic RVB theory we present is ensured to yield good results at half filling, and provides a natural generalization away from half filling.

We now show how the spin degrees of freedom are treated within the bosonic RVB theory. The superexchange Hamiltonian is rewritten as

$$H_s = \sum_{\langle ij \rangle} J_s \frac{X}{E} e^{i A_{ij}^h} b_i^y b_j^y + h.c. : \sum_i \left(b_i^y b_i + \frac{1}{2} \right); \quad (31)$$

where $J_s = J^s/2$, and s is assumed to be real and positive. The Lagrangian multiplier is introduced to implement the single occupancy constraint (4) on the average. We reemphasize that H_s is not a mean field Hamiltonian in the conventional sense. Spinons are coupled to a gauge field A_{ij}^h , related to the charge degrees of freedom, and the Hamiltonian is invariant under the internal gauge transformation (16). So H_s is still a gauge model, describing low energy, long wavelength spin fluctuations, underpinned by the bosonic RVB order parameter (30). The link field A_{ij}^h defined in (14) describes holons as vortices perceived by spinons in (31). In the ground state, A_{ij}^h will be treated as a constant field, as though the holons are Bose condensed (which condition will be determined self consistently, when we discuss the charge degrees of freedom), with

$$A_{ij}^h = \frac{X}{m} \quad (32)$$

for each plaquette, where $\frac{X}{m}$ is the hole concentration.

Then, H_s can be diagonalized by a Bogoliubov transformation¹⁷

$$b_i = \sum_m w_m(i) u_m + \sum_m v_m \frac{y}{m}; \quad (33)$$

as

$$H_s = \sum_m E_m \frac{y}{m} + \text{const}; \quad (34)$$

Here, $u_m = \frac{1}{\sqrt{2}} \sqrt{\frac{1}{E_m} + 1}$; $v_m = \frac{1}{\sqrt{2}} \sqrt{\frac{1}{E_m} - 1} \text{sgn}(m)$; and $E_m = \frac{P}{2} \frac{1}{m^2}$. The Lagrangian multiplier is determined by enforcing $\sum_i b_i^y b_i = (1 - \frac{X}{m})N$. The wave function w_m and the spectrum E_m are determined

by the following eigen equation,

$$\sum_m w_m(i) = \sum_{j=nn(i)}^X e^{i A_{ij}^h} w_m(j) : \quad (35)$$

The ground state of H_s is constructed in the usual way by imposing the condition, $\langle \text{RVB} |_{MF} = 0$,

$$\langle \text{RVB} |_{MF} = \exp \left[\sum_{ij}^X W_{ij} b_{i\alpha}^\dagger b_{j\beta}^\dagger A_{ij} \right] |i\rangle ; \quad (36)$$

where the RVB amplitude W_{ij} is given by

$$\begin{aligned} W_{ij} &= \sum_m \frac{v_m}{u_m} w_{m\alpha}(i) w_{m\beta}(j) \\ &= \sum_m \frac{v_m}{u_m} w_{m\beta}(i) w_{m\alpha}(j) : \end{aligned} \quad (37)$$

In deriving the above, we use $w_m(i) = w_m(i)$, which follows from (35). Furthermore, it is easy to show that for each state labeled by m , there is always a state labeled by m , for which $m = m$ and $w_m(i) = (-1)^i w_m(i)$, such that

$$\begin{aligned} W_{ij} &= \frac{1}{2} (-1)^j W_{ij} \\ &\neq 0 \quad \text{only if } i, j \text{ different sublattices,} \end{aligned} \quad (38)$$

i.e., the RVB amplitude W_{ij} only connects α and β spins on opposite sublattices.

The nn spin correlations are determined by

$$\langle S_i \cdot S_{nn} \rangle = \frac{3}{8} \langle S_j^2 \rangle < 0 \quad (39)$$

which is antiferromagnetic in nature [i.e., in the resulting bosonic RVB theory, $\langle S_i \cdot S_j \rangle > 0$ (< 0), if i and j belong to the same (opposite) sublattice]. Therefore, $\langle S_j^2 \rangle \neq 0$ generally characterizes a regime with short range AF correlations, present in a wide range of temperature in the t - J model at low doping (e.g., $T = 1500$ K in the zero-doping limit). Fig. 2 sketches a phase diagram in which $\langle S_j^2 \rangle$ or short-range AF correlations controls the high-energy, short-distance correlations in a low-doping regime ($< x_{RVB}$)¹⁷. Such a spin singlet-pairing regime, characterized by $\langle S_j^2 \rangle \neq 0$, defines the "pseudogap" regime of the bosonic RVB state, with a characteristic temperature T_0 , as illustrated in Fig. 2.

In the following two subsections, we further examine the spin correlations at half-filling and finite doping within this bosonic RVB description.

1. Antiferromagnetism at half-filling

At half-filling, the bosonic RVB ground state $\langle \text{RVB} | = P_G \langle \text{RVB} |_{MF}$, which satisfies the no-double-occupancy constraint (P_G denotes the Gutzwiller projection operator), can be explicitly written as

$$\langle \text{RVB} | = \sum_{fsg \text{ pair}(ij)}^X \exp \left[\sum_{ij}^X W_{ij} A_{ij} b_{i1}^\dagger h_1 b_{i2}^\dagger h_2 \dots b_{iN}^\dagger h_N \right] |i\rangle ; \quad (40)$$

where each site is occupied only by one spinon, with $fsg = 1, 2, \dots, N$ being partitioned into RVB pairs connected by the RVB amplitudes, W_{ij} . Here each RVB pair is denoted by (ij) , where $i(j)$ refers to the position of an α (β) spin, as before. The summation runs over all possible RVB pairing partitions. The holon annihilation operators in (40) are introduced since the spinon vacuum $|i\rangle$ is defined as a state where every site is occupied by a holon, for later convenience.

$\langle \text{RVB} |$ given in (40) is equivalent to the class of variational wave functions (cf Eq.(1)) proposed by Liang, Doucot, and Anderson (LDA)⁸. The RVB amplitudes can be determined from the bosonic mean field theory outlined above. At half-filling, $A_{ij}^h = 0$, and the mean field Hamiltonian (31) reduces to the Schwinger-boson mean field Hamiltonian.




FIG. 2: Schematic phase diagram of the b-RVB mean field theory: The bosonic RVB order parameter Δ^s underpins a pseudogap phase at $T < T_0$, in which the AF state, spontaneous vortex phase (a low- T pseudogap state), as well as the superconducting instability occur at low temperatures, separated by some critical dopings, x_c and x_{RVB} , respectively.

In this case, the RVB amplitude $W_{ij} / |j_{ij}|^3$, for $|j_{ij}| = a$ (a being the lattice constant), as shown in the inset of Fig. 3. We use the loop-gas method employed by LDA to determine the nn spin correlation (which also decides the ground state energy of the Heisenberg model), as well as the staggered magnetization, m . The maximum sample size is 64×64 . We obtained $\langle S_i \cdot S_{i+n} \rangle = 0.3344(2)J$, and a staggered magnetization, $m = 0.296(2)$. These results are essentially the same as the best variational result obtained by LDA, and also compare extremely well with exact numerical results⁸, $\langle S_i \cdot S_{i+n} \rangle = 0.3346$, and $m = 0.31$.

Therefore, at half filling, the insulating state f-RVB correctly reproduces both short range and long range AF correlations of the Heisenberg model. AFLRO appears naturally in the long wavelength limit, with a correct magnitude of magnetization; short range spin correlations are also well described as evidenced by the excellent agreement of our results with the ground state energy obtained from numerical studies.

2. Spin liquid at finite doping

At finite doping, the RVB amplitudes W_{ij} , determined by (31) and (37) decay exponentially for large spatial separations,

$$|W_{ij}| \sim e^{-\frac{|r_{ij}|^2}{2\xi^2}} \quad (41)$$

with a characteristic length scale

$$\xi = \frac{r}{a} \quad (42)$$

as shown in Fig. 3. Correspondingly, AFLRO disappears and a finite length scale, ξ , for equal time spin correlations emerges; viz., a featureless RVB spin liquid with short range correlations evolves continuously from an AF ordered state at half filling ($\delta = 1$).

Spin excitations can be created by the action of a spin raising/lowering operator on the spin liquid f-RVB in (40). In the bosonic mean field theory, spin excitations are described by the spinon RVB Hamiltonian, (31). The results from mean field theory have been reported elsewhere^{17,18}, and we briefly summarize the results below. At small doping, the spin excitations evolve (from the gapless magnon mode at half filling) into a sharp resonance at low energies,




FIG. 3: The RVB amplitudes, $|w_{ij}|$, obtained from the b-RVB theory for three different hole concentrations. The inset shows the result for half filling.

with a characteristic energy scale, E_g / J , around the momentum $(\pi; \pi)$: The spectral weight of this excitation is proportional to r , while its width in the momentum space is proportional to r^{-1} , which follows from (41). The evolution of the gapless magnon mode at half filling into a resonant mode at finite energies reflects the phase string effect on the spin degrees of freedom, through the effect of the gauge field, A_{ij}^h .

B. Charge degrees of freedom : optimizing the kinetic energy of doped holes

At finite doping, the mean field state $|\Psi_{MF}\rangle$ describes a condensate of RVB spinon pairs, and does not involve the charge degrees of freedom (holons) explicitly; i.e., the distribution of holons is independent of the positions of spinons. Since the single occupancy constraint is no longer crucial in the bosonic RVB theory, it is tempting to posit a mean field ground state

$$|\Psi\rangle = |\Psi_h\rangle |\Psi_{MF}\rangle \quad (43)$$

where $|\Psi_h\rangle$ describes the holon condensate. This approach, adopted in a previous work¹⁷, may be justified for a long wavelength, low energy theory as the local constraint is not crucial in the phase-string formalism¹⁹. However, to obtain the ground state wavefunction, it is desirable to employ a construction where the single occupancy constraint is satisfied exactly; viz., a method that treats spin backflow effects more carefully, which is also important for facilitating and optimizing the kinetic energy of doped holes. In the following, we show how this can be done differently.

First, we project out all double occupancies in $|\Psi_{MF}\rangle$, and define $|\Psi_i\rangle$ as given in (40), where each site is occupied only by one spinon, as if the system were at half filling. However, the RVB amplitudes, w_{ij} , will be determined by the bosonic RVB mean field theory of H_S at a fixed hole concentration.

Next, for every doped holon at site l , a spinon at the same site l is annihilated, by the action of $h_l^\dagger b_l$ on $|\Psi_i\rangle$, i.e.,

$$h_l^\dagger b_l |\Psi_i\rangle \quad (44)$$

such that the constraint is automatically satisfied. As discussed below, the construction (44) optimizes the kinetic energy of the holon-spinon composite, $h_l^\dagger b_l$, by maximizing the overlap between the RVB configurations before and after hopping. Clearly, the behavior of the spinon b_l characterizes the spinon backflow accompanying the motion of the holon in the RVB background.




FIG. 4: Ratio of the effective hopping integral $\frac{t_h}{t_c}$ obtained from (49).

Thus, the full ground state may be constructed as

$$|j_G\rangle = \sum_{f_{l_h}g}^X \langle f_{l_h}g | \langle l_1; l_2; \dots; l_{N_h} \rangle \sum_h^Y Z_h(l_h) h_{l_h}^y b_{l_h h} |RVBi\rangle; \quad (45)$$

where $\langle f_{l_h}g |$ is a bosonic (symmetric) many body wave function for holons, and $Z_h(l_h)$ denotes a single particle wave function for the spinon back flow $b_{l_h h}$, with the spin index h under the constraint $l=2$ $l_h h = S_z$ (note that $S_z = 0$ in $|RVBi\rangle$).

In the following, we describe how to determine $\langle f_{l_h}g |$ and $Z_h(l_h)$ in (45) based on a mean-field approximation. For this purpose, we introduce a mean-field ground state as follows

$$|j_G\rangle = P_G |j_G i_{MF}\rangle \quad (46)$$

where the mean-field state

$$|j_G i_{MF}\rangle = \sum_{f_{l_h}g}^X \langle f_{l_h}g | \langle l_1; l_2; \dots; l_{N_h} \rangle \sum_h^Y Z_h(l_h) h_{l_h}^y b_{l_h h} |RVBi_{MF}\rangle \quad (47)$$

By optimizing the kinetic energy of the holon-spinon composite, $h_l^y b_l$, based on (47), $\langle f_{l_h}g |$ and $Z_h(l_h)$ are determined as follows. As shown in Appendix A, one finds

$$\langle f_{l_h}g | \langle l_1; l_2; \dots; l_{N_h} \rangle = \text{const}; \quad (48)$$

namely, the holons are in a Bose condensed state; $Z_h(l_h)$ satisfies the following eigen equation

$$(-t_h)Z_h(i) = \frac{t}{4} \sum_{j=nn(i)}^X e^{i\phi_{ij}} e^{i\phi_{ij}^h} Z_h(j); \quad (49)$$

with $t_h > 0$ being the maximum eigenvalue such that the total hopping energy is given by

$$E_{tj} = E_{tj} = -4t_h N; \quad (50)$$

up with the RVB amplitudes, $(1/W_{ij})$. The latter depend upon doping as determined by the mean-field theory described in the previous section (see Fig. 3). The summation in Ψ_{RVB} runs over all possible pairing partitions for each configuration f_{sg} .

The ground state $|j_G\rangle$, reduces to $|\text{RVB}\rangle$ at half-filling, where it describes the antiferromagnetism extremely well as discussed in Sec. III A 1. Upon doping, $|\text{RVB}\rangle$ may be regarded as an insulating RVB vacuum. From (54), we see that each doped hole is "twisted" into a bosonic object e , via the nonlocal phase operator $\hat{\phi}_i$, in order to optimize its kinetic energy in such an RVB vacuum. The form of the state $|j_G\rangle$ suggests a generalized "spin-charge separation"; viz., the bosonic holes created by e , move freely in the background $|\text{RVB}\rangle$; they carry the charge while the vacuum is described by the pairing of neutral spins (spinons). These two degrees of freedom are indeed decoupled in the mean-field sense. But this is different from the typical spin-charge separation because the twisted holes are accompanied by spins. Indeed, annihilating a spinon at the site l in the RVB vacuum by c_l will always create an unpaired spinon within a scale where the spins were originally RVB paired (with no free spinons). So the state $|j_G\rangle$ is composed both of RVB paired spinons and twisted holes as the holon-spinon pairs as in (57) below.

Since the spins at the hole sites l 's in the vacuum $|\text{RVB}\rangle$ are automatically annihilated by c_l 's, the no double occupancy constraint is exactly imposed in $|j_G\rangle$ at finite doping. As shown in Appendix C 1, the ground state retains time reversal symmetry, which may not be obvious from the form of $|j_G\rangle$. It also respects global spin rotational symmetry under the approximation that the twisted holes are perfectly Bose condensed, as discussed in Appendix C 2.

The structure of $|j_G\rangle$ in the doped case looks rather simple in terms of the twisted holes moving on the RVB vacuum. But it would look more complicated when written in terms of the c -operators in (54) because $\hat{\phi}_i$ is highly nontrivial. Physically, the phase string $\hat{\phi}_i$ can be regarded as the nonlocal phase shift induced by doped holes. In the one dimensional case, where it is responsible for the Luttinger liquid behavior, the phase string can be identified precisely with the phase shift obtained from the Bethe ansatz solution for the large- U Hubbard model¹¹. In the two dimensional case, the role of the phase string operator $\hat{\phi}_i$ may be seen more clearly in the wavefunction form discussed in Sec. II, i.e., (24). Below we rewrite the wavefunction (54) in terms of the underlying electrons.

Starting from the wavefunction Ψ , corresponding to the ground state (45) in the basis of (21), and using the relation between the wavefunctions in the two basis sets (24), we get,

$$\Psi(f_{ij}g; f_{jl}g) = K \sum_{\text{pair}} \prod_{(ij)}^2 (1/W_{ij}) \prod_{(l_1 j_1)}^Y X_{l_1 j_1} \prod_{(i_1 l_1^0)}^Y X_{i_1 l_1^0} \prod_{(ll^0)}^Y Y_{ll^0}^5 : \quad (57)$$

Here, we have invoked the holon condensation condition (48). In (54), a given spin-hole configuration, $f_{ij}g$, $f_{jl}g$ and $f_{lh}g$, is partitioned into pairs, denoted by (ij) (the sites, i and j , are occupied by an up and a down spin, respectively), $(l_1 j_1)$ (sites l_1 and j_1 are occupied by a hole and a down spin, respectively), $(i_1 l_1^0)$ (two sites, i_1 and l_1^0 , are occupied by an up spin and a hole, respectively), and (ll^0) (a pair of holes). The latter three pairings have amplitudes, $X_{l_1 j_1} = Z^*(l_1)W_{l_1 j_1}$, $X_{i_1 l_1^0} = (1/W_{i_1 l_1^0})Z^*(l_1)$; and $Y_{ll^0} = Z^*(l)Z^*(l^0)W_{ll^0}$, respectively. Fig. 5 shows two possible partitions for a given spin-hole configuration, and the summation in (57) runs over all possible partitions. Thus, the wavefunction (57) shows that the phase string operator $\hat{\phi}_i$ leads to the generalized Jastrow-type factor K defined in (25). The remaining part of the wavefunction is bosonic, describing RVB pairing of spins, pairing between spins and holes as well as between holes.

B. Hidden ODLRO in the ground state: A pseudogap state

The properties of the RVB vacuum $|\text{RVB}\rangle$ were discussed in Sec. III A. In the following we shall focus on the charge degrees of freedom in $|j_G\rangle$.

>From the definition of (54), it follows that

$$|j_G\rangle = \sum_{\text{pair}} \prod_{(ij)}^2 (1/W_{ij}) \prod_{(l_1 j_1)}^Y X_{l_1 j_1} \prod_{(i_1 l_1^0)}^Y X_{i_1 l_1^0} \prod_{(ll^0)}^Y Y_{ll^0}^5 : \quad (58)$$

for finite doping: $N_h^{\text{net}} = 0$ ($N=2$). Thus the ground state exhibits an ODLRO as the twisted holes, described by e [or more precisely the bosonic object, $(1/Z^*(l))c_l$, which is gauge-invariant under (15) and (16)], are Bose condensed on the RVB vacuum $|\text{RVB}\rangle$ in (54). Since c_l is a nonlocal operator which differs from the electron operator by an infinite-body phase-string twist, the corresponding ODLRO in (58) is unlike the conventional ODLRO based on a local combination of electron operators, such as spin magnetization or the BCS superconducting pairing operator. In other




FIG. 5: Pairing among spins (full circles) and holes (open circles) are illustrated by the connection of lines. Two sets of bonds (solid and dashed lines) represent two different resonating pairing patterns, forming a loop covering of the lattice for a given spin-hole configuration.

words, it does not have a direct correspondence to a physical observable. Such a hidden ODLRO characterizes how doped charges (holes) are collectively reorganized and behave in the RVB background, RVBi. As the consequences, the ground state and excited states will exhibit some very unique novel properties. This will be the subject of our discussion below.

1. Pairing amplitude

We first show that the hidden ODLRO in j_Gi will result in a finite pairing amplitude between electrons in the singlet channel.

The electron singlet pair operator can be expressed as

$$\begin{aligned} \hat{c}_{ij}^{SC} &= \sum_i c_i c_j \\ &= \sum_i e^{i\hat{\phi}_i} c_i e^{i\hat{\phi}_j} c_j \\ &= \hat{c}_{ij}^0 e^{i\frac{1}{2}(\phi_i + \phi_j)} : \end{aligned} \quad (59)$$

In the last line, the pair operator is formally written in terms of an amplitude and a phase part, with the amplitude operator defined by²⁰

$$\hat{c}_{ij}^0 = \sum_i^h (1)^j e^{i\phi_j^0} e^{i\phi_{ij}^0} e^{iA_{ij}^h} c_i c_j \quad (60)$$

(a constant factor $(-1)^h$ is omitted).

We now demonstrate that $\hat{c}_{ij}^0 \neq 0$, i.e., the pairing amplitude is finite as the consequence of the hidden ODLRO. Since the twisted holes are Bose condensed, $(c_i c_j)^Y j_G \frac{1}{N_e - 2} = A j_G i_{N_e} + \text{excited states}$, where the coefficient A is nonzero and can be estimated by $A = Z(i)Z(j) \frac{N_h'' N_h^\#}{N_h}$ at small doping²³. The pair amplitude $\hat{c}_{ij}^0 = N_e - 2 j_G j_G \hat{c}_{ij}^0 j_G i_{N_e}$ is then given by

$$\begin{aligned} \hat{c}_{ij}^0 &= \frac{1}{N_h'' N_h^\#} \sum_i^h (1)^j e^{i\phi_j^0} e^{i\phi_{ij}^0} e^{iA_{ij}^h} Z(i)Z(j) \\ &= \frac{N_h''}{2} e^{i\phi_{ij}^0} e^{iA_{ij}^h} Z(i)Z(j); \end{aligned} \quad (61)$$

where we have used $N_h'' = N_h^\# = N_h = 2 = N$, and

$$Z(i)Z(j) = Z(j)(-1)^j e^{i\phi_j^0} : \quad (62)$$

The above equation can be obtained from (49), and by noting that $e^{i \frac{0}{i} i \frac{0}{j}} = e^{i 2 \frac{0}{ij}}$. In order to estimate the size of the pair amplitude, we focus on the pairing amplitude between nn sites, $(\frac{0}{ij})_{nn}$. Invoking translational invariance of the ground state, we get

$$\begin{aligned}
 & \frac{1}{2N} \sum_{\langle ij \rangle} \langle \frac{0}{ij} \rangle_{nn} \\
 &= \frac{1}{4} \sum_{\langle ij \rangle} \langle \frac{0}{ij} \rangle e^{i \frac{0}{ij} i A_{ij}^h Z(i) Z(j)} \\
 &= \frac{1}{4} \sum_i \langle \frac{0}{i} \rangle \frac{2t_h}{\tau} Z(i)^2 \\
 &= \frac{t_h}{\tau} : \tag{63}
 \end{aligned}$$

Therefore the ground state (54) indeed possesses a finite pairing amplitude as a direct consequence of the hidden ODLRO in j_{Gi} , which is proportional to the doping concentration for small doping. Note that a finite pairing amplitude exists only in the singlet channel. It is easily shown that the mean value of the pairing amplitude in the triplet channel vanishes identically [in this case, an additional sign appears inside the summation of (60)].

However, superconducting phase coherence is generally absent. The phase part of the pair operator (59) involves $\frac{s}{i}$ which is defined in (6) and is closely related to the spin degrees of freedom. More precisely, each unpaired spin contributes to a free $\frac{1}{2}$ vortex via $\frac{s}{i}$ in the pair operator (59). As will be discussed in subsection C, the presence of free spins in the ground state j_{Gi} prevents phase coherence in (59). Nevertheless, $\hat{\frac{0}{ij}} \neq 0$ in (59) is a meaningful definition and description of a low temperature pseudogap ground state, called the spontaneous vortex phase²¹. In this phase, each unpaired neutral spin (spinon) always carries a $\frac{1}{2}$ supercurrent vortex, known as a spinon-vortex²². We postpone further discussion of the spinon-vortices to Sec. IV C.

2. Fermionic nodal quasiparticles

In the following, we construct a quasiparticle excitation based on the ground state j_{Gi} . In particular, we show that the existence of nodes in the excitation spectrum is ensured by the hidden ODLRO.

Consider the action of c_i^y on the ground state j_{Gi} . Since $c_i = c_i e^{i \frac{0}{i}}$ and the twisted bosonic holes c_i are condensed, one finds

$$\begin{aligned}
 c_i^y j_{Gi} |_{N_e-1} &= e^{i \frac{0}{i}} c_i^y j_{Gi} |_{N_e-1} \\
 &= e^{i \frac{1}{2} \frac{s}{i} B_i} j_{Gi} |_{N_e} + \text{excited states}; \tag{64}
 \end{aligned}$$

where

$$\begin{aligned}
 B_i &= \frac{p}{N_h + 1} Z(i) e^{i \frac{0}{i}} e^{i \frac{1}{2} \frac{s}{i}} \\
 &= \frac{N}{2} (1)^{N_h} Z(i) e^{i \frac{1}{2} \frac{s}{i}} e^{i \frac{0}{i}} e^{i \frac{0}{i} \frac{h}{i}}; \tag{65}
 \end{aligned}$$

Let us focus on B_i first. Rewriting B_i using (62), we get

$$B_i = \frac{N}{2} Z_{\#}(i) e^{i \frac{1}{2} \left(\frac{0}{i} \frac{h}{i} \right)}; \tag{66}$$

and

$$B_{i\#} = \frac{N}{2} (1)^{N_h} Z_{\#}(i) e^{i \frac{1}{2} \left(\frac{0}{i} \frac{h}{i} \right)}; \tag{67}$$

Nontrivial spatial oscillations in B_i mainly originate from the phase factor $e^{i \frac{1}{2} \left(\frac{0}{i} \frac{h}{i} \right)}$. We rewrite this phase factor as a string operator,

$$e^{i \frac{1}{2} \left[\frac{0}{i} \frac{h}{i} \right]} = e^{i \frac{1}{2} \left[\frac{0}{i} \frac{h}{i} \right]} e^{i \frac{1}{2} \left[\frac{0}{i_1} \frac{h}{i_1} \right]} e^{i \frac{1}{2} \left[\frac{0}{i_1} \frac{h}{i_1} \right]} e^{i \frac{1}{2} \left[\frac{0}{i_2} \frac{h}{i_2} \right]}; \tag{68}$$

where $i; i_1; i_2; \dots$ is a sequence of nn sites on a path ending at site i . Using the relation $\langle c_{i_1}(i) = \dots \rangle$, and invoking the Bose condensation of the holes, we get

$$e^{-i\frac{1}{2}[\frac{0}{i} \frac{h}{i}]} e^{-i\frac{1}{2}[\frac{0}{i_1} \frac{h}{i_1}]} \dots e^{-i\frac{1}{2}(1)} e^{-i[\frac{0}{ii_1} A_{ii_1}^h]}; \text{ etc.}; \quad (69)$$

with \dots such that

$$B_i B_j^\dagger = D_{ij} e^{-ik_0 \cdot (r_i - r_j)} \quad (70)$$

where

$$k_0 = \frac{1}{2}[\dots]; \frac{1}{2}[\dots] \quad (71)$$

at four points in the Brillouin zone (setting the lattice constant $a = 1$). The quasiparticle weight is proportional to

$$\begin{aligned} D_{ij}'' &= D_{ij}^\# \\ &= \frac{N}{2} Z_\#(i) e^{-i\frac{1}{2}[\frac{0}{im} \frac{h}{im}]} Z_\#(j) \\ &\quad O(\dots); \end{aligned} \quad (72)$$

which is gauge invariant, as can be seen from (49). Here, i_j denotes a shortest-path connecting sites i and j . Therefore one finds

$$c_{i_1}^\dagger j_{G i_{N_e-1}}' e^{-ik_0 \cdot r_i} e^{-i\frac{1}{2}[\frac{0}{i} \frac{h}{i}]} j_{G i_{N_e}} : \quad (73)$$

If phase coherence is realized such that

$$e^{-i\frac{1}{2}[\frac{0}{i} \frac{h}{i}]} j_{G i} = j_{G i} \quad (74)$$

(see Sec. IV C), then $c_{i_1}^\dagger j_{G i_{N_e-1}}$ would have a finite overlap with $j_{G i_{N_e}}$; namely, the quasiparticle spectrum has four nodes at momenta, k_0 's, that approach $\frac{1}{2}; \frac{1}{2}$ in the limit of zero doping. Similar momentum structure can be identified in $c_k j_{G i_{N_e+1}}$. A true d-wave-like quasiparticle state in the superconducting state can be constructed as a linear combination of $c_k^\dagger j_{G i}$ and $c_{-k} j_{G i}$, which should be infinitesimally close to the ground state $j_{G i_{N_e}}$ at the four nodal points, k_0 's.

However, as discussed in Sec. IV C, the phase factor $e^{-i\frac{1}{2}[\frac{0}{i} \frac{h}{i}]}$ precludes phase coherence, and therefore the quasiparticle state $c_{i_1}^\dagger j_{G i_{N_e-1}}$ actually has a vanishing overlap with the ground state $j_{G i_{N_e}}$ because of such a phase factor. Thus, the low-lying fermionic quasiparticles near the four nodes, k_0 's, have a vanishing spectral weight $Z_k j_{G i} h_G j_{G i_{N_e-1}}^\dagger = 0$; viz., the non-superconducting (pseudogap) ground state $j_{G i}$ is a non-Fermi-liquid in nature. Long lived quasiparticle excitations are expected to emerge only when superconducting phase coherence is realized in (74).

We emphasize that the nodal structure of quasiparticles results from the ODLRO of the bosonic twisted holes, in which the phase string factor $e^{-i\hat{\phi}_i}$, reflecting the fermionic nature of the electron operator, determines the momentum positions of four nodes. Finally we note that in the previous effective phase string theory in terms of holons and spinons (see Appendix B), a similar d-wave quasiparticle excitation can be obtained as the consequence of forming the holon-spinon bound pairs by considering residual interactions beyond the effective model²⁰.

3. Two types of spin excitations

The hidden ODLRO in (54) is not only responsible for the existence of a finite pairing amplitude as well as d-wave like nodal quasiparticle excitations, it also predicts the presence of a new type of low-lying spin excitations which are quite distinctive as compared to the previously discussed spin excitations created in the RVB background $\mathcal{R}VB i$ (Sec. III A):

The latter may be called Type I spin excitations, which can be created by acting a spin operator, say, S_i^+ , directly onto the insulating spin liquid $\mathcal{R}VB i$, without involving the hole condensate in (54). Obviously, this type of spin excitations is the only elementary excitation at half-filling, where it reduces to a conventional spin wave in the long-wavelength limit as discussed in Sec. III A. At small doping, the majority of spins still remain RVB paired, characterized by $s \ll 0$ in $\mathcal{R}VB i$; Thus, this kind of spin excitation will naturally persist into the pseudogap regime,

where a spin gap will be opened up in its low-energy spectrum as the spin-spin correlations are short-ranged (here W_{ij} become exponential decay with the distance $|j-i|$; see Fig. 3). In fact, as mentioned in Sec. III A, the corresponding low-lying excitation reduces to a sharp resonance-like mode in the pseudogap regime, with a characteristic energy scale $E_g \sim J$ around the momentum (π, π) . Resonant spin wave like dispersion is still present at high energies, which disappears when the temperature reaches T_0 as shown in Fig. 2 where the short-range RVB pairing vanishes naturally.

Now let us discuss a new type (Type II) of $S = 1$ excitation which is associated with the twisted holes and may be regarded as composed of a pair of quasiparticle and quasihole, instead of a pair of neutral spinons in RVBi. To create a Type II spin excitation by, say, S_i^+ ; a pair of quasiparticles will be created and annihilated in $j \in i$, which involves a spin flip in the "hole condensate" part in (54).

To show this, let us rewrite $S_i^+ = c_{1i}^\dagger c_{i\#}$ by

$$\begin{aligned} S_i^+ &= e^{i\hat{\phi}_i} c_{1i}^\dagger c_{i\#} e^{-i\hat{\phi}_{i\#}} \\ &= (-1)^{i-h} e^{i\hat{\phi}_i} c_{1i}^\dagger c_{i\#} : \end{aligned} \quad (75)$$

Then, by using $(\text{hole})_{\text{part}}$ to denote Type II spin excitations created by S_i^+ in the condensed "hole" part in $j \in i$, we obtain

$$S_i^+ |j \in i\rangle_{\text{hole part}} = (-1)^{i-h} Z_{\#}(i) Z_{\#}(i) N_h^{\#} N_h^{\#} + 1 |j \in i\rangle : \quad (76)$$

The term on the right-hand side (rhs) has the same origin as the quasiparticles around nodal points discussed previously. But the phase factor $e^{i\hat{\phi}_i}$ in the quasiparticle channel [(64)] is now replaced by $(-1)^{i-h}$ in the spin channel, after a recombination of $e^{i\hat{\phi}_i}$ and $e^{-i\hat{\phi}_{i\#}}$ in (75)²⁴: The leading contribution to the spin-spin correlation at large distance is then given by

$$\langle S_i^+ |j \in i\rangle_{\text{hole part}} \langle S_j^+ |j \in j\rangle_{\text{hole part}} = (-1)^{i-j} Z_{\#}(j) Z_{\#}(j) N_{ij}^{\#} Z_{\#}(i) Z_{\#}(i) N \quad (77)$$

(using $N_h = \frac{P}{N}$) where

$$N_{ij}^{\#} = e^{i(\hat{\phi}_i - \hat{\phi}_j)} : \quad (78)$$

The momentum structure can be determined by the Fourier transformation of (77), in a way similar to that in the quasiparticle channel. Note that, an additional topological effect in $e^{i\hat{\phi}_i}$ (not in $e^{i\frac{1}{2}\hat{\phi}_i}$) requires a little more careful treatment. As detailed elsewhere²⁵, $(-1)^{i-h}$ will lead to the following incommensurate momenta, to leading order of approximation,

$$Q = [\quad ; \quad (1 \quad)] \quad \text{and} \quad [\quad (1 \quad) ; \quad] \quad (79)$$

with $\cdot = 2\pi$.

On the other hand, the dynamics of (77) is mainly decided by the vortices in $e^{i\hat{\phi}_i}$ which is in turn determined by the charge density. To leading order approximation, no superexchange interaction in the spin degrees of freedom influences (77). If one neglects the fluctuation in $e^{i\hat{\phi}_i}$; the term on the rhs of (76) would have a finite overlap with $|j \in i\rangle$ at the incommensurate momenta Q , which implies a zero mode or the presence of an incommensurate AFLRO. But the true long-range order will be prevented by the density fluctuations, because any small fluctuation in the charge density will be accumulated in the phase factor $e^{i\hat{\phi}_i}$ in a sufficiently long time and distance; which in turn will result in a slow-fluctuating incommensurate AF ordering above $|j \in i\rangle$, instead of a true static incommensurate AFLRO. This low-lying spin fluctuation can provide an alternative explanation, without invoking a dynamic stripe picture, of low-energy results in neutron scattering experiments on the cuprate superconductors.

Therefore, Type II spin excitations in (76) are quite distinct from Type I in nature. As a direct consequence of the hidden ODLRO, they are absent at half-filling [the weight of this term is proportional to the density of the charge condensate, from (76)]. One should thus properly distinguish these two in a neutron scattering measurement, as a unique prediction of this theory.

C. Superconducting phase coherence

As described above, the hidden ODLRO in $|j \in i\rangle$ leads to a nonzero pairing amplitude. But it does not necessarily imply superconductivity. Consider the action of the superconducting pairing operator on the N_e electron ground

state. The resulting state has an overlap with $N_e - 2$ electron ground state, determined by using (59) and (63), as follows:

$$\hat{\psi}_{ij}^{SC} |j_G\rangle_{N_e} = \frac{1}{\sqrt{2}} e^{i\frac{1}{2}(\phi_i + \phi_j)} |j_G\rangle_{N_e-2} + \text{excited states:} \quad (80)$$

So superconductivity is established only when

$$\frac{D}{E} e^{i\frac{1}{2}(\phi_i + \phi_j)} \neq 0; \quad (81)$$

viz., the onset of superconductivity is determined by phase coherence.

1. Spinon vortices

We noted earlier that each isolated ($S = 1/2$) spin in $|j_G\rangle$ means the creation of a ± 2 vortex in the phase of the superconducting order parameter $\hat{\psi}_{ij}^{SC}$, via ϕ_i , according to (80). Thus, each spin automatically carries a phase or supercurrent vortex, called a spinon-vortex, in the state described by $|j_G\rangle$. This was first identified in previous work based on the effective theory^{21,22}.

The ± 2 vortices centered at α and β spins in $|RVB\rangle$ are paired as vortex-antivortex pairs, since the underlying spins are always paired in the singlet RVB vacuum. Then, one has

$$\langle RVB | e^{i\frac{1}{2}(\phi_\alpha + \phi_\beta)} | RVB \rangle \neq 0. \quad (82)$$

This is in analogy with the two-dimensional XY model below the Kosterlitz-Thouless transition where vortex-antivortex pairs are bound together. In our case, vortices and antivortices are associated with α and β spins, respectively. The condition that $S^z = 0$ in the singlet background, $|RVB\rangle$, is equivalent to the charge neutrality condition in the XY model. Since α and β spins are RVB pairs with a finite range at finite doping [see (42)], the cancellation in $\frac{1}{2}(\phi_\alpha + \phi_\beta)$ is ensured at large distances. Thus, we see that the spin RVB pairing in $|RVB\rangle$ is very essential for phase coherence to occur. Note that this condition fails in the AF phase, where $\phi \neq 0$, or beyond the doping concentration x_{RVB} where the RVB pairing disappears.

However, the existence of free spins, associated with doped holes created by c_l , will imply the presence of free vortices in $|j_G\rangle$. To see this, following (54) and (82), one can write

$$e^{i\frac{1}{2}(\phi_\alpha + \phi_\beta)} |j_G\rangle = e^{i\frac{1}{2}(\phi_\alpha + \phi_\beta)} \prod_l Z_\alpha(l) c_l^{\dagger} \prod_l Z_\beta(l) c_l \frac{1}{\sqrt{2}} e^{i\frac{1}{2}(\phi_\alpha + \phi_\beta)} |RVB\rangle \\ e^{i\frac{1}{2}(\phi_\alpha + \phi_\beta)} |j_G\rangle \quad (83)$$

where $\phi_l = \sum_{i \in \text{path}} \phi_i$ with ϕ_i denoting the number operator of twisted holes of spin α at site l . Then, the Bose condensation of these free vortices in $|j_G\rangle$ will disorder the phase of the superconducting pairing order parameter according to (83) (although ϕ_l itself vanishes on average, its fluctuations will satisfy an area law in the Bose condensed case).

Therefore the ground state $|j_G\rangle$ is not a true superconducting ground state due to the lack of phase coherence, although it possesses a pairing amplitude as well as (incoherent) nodal quasiparticles. It is characterized by the presence of unpaired spinon-vortices as a direct consequence of the hidden ODLRO in $|j_G\rangle$. Nontrivial Nernst effect contributed by such unpaired spinon-vortices has been investigated based on the effective phase string theory elsewhere²¹.

2. Superconducting ground state

The superconducting instability will take place when those unpaired spins associated with twisted holes also become RVB paired. It reduces the superexchange energy when those backflow spins form the RVB pairs, and most importantly it removes the logarithmic-divergent energy associated with free vortices as vortices-antivortices are paired up at low temperature. Consequently the phase ϕ_l is cancelled out in (83) and the true phase coherence (81) of the superconducting order parameter can be finally established.

Corresponding to the pairing of those free spins associated with twisted holes, the twisted holes themselves are also effectively paired and the ground state should evolve from (54) into the following form

$$j_{G \text{ is } C} = \text{const: } \hat{D}^{\frac{N_h}{2}} \quad \text{RVBi} \quad (84)$$

in which

$$\hat{D} = \sum_{ij} g(i-j) \prod_{i,j} Z(i) Z(j) c_i c_j \quad (85)$$

with $g(i-j)$ characterizing the pairing of the twisted holes. Note that in the limit of no pairing, with $g(i-j) = \text{constant}$, equation (84) will recover the pseudogap (spontaneous vortex phase) ground state (54) at $N_h^* = N_h^\# = N_h = 2$.

According to (84), one has $\hat{D} \neq 0$. By using (59) – (62), \hat{D} can be further written as follows

$$\begin{aligned} \hat{D} &= \sum_{ij} g(i-j) \frac{2}{N} \sum_{i,j} e^{i \frac{0}{ij} - i \frac{A_{ij}}{2}} Z(i) Z(j) \frac{\hat{c}_{ij}^0}{2} \\ &= \frac{1}{2} \sum_{ij} g(i-j) \frac{0}{ij} \hat{c}_{ij}^0 \\ &= \frac{1}{2} \sum_{ij} g(i-j) \frac{0}{ij} e^{i \frac{1}{2} (\frac{s_i}{i} + \frac{s_j}{j})} \hat{c}_{ij}^{SC} \end{aligned} \quad (86)$$

such that

$$\hat{D}^D = \frac{1}{2} \sum_{ij} g(i-j) \frac{0}{ij} e^{i \frac{1}{2} (\frac{s_i}{i} + \frac{s_j}{j})} \hat{c}_{ij}^{SC} \neq 0 \quad (87)$$

Since the phase coherence $e^{i \frac{1}{2} (\frac{s_i}{i} + \frac{s_j}{j})} \neq 0$ is established self-consistently once $g(i-j)$ introduces the pairing among twisted holes as discussed above, one finally obtains $\hat{D}^{SC} \neq 0$ according to (87). Note that the phase factor $e^{i \frac{1}{2} (\frac{s_i}{i} + \frac{s_j}{j})}$ also determines the d-wave symmetry of \hat{c}_{ij}^{SC} as discussed previously in Ref.²⁰:

Therefore the superconducting ground state (84) can be obtained as the result of the pairing instability of twisted holes in the pseudogap ground state (54). Simultaneously the phase coherence condition (81) is achieved, and fermionic electron operators in \hat{D} become meaningful as

$$\hat{D} j_{G \text{ is } C} = \sum_{ij} g_{ij} c_i c_j j_{G \text{ is } C}; \quad (88)$$

with the d-wave symmetry characterized by

$$g_{ij} = \frac{1}{2} \frac{0}{ij} g(i-j) e^{i \frac{1}{2} (\frac{s_i}{i} + \frac{s_j}{j})} : \quad (89)$$

Equation (88) also implies the restoration of the coherent fermionic quasiparticle excitations in the superconducting state.

V. SUMMARY AND DISCUSSION

In this paper, we obtained and analyzed the ground state of a pseudogap phase (or spontaneous vortex phase) for a doped Mott insulator, starting from a generalized mean field treatment of the t-J model in the phase string representation. In this representation, the most singular effects of doping, viz., the no double occupancy constraint, and the phase string effect arising from the disordering of the Marshall sign, are sorted out through a systematic procedure. All nontrivial phases in the phase string representation are characterized by a topological gauge structure, which turns out to be well controlled in the regime of interest (low doping). The physical picture of the ground

state that emerges, is that of an underlying RVB vacuum (a spin liquid insulator) which makes the doped holes be twisted into bosonic objects, thereby gaining kinetic energy. The ground state retains time reversal and spin rotational symmetries, and has a simple form (54) in the electron second-quantization representation.

At half filling, the ground state describes an AF Mott insulator, and provides a very accurate description of both long and short range spin correlations of the Heisenberg model. At finite doping, the spin degrees of freedom are described by an RVB spin liquid (55); the mean RVB pair size (42) reduces continuously as doping increases. Charge degrees of freedom emerge at finite doping, with a hidden ODLRO (58). This leads to a finite (electron) pairing amplitude, which characterizes the pseudogap structure of the ground state. The hidden ODLRO in the charge sector is also found to be the underlying reason for the four nodes in the quasiparticle excitation spectrum, but the ground state is non-Fermi-liquid like in the absence of phase coherence. It is more appropriate to call the pseudogap ground state as a spontaneous vortex phase, since free $S = 1/2$ spins that act like vortices prevent superconducting phase coherence in this state.

Two types of low-lying spin excitations above the pseudogap ground state were identified. One is composed of neutral spinons and has a resonance-like feature around (π, π) . It reduces to the conventional gapless spin wave at half filling. The other type of spin excitation exists only in the doped regime and describes a slowly fluctuating incommensurate AF ordering. It provides an alternative explanation of the neutron scattering experiments without invoking a charge stripe scenario.

Superconducting phase coherence is established by an instability which causes RVB pairing of the free spins associated with the twisted holes. The ensuing ground state (84), is non-BCS-like in many aspects. The properties of the superconducting ground state (84), especially those involving non-BCS-like features, are very interesting problems to pursue further.

The class of wavefunctions discussed in this paper provides a unified description of Mott physics, antiferromagnetism, pseudogap physics, and d-wave superconductivity within a single framework. There are many basic distinctions between this ground state and BCS-like wave functions. An important distinction is the independence of the spin excitations from the quasiparticles. The presence of the pairing amplitude and spontaneous vortices associated with spin excitations is another unique feature and is independent of superconducting phase coherence. Furthermore, generalized Jastrow-like factors K in (23) that appear in the ground state wave function, are incompatible with the (Slater) determinantal wave functions of the BCS theory. Therefore, the pseudogap and superconducting ground states obtained in this work represent and describe a new state of matter.

Given that the state (54) undergoes a superconducting instability with phase coherence at low temperature, we may discuss further implications for the global phase diagram. Clearly, the phase coherence condition (82) fails in the spin ordered phase, when the spatial separation of the RVB pair, $\neq 1$; indeed, at half filling, the RVB amplitude W_{ij} shows a power law decay at $|i - j| \neq 1$, without a finite length scale, as discussed in Sec. III A. In this AF phase, phase coherence in (82) due to the tight binding of vortices and antivortices in \mathbb{S}^1 is disrupted. In general, such a spin ordered phase will persist into finite doping at sufficiently low hole concentration, x_c ²⁶. The condition of phase coherence is not satisfied also when the RVB pairing between spins in \mathbb{R}^{VB} is continuously reduced to zero, at sufficiently large doping x_{RVB} . In this case, the underlying rigidity for binding of vortices and antivortices in the ground state is also destroyed and the superconductivity should be suppressed beyond x_{RVB} . How the ground state wavefunction systematically evolves with the doping concentration and the possible existence of quantum critical points at, say, x_c and x_{RVB} , is an important subject for further investigation.

Finally, the results presented in this paper can be used as a starting point to pursue several interesting questions, and we indicate a few possible directions here: (i) The pseudogap (spontaneous vortex phase) ground state is inherently unstable to superconductivity as the result of pairing of the twisted holes. Nonetheless, the pseudogap ground state can still be regarded as a true ground state, for instance, when strong magnetic fields are applied to destroy the RVB pairing of the low-lying spins associated with the backflow of holes. The study of this state in the presence of strong magnetic fields is thus experimentally relevant. (ii) It will be very useful to develop a numerical scheme based on the wavefunction (57) for a variational study. At half-filling, we used the wavefunction to compute the ground-state energy, magnetization, and equal-time spin-spin correlations by employing the loop gas method. As pointed out, this gave us accurate results. However, away from half filling, the RVB amplitude W_{ij} is no longer a real function and loop products generate a fictitious flux proportional to doping concentration [cf. (C28)]. This leads to a sign problem and further work is necessary to overcome this barrier. Havilio and Auerbach studied correlations in doped antiferromagnets using a combination of numerical methods and the Gutzwiller approximation for a different class of RVB wave functions²⁸. A similar study of the wave function proposed in this paper should yield interesting results. (iii) Finally, we note that the form of the wave function (24) follows from very general consequences of a hole moving in an RVB background, since the disordering of the Marshall sign, i.e., the phase string effect, occurs in systems even without AF long range order. Thus the phase string representation and the bosonic RVB mean field theory can be used to study models such as the doped quantum dimer model, where the spin background is known to be RVB paired.

A cknow ledgm ents

We acknowledge helpful discussions with P.W. Anderson, S.P. Kou, H.T. Nieh, X.L. Qi, D.N. Sheng, X.G. Wen, and J. Zaanen. This work is partially supported by the grants of the NSFC, Grant No. 104008 and SRFDP from MOE of China.

APPENDIX A: DETERMINATION OF ρ_h AND Z_h (I)

From the mean-field state (47), one finds

$$\begin{aligned} h_{Gj} h_{GF} = & \sum_{l_1 < l_2}^X \int_h (f_{lh} g)^2 h_{RVB}^Y \sum_h Z_h^0(l_h) Z_h(l_h) h_{l_h} h_{l_h}^Y b_{l_h}^Y b_{l_h} h_{RVB} h_{GF} \\ & \sum_{l_1 < l_2}^X \int_h (f_{lh} g)^2 \sum_h Z_h(l_h) \int_h h_{RVB}^b h_{RVB} h_{GF}. \end{aligned} \quad (A1)$$

In obtaining the last line, we omit RVB pairings involving two holon sites, e.g., the second term on the rhs of the following expression (note that l_h refers to a holon site):

$$h_{RVB}^b h_{l_1}^b h_{l_2}^b h_{RVB} h_{GF} = h_{RVB}^b h_{l_1}^b h_{RVB} h_{GF} h_{RVB}^b h_{l_2}^b h_{RVB} h_{GF} + h_{RVB}^b h_{l_1}^b h_{l_2}^b h_{RVB} h_{GF} h_{RVB}^b h_{l_1}^b h_{l_2}^b h_{RVB} h_{GF} \quad (A2)$$

The essence of this approximation is that for a dilute concentration of holes, the hole-hole correlation induced by the background spin RVB pairing can be neglected, since the average separation between holes sets the upper bound for the size of the RVB pair wave function. For the sake of clarity, we shall invoke this approximation in determining $\rho_h(f_{lh} g)$ and $Z_h(l_h)$ below. In principle, these effects can be incorporated without affecting our results qualitatively.

Now, consider the hopping term (9), which is rewritten as

$$H_t = \sum_{hij}^X \tau e^{iA_{ij}^s - iA_{ij}^0 - iA_{ij}^h} h_i^Y b_i h_j^Y b_j + h.c. \quad (A3)$$

It can be interpreted as describing the hopping of a holon-spinon composite $h_i^Y b_i$ under the influence of the gauge field $A_{ij}^s - A_{ij}^0 - A_{ij}^h$ in the RVB background h_{RVB} ; b_i represents the spinon backflow accompanying the hopping of the holon h_i^Y .

The \circ and $\#$ spinons are paired in the RVB background h_{RVB} , and it costs finite energy to break up an RVB pair at finite doping. Correspondingly, $A_{ij}^s = 0$ and A_{ij}^s remains small in (A3) because A_{ij}^s depicts excitations bound to ± 1 spins. We point out that this condition could fail both, at very low doping, ($x < x^{26}$), where a spin ordered phase exists, and at large doping ($x > x_{RVB}$), where the RVB pairing disappears and A_{ij}^s fluctuates strongly. In both limits, we expect different phases.

Neglecting A_{ij}^s and the hard core repulsion among holons, one may simply choose

$$\rho_h(l_1; l_2; \dots; l_h) = \text{const} \quad (A4)$$

as if the holons are in a Bose condensed state. Consistent with this picture of holon condensation, A_{ij}^h can indeed be treated as describing a constant flux per plaquette. We recall that such an approximation [cf. (32)] was made in determining the RVB amplitudes, W_{ij} .

We can now determine $Z_h(l_h)$ by minimizing $h_{Gj} h_{GF} = h_{Gj} h_{GF}$. Upon invoking the approximations detailed above, we get

$$\frac{h_{Gj} h_{GF}}{h_{Gj} h_{GF}} = \sum_{h=1}^X \sum_{ij}^X \tau e^{-iA_{ij}^0 - iA_{ij}^h} Z_h(i) Z_h(j); \quad (A5)$$

where

$$\tau = \frac{n^b}{2} + \frac{j^s j^2}{2n^b} \tau; \quad n^b = 1; \quad (A6)$$

By optimizing the hopping energy under the condition $\sum_i |Z_h(i)|^2 = 1$; we get the hopping energy

$$h_{Gj} H_{tj} G_{MF} = h_{Gj} G_{MF} = 4t_h N; \quad (A7)$$

with $t_h > 0$ being the maximum eigenvalue of the eigen equation

$$(t_h) Z_h(i) = \frac{t}{4} \sum_{j=nn(i)}^X e^{i \phi_{ij}^0 - i h A_{ij}^h} Z_h(j); \quad (A8)$$

and Z_h , its eigen wave function.

From (A6) we see that t has two contributions: the first term, proportional to $(1 - \langle n \rangle) = 2$; is the probability that the hole hops between nn sites, in a background of uncorrelated spins; the second term, proportional to $\langle j^s j^s \rangle$; is the enhancement from RVB assisted hopping – the initial state (before hole hopping) and the final state have a non vanishing overlap owing to the RVB pairing in the spin background. Note that t' is at small as $n^b \rightarrow 1$; $s \rightarrow 1$. The effective hopping integral t_h can be obtained from (A8), for a fixed hole concentration and under the approximation (32). The numerical results are shown in Fig. 4.

Finally, we note that the eigen values of (A8) would form a Hofstadter spectrum, if the gauge field A_{ij}^h were treated at a mean field level, as a uniform flux (32). However, the density of the back flow spinons is tied to that of the holons and so the fluctuations in A_{ij}^h due to fluctuations in the holon density are self consistently related to the density fluctuations of the spinon back flow. Thus, (A8) resembles an anyon (semion) system in the bosonic representation with a spin index²⁷. The mean field approximation is usually no longer valid in determining its excitation spectrum. This is a nontrivial issue. Nevertheless, since we shall be only interested in its ground state, which is a condensate with a uniform density distribution, all we need to know is that there is a renormalized constant t_h for the effective holon hopping term, estimated as the maximum eigen-value at the mean-field level in (A8). The wave packet $Z_h(i)$ is determined as a linear combination of the wave packets of a cyclotron radius $a_c = a = \frac{b}{\phi}$ in accord with (32) and is made maximally uniform in space, in order to accommodate the condensation of the holes.

APPENDIX B: EFFECTIVE THEORY

In the bosonic RVB mean field theory, the holon many-body wave function ψ_h has been assumed to satisfy the ideal Bose condensation condition (48) at $T = 0$. As we noted in the paper, two effects are omitted in (48), viz., the hard core correlation among holons and the fluctuations of the link field A_{ij}^s . The latter effect should become important when excited spinons are present. Furthermore, ψ_h will also become nontrivial in the presence of an external electromagnetic field.

A general form of the wave function ψ_h in (47) under the influence of A_{ij}^s , and in the presence of an external electromagnetic field A_{ij}^e , can be determined by the following effective hopping term

$$H_h = \sum_{hiji}^X e^{i A_{ij}^s + i A_{ij}^e} h_i^\dagger h_j + h.c.; \quad (B1)$$

If the hard core condition of the holon field is neglected, and if $A_{ij}^e = 0$, (B1) leads to the Bose condensation solution (48) in the ground state. Note that H_h is a gauge model consistent with the gauge invariance (15) in the original Hamiltonian. H_h also respects the spin rotational symmetry as one can easily check that $[H_h, S] = 0$, where S is the total spin operator. Furthermore, in the absence of A_{ij}^e , the time reversal symmetry of H_h can also be shown, by noting that $A_{ij}^s \rightarrow -A_{ij}^s$ under the flip of the spins.

The effective hopping integral t_h appears in (B1) as a renormalized t , which is decided by the spinon back flow according to (49). Here the spinons do not directly see the gauge field A_{ij}^s and the external electromagnetic field A_{ij}^e , because they satisfy a different gauge transformation (16) in (49). Thus, the spinons are truly charge neutral and only carry $S = 1/2$ in the phase string formalism, in contrast to, say, the slave boson gauge theories where both holons and spinons are coupled to the same external electromagnetic field through the Lo e-Larkin rule.

An effective Hamiltonian H_{string} based on H_h and H_s can be written down as

$$H_{string} = H_h + H_s; \quad (B2)$$

Such an effective Hamiltonian has been derived earlier¹⁷, and is the basic low energy effective model for the doping and temperature regimes underpinned by the bosonic RVB order parameter $\phi \neq 0$ in the phase diagram of Fig. 2.

We remark that H_{string} in (B2) is obtained under the assumption that the spinon back flow accompanying the hopping of the holons only provides a renormalized hopping integral t_h . Here t_h is determined at the mean field level,

as the minimum eigen value of (49). However, in principle, the internal excitations of the spinon backflow can effectively reduce t_h according to (49) and thus increase the kinetic energy of H_{string} . This should be taken into account when the high energy part of the charge degrees of freedom are studied. Finally, we point out that the holon and its spinon backflow is actually bound together, as previously shown²⁰ by including the residual interactions of the $t-J$ model beyond the effective Hamiltonian. Such a twisted hole as a bound holon-spinon pair emerges naturally in the present wavefunction approach given in the main text.

APPENDIX C: TIME REVERSAL AND SPIN ROTATIONAL SYMMETRIES OF THE PSEUDOGAP GROUND STATE (??)

1. Time reversal symmetry

The time reversal relation for an $S = 1/2$ single-particle wavefunction is

$$T = \quad ; \quad (C1)$$

Correspondingly, in second quantized language, the time reversal of the electron operators read

$$T c_i^\uparrow T^{-1} = c_i^\downarrow ; \quad (C2)$$

$$T c_i^\downarrow T^{-1} = c_i^\uparrow ; \quad (C3)$$

It is easy to check that

$$T S_i T^{-1} = -S_i ; \quad (C4)$$

Then the time reversal of the RVB state $|\text{RVB}\rangle$ is given by

$$\begin{aligned} T |\text{RVB}\rangle &= \text{const} \sum_{\mathbf{X}} \text{RVB}(\mathbf{1}; \mathbf{2}; \dots; \mathbf{N}) [\mathbf{1} \mathbf{2} \dots \mathbf{N}] c_{\mathbf{1}}^\uparrow c_{\mathbf{2}}^\uparrow \dots c_{\mathbf{N}}^\uparrow \prod_i \mathcal{D}_i \\ &= \text{const} \sum_{\mathbf{f}_{sg}} \text{RVB}(\mathbf{1}; \mathbf{2}; \dots; \mathbf{N}) c_{\mathbf{1}}^\downarrow c_{\mathbf{2}}^\downarrow \dots c_{\mathbf{N}}^\downarrow \prod_i \mathcal{D}_i \\ &= |\text{RVB}\rangle \end{aligned} \quad (C5)$$

by noting that $\mathbf{1} \mathbf{2} \dots \mathbf{N} = (-1)^{N/2}$ in a bipartite lattice; and

$$\begin{aligned} \text{RVB}(\mathbf{1}; \mathbf{2}; \dots; \mathbf{N}) &= \sum_{\mathbf{X}} \sum_{\mathbf{Y}} (-1)^i (-1) W_{ij} \\ &= \sum_{\mathbf{X}} \sum_{\mathbf{Y}}^{\text{pair}(ij)} (-1)^i (-1) W_{ji} \\ &= \sum_{\mathbf{X}} \sum_{\mathbf{Y}}^{\text{pair}(ij)} (-1)^j W_{ji} \\ &= \text{RVB}(\mathbf{1}; \mathbf{2}; \dots; \mathbf{N}) ; \end{aligned} \quad (C6)$$

Furthermore, by using

$$\begin{aligned} T e^{i \hat{S}_i} T^{-1} &= e^{i \frac{1}{2} [\begin{smallmatrix} s_i & 0 \\ i & h_i \end{smallmatrix}]} (-1)^{\hat{N}_h} (-1)^i \\ &= e^{i \hat{S}_i} e^{i \frac{0}{2}} (-1)^{\hat{N}_h} (-1)^i \end{aligned}$$

according to Eq.(5) and $c_i = e^{i \hat{S}_i} c_i$, one finds that

$$T c_i T^{-1} = c_i e^{i \frac{0}{2}} (-1)^{\hat{N}_h} (-1)^i ; \quad (C7)$$

Finally, the time reversal of the ground state $|\psi_g\rangle$ is given by

$$\begin{aligned}
T j_G i &= T \sum_1 X_{Z''(l) e_{1''}}^{! N_h''} X_{Z_{\#}(l^0) e_{1^0 \#}}^{! N_h^{\#}} T^{-1} T \mathcal{R} V B i \\
&= \sum_1 X_{Z''(l) e_{1''}}^{! N_h''} e^{i \frac{0}{1} (1)^{N_h''} (1)^1} X_{Z_{\#}(l^0) e_{1^0 \#}}^{! N_h^{\#}} e^{i \frac{0}{1^0} (1)^{N_h^{\#}} (1)^0} \mathcal{R} V B i \\
&= (1)^{N_h''} (1)^{N_h^{\#}} (N_h'' - 1) = 2 \sum_1 X_{Z_{\#}(l) e_{1\#}}^{! N_h''} X_{Z''(l^0) e_{1^0''}}^{! N_h^{\#}} \mathcal{R} V B i \\
&= j_G i
\end{aligned}$$

at $N_h'' = N_h^{\#} = N_h = 2$ (or $S^z = 0$). In obtaining the third line, the condition (62) is used.

Thus, we have proved the time reversal symmetry of the pseudogap ground state (54), for $S^z = 0$.

2. Spin rotational symmetry

The condition for a state with $SU(2)$ spin rotational symmetry is

$$S^2 j_i = 0 \quad (C8)$$

or equivalently,

$$S^z j_i = 0; \quad (C9)$$

$$S j_i = 0; \quad (C10)$$

Here S is the total spin and $S = S^x i^x + S^y i^y + S^z i^z$, where S^x, S^y and S^z are the three components of S . In the following we shall consider $j_G i$ with $S^z = 0$; and show that $S j_G i = 0$:

By noting that, for $i \notin l$,

$$S_i^+ e^{i \frac{1}{2} s_i} = e^{i \frac{1}{2} s_i} S_i^+ e^{i i(i)} \quad (C11)$$

$$S_i^+ e^{i \frac{1}{2} h_i} = e^{i \frac{1}{2} h_i} S_i^+ \quad (C12)$$

we have

$$\begin{aligned}
S_i^+ e_l &= S_i^+ e^{i \wedge_1} e_l \\
&= e^{i i(i)} e_l S_i^+ (1 - i_l)
\end{aligned} \quad (C13)$$

and

$$S^+ e_l = e_l \sum_{i \notin l} S_i^+ e^{i i(i)}; \quad (C14)$$

Then according to (52) and (48),

$$\begin{aligned}
S^+ j_G i &= S^+ \sum_{f l_h g h} X_{Z_h(l_h) e_{l_h h}}^Y \mathcal{R} V B i \\
&= \sum_{f l_h g h} X_{Z_h(l_h) e_{l_h h}}^Y \otimes \sum_{i \notin l_h} S_i^+ e^{i \frac{P}{l_h} i l_h(i)} \mathcal{R} V B i^A; \quad (C15)
\end{aligned}$$

In terms of (55) and (56), one has

$$\begin{aligned}
\mathcal{R} V B i &= \sum_{f s g} X_{\mathcal{R} V B (1; 2; N)}^Y C_{11}^Y C_{22}^Y \frac{Y}{N} \mathcal{D} i \\
&\text{const:} \sum_{f s g \text{ pair } (ij)} X_{(1)^j} X_{ij}^Y (1)^j w_{ij} j; i''; j \#; i; \quad (C16)
\end{aligned}$$

Substitute the above solution into W_{ij} ,

$$\begin{aligned} W_{ij} &= \sum_m \frac{v_m}{u_m} w_m(i) w_m(j) \\ &= \sum_m \frac{v_m}{u_m} e^{im_y(i_y - j_y)} g_{m_x, m_y}(i_x) g_{m_x, m_y}(j_x): \end{aligned}$$

Then

$$\begin{aligned} W_{i+; j+} &= \sum_m \frac{v_m}{u_m} e^{im_y(i_y - j_y)} g_{m_x, m_y}(i_x + x) g_{m_x, m_y}(j_x + x) \\ &= \sum_m \frac{v_m}{u_m} e^{im_y(i_y - j_y)} g_{m_x, m_y}(i_x) g_{m_x, m_y}(j_x) \\ &= \sum_m \frac{v_m}{u_m} e^{i(m_y + x)(i_y - j_y)} g_{m_x, m_y}(i_x) g_{m_x, m_y}(j_x) \\ &= e^{i x (i_y - j_y)} W_{ij}: \end{aligned}$$

and it is straightforward to show

$$W_{ij} = \prod_{\text{loop}} w_{ij} e^{i \sum_{\text{loop}} A_{ij}^h}; \quad (\text{C } 28)$$

on a loop $(i \rightarrow j \rightarrow j+ \rightarrow i+ \rightarrow i)$. One can always choose a proper gauge such that (C 18) holds on any link $(i; j)$.

- ¹ P. W. Anderson, Science 235, 1196 (1987).
- ² G. Baskaran, Z. Zou, and P. W. Anderson, Solid State Commun. 63, 973 (1987).
- ³ C. Gros, Phys. Rev. B 38, 931 (1988).
- ⁴ F. C. Zhang, C. Gros, T. M. Rice, and H. Shiba, Supercond. Sci. Technol. 1, 36 (1988).
- ⁵ H. Yokoyama and M. Ogata, Phys. Rev. B 65, 3615 (1996).
- ⁶ A. Paramekanti, M. Randeria, and N. Trivedi, Phys. Rev. Lett. 87, 217002 (2001).
- ⁷ P. W. Anderson, P. A. Lee, M. Randeria, T. M. Rice, N. Trivedi, and F. C. Zhang, J. Phys.: Condens. Matter 16, R 755 (2004).
- ⁸ S. Liang, B. Doucot, and P. W. Anderson, Phys. Rev. Lett. 61, 365 (1988).
- ⁹ W. Marshall, Proc. Roy. Soc. (London) A 232, 48 (1955).
- ¹⁰ B. Sutherland, Phys. Rev. B 37, 3786 (1988).
- ¹¹ Z. Y. Weng, D. N. Sheng, Y. Chen, and C. S. Ting, Phys. Rev. B 55, 3894 (1997); D. N. Sheng, Y. C. Chen, and Z. Y. Weng, Phys. Rev. Lett. 77, 5102 (1996).
- ¹² Z. Zou and P. W. Anderson, Phys. Rev. B 37, 627 (1988).
- ¹³ A. Auerbach and D. P. Arovas, Phys. Rev. Lett. 61, 617 (1988); Sanjy Sarker, C. Jayaprakash, H. R. Krishnamurthy, and Michael Ma, Phys. Rev. B 40, 5028 (1989); Daijiro Yoshioka, J. Phys. Soc. Jpn. 58, 32 (1989).
- ¹⁴ Z. Y. Weng, D. N. Sheng, and C. S. Ting, Phys. Rev. B 52, 637 (1995); Mod. Phys. Lett. B 8, 1353 (1994).
- ¹⁵ Q. H. Wang, Phys. Rev. Lett. 92, 057003 (2004); Chin. Phys. Lett. 20, 1582 (2003).
- ¹⁶ S. P. Kou, X. L. Qi, and Z. Y. Weng, Phys. Rev. B 71, 235102 (2005).
- ¹⁷ Z. Y. Weng, D. N. Sheng, and C. S. Ting, Phys. Rev. B 59, 8943 (1999); Phys. Rev. Lett. 80, 5401 (1998).
- ¹⁸ W. Q. Chen and Z. Y. Weng, Phys. Rev. B 71, 134516 (2005).
- ¹⁹ Note that the naive form of $\mathcal{H}S_{\text{H}} = \sum_{i_h} \mathcal{R}VB_{i_M F}$ does not optimize the kinetic energy of the hole. This can be seen most directly by calculating $\mathcal{H}S_{\text{H}} \mathcal{H}_t \mathcal{H}S_{\text{H}}$, where \mathcal{H}_t is the hole hopping term in the tight-J Hamiltonian, (9). Since $\mathcal{H}_t^Y b_{i_h n} = 0$, the kinetic energy of the hole vanishes. In the earlier approach¹⁷, to avoid this difficulty, a hidden symmetry in $\mathcal{R}VB_{i_M F}$ has to be used to get a finite and optimized $\mathcal{H}S_{\text{H}} \mathcal{H}_t \mathcal{H}S_{\text{H}}$; which makes $\mathcal{R}VB_{i_M F}$ itself also explicitly depend on the hole configuration.
- ²⁰ Yi Zhou, V. N. Muthukumar, and Zheng-Yu Weng, Phys. Rev. B 67, 064512 (2003).
- ²¹ Z. Y. Weng and V. N. Muthukumar, Phys. Rev. B 66, 094509 (2002).
- ²² V. N. Muthukumar and Z. Y. Weng, Phys. Rev. B 65, 174511 (2002).
- ²³ With the increase of the hole concentration, the "hard-core" condition among bosonic c_{H} 's will gradually become important and eventually the above "ideal" Bose condensation approximation has to be modified.
- ²⁴ This kind of spin excitation involving the twisted holes has some similarity to the usual d-wave BCS theory as in both cases the spin excitation is made of nodal quasiparticles. However, there will be an important distinction since quasiparticles in the present case are no longer elementary in the conventional sense as pointed out in the text.

- ²⁵ Z. Y. W eng, D. N. Sheng, and C. S. T ing, Phys. Rev. B 59, 11367 (1999).
- ²⁶ S. P. K ou and Z. Y. W eng, Phys. Rev. Lett. 90, 15700 (2003).
- ²⁷ X. G. W en and A. Zee, Phys. Rev. B 41, 240 (1990).
- ²⁸ M. oshe H avilio and A. ssa A uerbach, Phys. Rev. B 62, 324 (2000).