

Conformal Field Theory Approach to the Two-Channel Anderson Model

H. Johannesson^a, N. Andrei^b, and C. J. Boolech^c

^aTheoretical Physics, Chalmers and Goteborg University, Goteborg, Sweden

^bCenter for Materials Theory, Rutgers University, U.S.A.

^cSchool of Physics, University of Geneva, Switzerland

Abstract

The two-channel Anderson impurity model serves as a prototype for describing heavy-fermion materials with a possible mixed-valent regime with both quadrupolar and magnetic character. We report on the low-energy physics of the model, using a conformal field theory approach with exact Bethe Ansatz results as input.

Key words: Heavy fermions, Anderson model, two-channel Kondo physics

PACS: 71.27.+a, 75.20.Hr, 75.40.-s

As is well-known, several uranium-based heavy fermion materials exhibit manifest non-Fermi liquid behavior at low temperatures. A particularly intriguing case is that of the UBe_{13} compound. It has been suggested that the electric quadrupolar degrees of freedom of the uranium ions (in $5f^2$ configurations) in the Be_{13} host are overscreened by the local orbital motion of the conduction electrons, producing a two-channel Kondo response of this material [1]. Taking into account the possibility that the magnetic $5f^3$ configurations may also be at play, one is led to consider the two-channel Anderson impurity model [1]

$$H = H_0 + \sum_s f^y f + \sum_q b^y b + V \left(\sum_s (0) b^y f + f^y b \right) \quad (0); \quad (1)$$

where H_0 is the free part of the Hamiltonian. The conduction electrons f^y carry spin ($= \uparrow; \downarrow$) and quadrupolar ($= \square$) quantum numbers, and hybridize with a local uranium ion via a matrix element V . The ion is modeled by a quadrupolar [magnetic] doublet of energy ϵ_q [ϵ_s], created by a boson [fermion] operator b^y [f^y]. Strong Coulomb repulsion implies single occupancy of the localized levels: $f^y f + b^y b = 1$.

We have carried out a nonperturbative analysis of the model, exploiting boundary conformal field theory [2] to trade the hybridization interaction in (1) for a scale invariant boundary condition on the conduction electrons. Identifying the proper boundary condition immediately identifies the critical theory to which the model belongs. By a numerical fit to the exact Bethe Ansatz solution [3] of (1) we can determine all parameters and scales of this theory, thus obtaining a complete descrip-

Corresponding author.

E-mail: johannesson@fy.chalmers.se

tion of the low-energy dynamics of the model. Our most important results can be summarized:

The model renormalizes to a line of fixed point Hamiltonians parameterized by the average charge n_c of f orbitals at the impurity site (Fig. 1), all exhibiting the same zero-temperature impurity entropy $S_{imp} = k_B \ln \sqrt{2}$, typical of two-channel Kondo physics [4]. The low-temperature specific heat induced by the impurity takes the form $C_{imp} = \gamma_s T \ln(T_s/T) + \gamma_q T \ln(T_q/T) + O(T^2)$, where γ_s and T_s are amplitudes and temperature scales respectively, measuring the participation of the spin [quadrupolar] degrees of freedom in the screening process. Similarly, the impurity response to a magnetic [quadrupolar] field is also of two-channel Kondo type, $\chi_{imp} = \chi_s \ln(T_s/T) + \chi_q \ln(T_q/T)$, with the amplitudes χ_s decreasing exponentially with n_c . We have also calculated the Fermi edge singularities caused by time-dependent hybridization between conduction electrons and impurity. These are coded by the scaling dimensions of the pseudo-particles b^y and f^y in (1), and we find that $x_b = (3 + 2n_c^2)/16$ and $x_f = (5 - 4n_c + 2n_c^2)/16$ respectively. We point out that the values of the pseudo-particle scaling dimensions as obtained by various approximation schemes, e.g. NCA and large-N calculations [5], deviate from our exact result.

The conformal field theory formalism allows us to determine also the asymptotic dynamical properties of the model, incl. Green's functions and resistivities. We have calculated the self-energy of the electrons and find that it leads to a zero-temperature resistivity independent of the level separation: $\rho(T=0) = 3\eta_1/4 (e v_F)^2$, with η_1 the (dilute) impurity concentration, ρ_0 the density of states at the Fermi level, and v_F the Fermi velocity. The finite-temperature calculation is in progress.




Fig. 1. Schematic renormalization group flow diagram of the two-channel Anderson impurity model. The flow connects the line of microscopic theories characterized by the ratio $n_c = (n_s - n_q) / (n_s + n_q)$ with the line of infrared fixed points parameterized by n_c .

References

- [1] For a review, see D. L. Cox and A. Zawadowski, *Adv. Phys.* 47, 599 (1998).
- [2] I. A. A. Eck and A. W. W. Ludwig, *Nucl. Phys. B* 360, 641 (1991).
- [3] C. J. Bolech and N. Andrei, *Phys. Rev. Lett.* 88, 237206 (2002).
- [4] H. Johannesson, N. Andrei, and C. J. Bolech, *Phys. Rev. B* 68, 075112 (2003).
- [5] D. L. Cox and A. E. Ruckenstein, *Phys. Rev. Lett.* 71, 1613 (1993).