Exact Analysis of Function Attractive Fermions and Repulsive Bosons in One D imension

Toshiaki Iida and M iki W adati

Department of Physics, Graduate School of Science, University of Tokyo, Hongo 7–3–1, Bunkyo-ku, Tokyo 113–0033, Japan

(M arch 2, 2005)

A bstract

The G audin integral equation for the ground state of a one-dimensional -function attractive spin-1=2 fermions is solved in the form of power series. The rst few terms of the asymptotic expansions for both strong and weak coupling cases are calculated analytically. The physical quantities such as the ground state energy are expressed in terms of a single dimensionless parameter = c=D, where c is the coupling constant and D is the number density. The results agree with those obtained from the perturbation calculations, which include the one in the classical electrostatics originally by K include . In the strong coupling limit, the connection to the solutions of the Lieb-Liniger integral equation for the ground state of a one-dimensional -function repulsive bose gas is shown explicitly.

KEYW ORDS: -function gas, one-dimensional integrable systems, BCS-BEC crossover, Lieb-Liniger integral equation, Gaudin integral equation, Tonks-Girardeau gas, hard-core bose gas

1 Introduction

We discuss a one-dimensional system of quantum N particles with -function interaction in a periodic box of length L. The Ham iltonian of the system is

$$H = \sum_{j=1}^{X^{N}} \frac{2m}{2m} \frac{e^{2}}{e^{2}x_{j}^{2}} + \sum_{i \in j}^{X} (x_{i} - x_{j}); \qquad (1.1)$$

where m is the mass and c is the coupling constant. Hereafter, we choose a unit, $\sim = 1$ and 2m = 1 to simplify the expressions.

The system (1.1) has been studied extensively by use of the Bethe ansatz method.¹⁹ For N repulsive (i.e. c > 0) bosons, the ground state is described by the integral equation for the distribution function (k) of the quasi-m omenta,¹

$$(k) = \frac{1}{2} + \frac{c}{k} \sum_{K}^{K} \frac{(q) dq}{c^{2} + (k - q)^{2}}; \qquad (1.2)$$

where K is a cut-o momentum, which is xed by the normalization condition, $Z_{\rm rec}$

$$(k) dk = D$$
: (1.3)

iida@monet.phys.s.u-tokyo.ac.jp

^ywadati@monet.phys.su-tokyo.ac.jp

Here, D N=L is the number density of the particles. The interval [K;K] is led with quasi-m on enta k. The ground state energy is given by

$$\frac{E_{B}}{L} = \int_{K}^{L} k^{2} (k) dk:$$
(1.4)

W e call (12) the Lieb-Liniger integral equation.

In 1967, Yang⁴ proposed a method, which we now call the generalized Bethe ansatz, or \nested " Bethe ansatz, to study the system (1.1) with arbitrary internal degree of freedom. For N attractive (i.e. c < 0) spin-1=2 fem ions, the ground state with total spin 0 is described by³

$$f(q) = 2 + \frac{c}{2} \int_{Q}^{Z} \frac{f(q^{0}) dq^{0}}{c^{2} + (q - q^{0})^{2}}:$$
 (1.5)

Here, f(q) is the distribution of spins with spin-wave rapidity q, and a cut-o rapidity Q is xed by the norm alization condition for f(q):

$$\frac{1}{2} \int_{Q}^{Z} f(q) dq = D; \qquad (1.6)$$

while the ground state energy is given by

$$\frac{E_{F}}{L} = \frac{1}{2} \int_{Q}^{2} q^{2} \frac{c^{2}}{4} f(q) dq; \qquad (1.7)$$

For our purpose, it is convenient to introduce the e ective ground state energy,

$$\frac{E_{F}^{e}}{L} = \frac{E_{F}}{L} + \frac{c^{2}}{4}D = \frac{1}{2} \int_{Q}^{2} q^{2}f(q) dq; \qquad (1.8)$$

W e call (1.5) the G audin integral equation.

By use of a method proposed in the previous paper¹⁰ to solve (1.2), we explicitly calculate the rst few terms of the asymptotic expansions of f(q) for both strong and weak coupling cases. It is essential to use the single dimensionless coupling constant , de ned by

$$= \frac{c}{D} = \frac{2m c}{\sim^2 D}:$$
 (1.9)

C on bining the results with those in the previous papers, 10,11 we discuss in particular four cases, ! 1, ! 0, ! + 0, and ! + 1. We also study the physical quantities, such as the ground state energy and the chem ical potential, as functions of .

This paper is organized as follows. In x 2, we solve (1.5) for large and small j j. The similarities and the connections of the obtained expressions in the case of ! 1 for (1.5) and

! +1 for (12) are investigated in x 3. The recent discussion on $\BCS-BEC$ crossover¹²⁽¹⁴ is made clear for these integrable models in one-dimension. For the small j jcase, it is known to be di cult to solve (12) and (1.5). Analogies with the perturbation calculation in the classical electrostatics originally by K irchho are discussed in x 4. The last section is devoted to the conclusions. To concentrate on physical discussions, most of the calculations are given in Appendices A C.

2 Solutions of the G aud in Integral E quation

2.1 Form ulation

Following a method presented in the previous papers, 10,11 we solve the G audin integral equation (1.5). In terms of new variables,

$$q = Qx;$$
 $c = Q;$ $q' = Qy;$ (2.1)

$$f(Qx) = F(x);$$
 (2.2)

(1.5) becom es

$$F(y) + - \frac{Z_{1}}{\frac{F(x) dx}{2 + (x - y)^{2}}} = 2:$$
 (2.3)

W e also refer to (2.3) as the G audin integral equation. A sm entioned, we use the num ber density D and the dimensionless coupling constant ,

$$D = \frac{N}{L}; \qquad (2.4)$$

$$= \frac{C}{D}:$$
 (2.5)

The large (sm all) corresponds to the large (sm all) j j. By expanding

$$F(x) = \int_{n=0}^{X^{2}} a_{n} x^{2n}; \qquad (2.6)$$

we have

$$Z_{1}^{2} dx \frac{F(x)}{2 + (x - y)^{2}} = \frac{X^{l} X^{l} X^{p}}{n = 0 p = 0 m \frac{m}{p} 0} \frac{a_{n} 22 + 1}{(2m)! (p - m)!} + (1)^{m + n} 2(m + n)! arctan \frac{1}{2} y^{2p}; \quad (2.7)$$

where $s = {}^{2}$. Substituting (2.6) and (2.7) into (2.3), we get

In order that (2.8) holds for any y, 1 y 1, we equate the term s of the same power of $_{Y}^{2p}$. At y^{0} and y^{2p} (p 1), we respectively obtain

$$\begin{aligned} a_{0} + \frac{2}{n} \arctan \frac{1}{n} = 0 \\ &+ \frac{2}{n} \frac{x^{4}}{n} = 1 \\ a_{n} = 1 \\ &= 1 \end{aligned} \begin{pmatrix} x^{n} & \frac{(1)^{j+1} 2(1-1)}{2(n-1)+1} + (1)^{n-2n-1} \arctan \frac{1}{n} = 2; \\ &= 2; \\ a_{p} + \frac{2}{n} \frac{x^{4}}{n} \frac{x^{p}}{(2m)! (p-m)! (p-m)$$

In this way, a solution of (2.9) gives a solution F(x) of (2.3).

 $\tt U\,ntil\,now$, all the expressions are exact and no approximation is employed. We summarize here the method of solution:

1. Solve (2.9) to determ ine fang of (2.6) in the case of large $% \left[\left(2.9\right) \right) =\left[\left(2.9\right) \right] \left(1.9\right) \left(2.9\right) \right]$ and small $\left(1.9\right) \left(2.9\right) \left(1.9\right) \left(1.9$

2. Use (1.6) with (2.1), which can be rewritten as

$$= \frac{2}{1} \frac{1}{1} ; (2.10)$$

to obtain as a function of .

3. Calculate other physical quantities from the expressions of (1.6), (2.1), (1.8) and therm odynam ic identities² in the following form;

$$D = \frac{Q}{2} \int_{1}^{Z} F(x) dx; \qquad (2.11)$$

$$\frac{E_{F}^{e}}{L} = \frac{Q^{3}}{1} x^{2} F(x) dx; \qquad (2.12)$$

$$= D^{2} 3e \frac{de}{d}$$
; (2.13)

$$v = 2D$$
 3e $2\frac{de}{d} + \frac{1}{2} \frac{2^2 d^2 e}{d^2}$ ¹⁼²; (2.14)

where $E_F = (N D^2)$, and v are the scaled energy of the ground state, the chem ical potential and the sound velocity, respectively.

2.2 Strong coupling case

We consider the large case. In this case, only a few a_n 's are enough to have good approximations. Indeed, it can be shown that $a_n = 0$ (1 = 2n+1) for n = 1. Therefore, we only keep a_0 , a_1 , and set $a_n = 0$ (n = 2),

$$F(x) = a_0 + a_1 x^2$$
: (2.15)

Equations (2.9) give

$$a_0 + \frac{2}{3} \arctan \frac{1}{2} a_0 + \frac{2}{3} \frac{1}{5^2} a_1 = 2;$$
 (2.16a)

$$a_0 \quad \frac{2}{3}a_0 \quad \frac{2}{3}a_1 = 0;$$
 (2.16b)

and consequently,

$$a_0 = 2 \quad 1 \quad \frac{2}{2} + \frac{4}{2 \quad 2} + \frac{2 \quad 2}{3 \quad 3 \quad 3} \quad ;$$
 (2.17a)

$$a_1 = -\frac{4}{3}$$
: (2.17b)

Using (2.15) with (2.17) in (2.10), we obtain as a function of . The result is

$$= \frac{2(2+1)}{6(2+1)^3} \quad 1 \quad \frac{2}{6(2+1)^3} \quad (2.18)$$

 $T \ he \ expressions \ of \ a_0 \ and \ a_1 \ in \ term \ s \ of \ are$

$$a_0 = \frac{2 + 1}{6(2 + 1)^4}$$
; (2.19a)

$$a_1 = \frac{2}{2(2 + 1)^3}$$
: (2.19b)

The ground state energy is calculated from (2.12),

$$\frac{E_{F}^{e}}{L} = D^{3} \frac{2}{12} \frac{2}{2+1} + \frac{4^{2}}{15(2+1)^{3}} : \qquad (2.20)$$

The cut-o rapidity Q, the chemical potential and the sound velocity v are given by

$$Q = \frac{D}{2} \frac{1}{2 + 1} + \frac{2}{6(2 + 1)^3} ; \qquad (2.21)$$

$$e = D^{2} \frac{2}{12} \frac{6}{2} + \frac{1}{2} \frac{6}{2} + \frac{1}{1} \frac{2}{2} + \frac{1}{1} \frac{2}{1} + \frac{8^{2}}{5(2+1)^{2}(6+1)} ; \qquad (2.22)$$

$$v = 2 D \frac{2}{2 + 1} + \frac{2^2}{3(2 + 1)^3}$$
; (2.23)

where we have introduced the elective chemical potential e^{-} , e^{-} + (1=4)D²².

2.3 W eak coupling case

This is the small case. The low est order approximation of (2.9) is

$$2a_0 = 2;$$
 (2.24a)

$$2a_p = 0; p = 1:$$
 (2.24b)

A derivation of (2.24) from (2.9) is the same as shown before.^{10,11} Another derivation is given in Appendix A, where we treat the integral in a more direct manner in the small case. Equations (2.24) simply gives

$$F(x) = 1$$
: (2.25)

W e list here the analysis based on (2.25):

$$= \frac{2}{2};$$
 $Q = \frac{D}{2};$ (2.26)

$$\frac{E_{F}^{e}}{L} = D^{3} \frac{2}{12}; \qquad e = D^{2} \frac{2}{4}; \qquad v = D: \qquad (2.27)$$

It should be noted that, in contrast to the solution of the Lieb-Liniger integral equation (1.2), all the coe cients a_n are nite at ! 0.¹⁰ This nice analytic property can be seen from (2.3), since the kernel = f [² + (x y²)]g becomes an expression of (x y) in the limit ! 0.

The second order approximation of (2.9) is

$$2a_{p} + \frac{2}{n} \frac{\chi}{2n} \frac{2p+1}{2n} a_{n} = 2_{p0}; p = 0;$$
 (2.28)

Here, $_{p0}$ is the K ronecker's delta. A derivation of (2.28) is also given in the previous papers^{10,11} and in Appendix A.W e set

$$a_n = a_n + a_n^{(1)} + o(); n 0;$$
 (2.29)

where $_{n0}$ stands for the solution of (2.24) and $a_n^{(1)}$ denote the corrections of the 1 order. Substitution of (2.29) into (2.28) yields

$$2a_{p}^{(1)} = 0; p = 0; (2.30)$$

which leads to

F (x) =
$$1 + -\frac{x^{2}}{x^{2n}} + o()$$

= $1 + -\frac{1}{1 + x^{2}} + o()$: (2.31)

This agrees with the approximation shown in the Gaudin's book.⁷ From (2.10), F (x) is integrated to give

$$=$$
 2 + $\frac{2b}{2}$; (2.32)

where the divergent sum b is de ned by

$$b = \sum_{n=0}^{X} \frac{1}{2n+1}$$
 (2.33)

Equation (2.32) is solved to obtain as a function of:

$$=$$
 $\frac{2}{-1}$ $\frac{2b}{2}$: (2.34)

W e proceed to calculate the physical quantities. From (2.11) and (2.34), we have

$$Q = \frac{D}{2} \quad 1 \quad \frac{2b}{2} \quad :$$
 (2.35)

The ground state energy is obtained from (2.12), (2.31) and (2.35):

$$\frac{E_{F}^{e}}{L} = \frac{Q^{3}}{2} + \frac{2}{3} + \frac{2}{2} + \frac{X^{4}}{2n+3} + \frac{1}{2n+3}$$

$$= D^{3} \frac{2}{12} + \frac{12}{2} + \frac{X^{4}}{2n+3} + \frac{1}{(2n+1)(2n+3)}$$

$$= D^{3} \frac{2}{12} + \frac{6}{2} + \frac{1}{2} + \frac{6}{2} + \frac{1}{2} + \frac{1}{$$

This recovers the previous result¹⁶ where the integrals are directly treated without obtaining the explicit form of F (x). It is remarkable that we observe the cancellation of the divergent sum s as before.^{10,11} That is, the physical quantities are expressed in power series of without any divergent sum. The chem ical potential and the sound velocity can be calculated from the form ulae (2.13) and (2.14),

$$e = D^2 \frac{2}{4} + \frac{4}{2}$$
; (2.37)

$$v = D 1 + \frac{1}{2}$$
: (2.38)

The limit ! 0 gives the well-known results for free ferm ions.

In g. 1, we plot the solutions (2.15) with (2.19) and (2.31) with (2.34), for several values of . For comparison, we also plot the numerical solutions based on the classical quadrature method.^{1,17} In this gure, we choose a unit, D = 1 so that Q=D takes a value between =4 and =2 (see (2.21) and (2.35)). For a xed Q and 1 x 1, q and f (q) are determined by (2.1) and (2.2), respectively. Numerical values of are obtained through the relation =

Q=D. In the weak coupling case, it is fair to say that the error is large near the end-points, $jq \neq D / Q=D$. This can be observed analytically when the solution (2.31) is substituted into the integral equation (2.3).

3 The BCS-BEC Crossover

Recently, there is a renewed interest in the crossover from a Bardeen-Cooper-Schrie er (BCS) super uid with Cooper pairs to a Bose-Einstein Condensate (BEC) of molecules composed of



Figure 1: The normalized distribution function of spins f (q) in the ground state for various values of . Num erical results are indicated by points (+, etc.), while the analytical ones, (2.15) and (2.31), are by lines ($| \{, | \{ etc). Num erical solutions are obtained by a classical method.^{1,17} For sm all , analytical solutions are shown for the intervals [Q=D + j j=4;Q=D j j=4] of q=D, which correspond to [1+=4;1==4] of x. For intermediate , = 1:136 and = 2:239, only the num erical results are shown. For large , analytical and num erical results are indistinguishable.$

tightly bound ferm ion pairs.^{13,18} The experim ental realizations of low-dimensional trapped bose gases at ultracold temperatures also inspired the controversy even in one-dimension.^{12,14} The analytic solutions in x 2 of the G audin integral equation and in the previous paper¹⁰ of the Lieb-Liniger (LL) integral equation enable us to make the quantitative discussion. To avoid confusion, physical quantities, such as the mass m, the number of particles N, the number density D, the coupling constants , , and the ground state energy E are denoted with the subscripts F and B for a ferm igas with spin-1=2 and a bose gas, respectively.

The dimensionless form of the LL integral equation (12) reads 1,10

$$g(y) \quad \frac{B}{2} \int_{1}^{2} \frac{g(x) dx}{B} = \frac{1}{2}; \quad (3.1)$$

where

$$g(x) = (K x); \quad k = K x; \quad 2m_B c = K_B:$$
 (3.2)

The relation between $_{\rm B}$ and $_{\rm B}$ is

$$_{B} = \frac{2m_{B}C}{D_{B}} = \frac{Z_{1}}{Z_{1}};$$
(3.3)
$$g(x) dx$$

where $D_B = N_B = L$. The ground state energy is given by

$$\frac{E_{B}}{L} = \frac{K^{3}}{2m_{B}} \int_{-1}^{L} x^{2} g(x) dx: \qquad (3.4)$$

A salready mentioned in the G audin's paper,³ strong coupling ferm ions described by the G audin integral equation (2.3) may be regarded as the repulsive bosons of dimers. To exam ine whether it is indeed true, we establish a relation between the exact solutions of the G audin integral equation (2.3) and the LL integral equation (3.1). In the following, we concentrate ourselves on the strong coupling case, i.e. the large $_{\rm F}$ case for ferm ions and large $_{\rm B}$ case for bosons are considered. For explicit com parison, we sum marize here the results shown in x 2.2 and the previous papers.^{10,11}

For $_{\rm F}$ 1 (ferm ions with strong attraction),

$$f(q) = \frac{2_{F} + 1}{F} - 1 - \frac{2_{F}}{6(2_{F} + 1)^{4}} - \frac{1}{\frac{3}{F}D_{F}^{2}}q^{2};$$
(3.5a)

$$_{\rm F} = \frac{4_{\rm F} + 2}{6(2_{\rm F} + 1)^3} ; \qquad (3.5b)$$

$$Q = \frac{D_F}{2} \frac{F_F}{2_F + 1} + \frac{2}{6(2_F + 1)^3} ;$$
 (3.5c)

$$\frac{E_{F}^{e}}{L} = \frac{D_{F}^{3}}{2m_{F}} \frac{2}{12} \frac{F}{2F+1} + \frac{4^{2}}{15(2F+1)^{3}}; \qquad (3.5d)$$

$${}^{e}_{F} = \frac{D_{F}^{2}}{2m_{F}} \frac{{}^{2}}{12} \frac{6_{F}+1}{2_{F}+1} \frac{}{2_{F}+1} \frac{}{2_{F}+1} \frac{}{2_{F}+1} \frac{}{1+\frac{8_{F}^{2}}{5(2_{F}+1)^{2}(6_{F}+1)}};$$
(3.5e)

$$v_{\rm F} = \frac{2 D_{\rm F}}{2m_{\rm F}} \frac{F}{2 F + 1}^2 1 + \frac{2^2}{3(2 F + 1)^3}$$
: (3.5f)

For $_{\rm B}$ 1 (bosons with strong repulsion),

$$(k) = \frac{B+2}{2B} \quad 1 \quad \frac{2^{2}}{3B(B+2)^{2}} \quad \frac{1}{B^{3}D_{B}^{2}}k^{2}; \quad (3.6a)$$

$$_{\rm B} = \frac{_{\rm B} + 2}{3(_{\rm B} + 2)^3} \quad ; \tag{3.6b}$$

$$K = D_{B} \frac{B}{B+2} + \frac{4^{2}}{3(B+2)^{3}} ; \qquad (3.6c)$$

$$\frac{E_{B}}{L} = \frac{D_{B}^{3}}{2m_{B}} \frac{2}{3} - \frac{B}{B+2} \frac{2}{1+\frac{32^{2}}{15(B+2)^{3}}}; \qquad (3.6d)$$

$$_{B} = \frac{D_{B}^{2}}{2m_{B}} \frac{3_{B} + 2}{B + 2} \frac{2}{3} - \frac{B}{B + 2} \frac{2}{1 + \frac{64^{2}}{5(B + 2)^{2}(3_{B} + 2)}}; \quad (3.6e)$$

$$v_{\rm B} = \frac{2 D_{\rm B}}{2m_{\rm B}} - \frac{B_{\rm B}}{B+2} + \frac{16^2}{3(B+2)^3}$$
 : (3.6f)

Identifying N $_{\rm F}$ ferm ions with mass m $_{\rm F}$ with N $_{\rm B}$ bosons of dimers with mass m $_{\rm B}$, we have

$$m_{B} = 2m_{F};$$
 $N_{B} = \frac{N_{F}}{2};$ $D_{B} = \frac{D_{F}}{2}:$ (3.7a)

Hence, from (1.9) we have

$$_{\rm B} = 4_{\rm F}$$
: (3.7b)

Substitution of (3.7) into (3.5d) gives

$$\frac{E_{F}^{e}}{L} = \frac{(2D_{B})^{3}}{2m_{B}=2} \frac{2}{12} \frac{B_{B}=4}{B=2+1} + \frac{4^{2}}{15(B=2+1)^{3}}$$

$$= \frac{D_{B}^{3}}{2m_{B}} \frac{2}{3} \frac{B_{B}+2}{B+2} + \frac{32^{2}}{15(B+2)^{3}} = \frac{E_{B}}{L};$$
(3.8)

The elective ground state energies are exactly the same through the relation (3.7) in the strong coupling case. We can easily verify that other physical quantities are also sm oothly connected through the relation (3.7) at the point 1 = F = 1 = B = 0. In fact, in the case being considered, equations (2.3) and (3.1) can be treated in a uni ed way, if we regard as a parameter and D as a normalization condition for these integral equations. We perform such calculations in Appendix B.

W e note that the quasi-m om enta for bosons and ferm ions are related as

$$K = 2Q; k = 2q:$$
 (3.9)

These relations are consistent with the assumptions that lead us to the G audin integral equation. For negative c, quasimom entum fkg of the fermions (not to be confused with k for the repulsive bosons in (3.9)) become imaginary. In the therm odynamic limit, two k s with a real part q, which are complex conjugates of each other, form a bound state of two fermions with the binding energy $c^2=2$ (see (1.7)).⁹ This kind of argument usually starts with the c = 0 limit. Rem arkably, it holds also in the strong coupling case.

Equations (3.5d) and (3.6d) need some explanation. Since we are seeking for the solution of 0 (1= ³), the expansion must be in powers of either 1= ' 1=(_B + 2) ' 1=(_F + 1=2) or in 1= _B ' 1= _F. However, it occurs that the solution of the rst term should be (_B=(_B + 2))² ' (j_F j=(j_F j 1=2)², not be simply 1. This factor can be counted as the rem iniscent of the e ective \diam eter" a of each particle, in the lim it of the hard-core bose gas, or, in other words,

the quantized version of the classical Tonks gas.¹⁵ In fact, the energy at zero tem perature for the hard-core bose gas is calculated as¹⁹

$$\frac{E_{hc}}{L} = \frac{D_B^3}{2m_B} \frac{2}{3} - \frac{1}{1 - aD_B}^2$$
(3.10)

Besides, equating the scattering phase shifts in the low energy lim it, we nd²⁰

$$a = \frac{1}{2m_B} \frac{2}{c} = \frac{1}{2m_F} \frac{1}{c}$$
 (3.11)

W ith the relation (3.11), we see that (3.10) and the nst terms of (3.6d) and (3.5d) become exactly the same. The positiveness (negativeness) of the diameter a for the attractive fermions (repulsive bosons) is due to its statistical property. The negativeness for the repulsive bosons can also be interpreted as the transparency e ect for a nite c > 0. It is easy to verify that the chem ical potential $_{hc}$ and the sound velocity v_{hc} of the hard-core bose gas¹⁹ are obtained by the substitution of (3.11) with (3.7) into either (3.5e) and (3.5f) or (3.6e) and (3.6f). D iscarding the second term in the last brackets, we obtain

$$_{\rm hc} = \frac{D_{\rm B}^{2}}{2m_{\rm B}} \frac{^{2}}{3} \frac{3}{1} \frac{aD_{\rm B}}{aD_{\rm B}} \frac{1}{1} \frac{^{2}}{aD_{\rm B}} ; \qquad (3.12)$$

$$v_{hc} = \frac{2 D_B}{2m_B} \frac{1}{1 aD_B}^2$$
 (3.13)

Note that h_c corresponds to 2 $\frac{e}{F}$.

Now we are ready to describe the BCS-BEC crossover in one-dimension. The discussion above for smooth connections in the strong coupling cases justiles a link between the two systems. In view of the physical quantities, we can map two fermions of opposite spins forming a bound state whose quasimomenta are q ic=2 onto one boson with quasimomentum k = 2q. In such a way, the elective ground state energy per volume, the chemical potential and the velocity of sound can be identied. If we take $_{\rm F}$ as the coupling constant common to the both system s and vary $1=_{\rm F}$ from 1 to +1 through 0, we not the following asymptotic expressions of the elective energy:

$$\frac{E_{F}^{e}}{N_{FF}} = \bigwedge_{F}^{0} \frac{1}{3} + \frac{6}{2_{F}} ; \qquad 1 = F \quad 1;$$

$$\frac{1}{3} - \frac{F}{2_{F} + 1} + \frac{4^{2}}{15(2_{F} + 1)^{3}} ; 1 = F \quad 0; \qquad (3.14)$$

$$\frac{F}{1} + \frac{1}{3} + \frac{1}{3} + \frac{1}{5} + \frac{$$

where FF is the Fermi energy of the free ferm ions,

$$_{\rm FF} = \frac{1}{2m_{\rm F}} - \frac{D_{\rm F}}{2}^2$$
: (3.15)

The meanings of the above three expressions are clear. They are the energy of the Hartree-Fock approximation for weakly interacting fermions,¹⁶ the energy of either the hard-core bose gas or the spinless non-interacting fermions,²⁶ and the energy due to the Bogoliubov theory for weakly interacting bosons,¹ for 1 = F 1, 1 = F 0, and 1 = F 1, respectively. The asymptotic solutions of the LL and the G audin integral equation reproduce the results of these three \phases" rigorously. Two crossovers for the intermediate value of F are by no means a phase transition. The solutions are continuous and analytic functions of 1 = F except the two points 1 and +1. Those can be proved mathematically from the properties of the integral equations.



Figure 2: A symptotics of the ground state energy E_F^e in units of the Ferm i energy of the free ferm ions as a functions of the inverse of the coupling constant, 1 = F.



Figure 3: Three regimes of one-dimensional quantum gas: for $_{\rm F}$ < 0, spin-1=2 ferm i gas and for $_{\rm F}$ = $_{\rm B}$ =4 > 0, -function bose gas.

Let us explain these three regimes in detail. When 1 = F 1, the ground state is a BCSlike state with Cooper pairs.¹⁶ The elective size of this bound state is much larger than the average interparticle distance. The existence of the therm odynam ic limit is guaranteed by the Pauli exclusion principle; i.e. there is no bound state of more than two ferm ions.

The interm ediate regime of the hard-core bose gas around 1 = F = 0 should be distinguished from the G irardeau gas, which corresponds only to the limit 1 = F = 0. Indeed, when we compare our result with the numerical one, as in g. 2, it traces well the sudden change around 1 = F = 0. In this region of the bosonic side, the strong mutual repulsion compels the bosonic particles to exhibit the ferm ionic properties due to one-dimensional kinematics. On the other hand, even the in nitely strong attraction cannot causes the ferm ions to condense. Namely, the distribution and the energy spectrum of the tightly bound ferm ion pairs can be identied with those of spinless hard-core bosons with a positive \diameter".

In the regime of weakly interacting bose gas, the lowest order solution of the distribution $(k) \mbox{ is}^{1,10}$

$$(k) = \frac{1}{2 D_{F} F} D_{F}^{2} F q^{1=2} : \qquad (3.16)$$

This indicates that the weaker the interaction, the narrower the quasi-m om entum distribution region and the higher the peak of the distribution. In the recent experiment this behavior was observed for $_{\rm B}$ = 0.5.²¹ In the limit 1= $_{\rm F}$! +1, all the particles condense into the zero quasi-m om entum state.

In the same m anner as we have done in the e ective ground state energy, we list here the asymptotic expressions of the velocity of sound and the e ective chem ical potential:

$$\frac{v_{\rm F}}{v_{\rm FF}} = \begin{pmatrix} 2 & \frac{F}{2_{\rm F}+1} & 1 + \frac{2^{2}}{3(2_{\rm F}+1)^{3}} & 1 = F & 1; \\ \frac{1}{2_{\rm F}} & \frac{1}{2_{\rm F}} & 1 & \frac{1}{2_{\rm F}} & \frac{1}{2_{\rm F}} & 1 = F & 0; \\ \frac{1}{2_{\rm F}} & \frac{1}{2_{\rm F}} & 1 & \frac{1}{2_{\rm F}} & \frac{1}{2_{\rm F}} & 1 = F & 1: \\ \end{pmatrix}$$
(3.17)

where v_{FF} is the Ferm i velocity of the free ferm ions,

$$v_{\rm FF} = \frac{D_{\rm F}}{2m_{\rm F}} :$$
 (3.18)

and

$$\frac{e}{F}_{FF} = \begin{cases}
\frac{8}{F} & 1 + \frac{4}{2}_{F}; \\
\frac{1}{32} & F + 1 \\
\frac{2}{2} & F + 1 \\
\frac{2}{2} & F + 1 \\
\frac{2}{5} & \frac{1}{5} & \frac{2}{5} & \frac{1}{5} \\
\frac{2}{5} & F + 1 \\
\frac{2}{5} & \frac{1}{5} & \frac{2}{5} & \frac{1}{5} \\
\frac{2}{5} & F + 1 \\
\frac{2}{5} & \frac{1}{5} & \frac{2}{5} & \frac{1}{5} \\
\frac{2}{5} & F + 1 \\
\frac{2}{5} & \frac{1}{5} & \frac{2}{5} & \frac{1}{5} \\
\frac{2}{5} & F + 1 \\
\frac{2}{5} & \frac{1}{5} & \frac{2}{5} & \frac{1}{5} \\
\frac{2}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} \\
\frac{2}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} \\
\frac{2}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} \\
\frac{2}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} \\
\frac{2}{5} & \frac{1}{5} & \frac{1}{5} \\$$

These expressions are plotted in g. 4 and g. 5 as functions of 1 = F.

4 A nalogy with the C ircular P late C ondenser

An intriguing fact is that the G audin integral equation and the LL integral equation also appear in a completely di erent subject, the circular plate condenser.^{5,7,22} W e consider the condenser of the radius 1 whose two plates are separated by a distance . Cylindrical coordinates are taken as follows: the radial distance from the common axis of the disks is taken to be , the distances from the disks to be , ⁰, respectively. E.R. Love²² (1949) proved that the potential (;; ;⁰) due to the disks is expressed as

$$(;;)^{0} = \frac{V_{0}}{1} \sum_{1}^{Z_{1}} \left(\frac{1}{\left[\frac{2}{2} + (\cdot + it)^{2}\right]^{1-2}} - \frac{1}{\left[\frac{2}{2} + (\cdot 0)^{2}\right]^{1-2}} \right) h(t) dt; \qquad (4.1)$$



Figure 4: A symptotics of the sound velocity v_F in units of the Ferm i velocity of the free ferm ions v_{FF} as a function of the inverse of the coupling constant, 1 = F.



F igure 5: A sym ptotics of the chem ical potential $_{\rm F}^{\rm e}$ in units of the Ferm ienergy of the free ferm ions $_{\rm FF}$ as a function of the inverse of the coupling constant, 1= $_{\rm F}$.

where h (t) is determined by

h (t)
$$-\frac{\sum_{1}^{2} h(s) ds}{\frac{1}{(t - s)^{2} + 2}} = 1$$
: (4.2)

The capacitance C of the condenser is given by

$$C = \frac{1}{2} \int_{-1}^{Z} h(t) dt; \qquad (4.3)$$

In the above expressions, the upper signs refer to the case of equally charged disks at potential V_0 , and the lower signs to that of oppositely charged disks, namely, disks at potential V_0 and

 V_0 . The Love equation (4.2) is exactly in the same form as the G audin and the LL integral equation: to be precise, h(x) in (4.2) corresponds to (1=2)F (x) and 2 g(x) in (2.3) and (3.1), respectively.

In the case of oppositely charged disks (lower signs), the solution of (4.2) is calculated for small (sm all separation). By the heuristic argum $ent_r^{23,24}$ away from the edges h(t) to the o(1) order is written as

$$h(t) = \frac{1}{2} (1 + t)^{1-2} + \frac{(1 + t)^{1-2}}{2} \log \frac{16 + e}{1 + t} + o(1); \qquad (4.4)$$

and with (4.3), the capacitance is

$$C = \frac{1}{4} + \frac{1}{4} \log \frac{16}{e} + o(1):$$
(4.5)

This result is originally by G.K irchho (1877).²⁵

On the other hand, we have 10,20

$$g(\mathbf{x}) = \frac{1}{2} (1 \quad x^{2})^{1=2} \quad \frac{1}{4} \frac{x^{4}}{\sum_{n=0 \text{ m}=0}^{\infty} \frac{(2m+1)!!}{[(2m)!!]^{2}} (2n)!!} (2n \quad 1)!! (2n \quad 1)!!}{(2n \quad 2m \quad 1)} x^{2n}; \quad (4.6)$$

and

$$C = \int_{1}^{Z_{1}} g(x) dx = \frac{1}{4} - \frac{1}{2} + \frac{1}{4}b; \text{ where } b = \int_{m=0}^{X^{2}} \frac{(2m-1)!!}{(2m)!!}^{2} :$$
(4.7)

W e regularize the divergent sum b as follows. W e set

b(k)
$$\frac{X^{4}}{m=0} \frac{(2m-1)!!}{(2m)!!} k^{2m} = \frac{2}{-K} (k):$$
 (4.8)

The function K (k) is the complete elliptic integral of the rst kind with the modulus k. A sym ptotic behavior of K (k) is

K (k) =
$$\frac{1}{2}\log\frac{16}{1-k^2}$$
 as $k^2 \approx 1$: (4.9)

If we choose 1 $k^2 = = (e)$ in (4.9) with (4.8), we get

$$b = \lim_{k^2 \le 1} b(k) = -\frac{1}{2} \log \frac{16 e}{16} :$$
 (4.10)

Using the expression (4.10), we may rewrite (4.6) as

$$g(\mathbf{x}) = \frac{1}{2} (1 \quad \mathbf{x}^{2})^{1=2} + \frac{1}{4} \frac{\mathbf{x}^{1}}{\mathbf{x}^{1}} \frac{\mathbf{x}^{1}}{(2m)!!} \frac{\mathbf{x}^{2}}{(2m)!!!} \frac{(2m-1)!!}{(2m)!!!} \frac{(2m-1)!!}{(2m)!!!} \mathbf{x}^{21} + \frac{1}{4} \frac{\mathbf{x}^{1}}{\mathbf{x}^{1}} \frac{\mathbf{x}^{1}}{(2m)!!!} \frac{(2m-1)!!}{(2m)!!!} \frac{(2m-1)!!}{(2m)!!!} \frac{(2m-1)!!}{2m} \mathbf{x}^{21} = \frac{1}{2} (1 \quad \mathbf{x}^{2})^{1=2} + \frac{1}{4} \frac{1}{2} \log \frac{16}{2} \frac{e}{\mathbf{x}} \log \frac{1+\mathbf{x}}{1-\mathbf{x}} \frac{1}{(1-\mathbf{x}^{2})^{1=2}}$$
(4.11)

Here, we have used the power series form ulae,

$$\sum_{p=0}^{X} \frac{(2p \ 1)!!}{(2p)!!} x^{2p} = (1 \ x^2)^{1=2}; \quad jxj < 1;$$
(4.12)

$$\overset{X^{1}}{\underset{n=0}{\sum}} \frac{1}{2n+1} x^{2n+1} = \frac{1}{2} \log \frac{1+x}{1-x}; \quad jx j < 1;$$
 (4.13)

and the following sum mation form ula,

$$\frac{X^{1}}{m=0} \frac{(2m \ 1)!!}{(2m)!!} \frac{2}{2m} \frac{1}{(2l+1)} = \frac{2}{(2l)!!} \frac{(2l)!!}{(2l+1)!!} \frac{X^{1}}{2l} \frac{1}{2l} \frac{1}{2k+1} \frac{(2k \ 1)!!}{(2k)!!};$$

$$1 = 0;1;2;:::; (4.14)$$

which is proved in Appendix C.

Therefore, by the regularization of the divergent sum (4.10), we exactly reproduce Hutson's solution (4.4) for x = 0 for o(1). This coincidence is reasonable since we have started from the expansion of g(x) around x = 0.

We may also calculate the physical quantities from (4.11). At our consideration, to the ⁰ order, we can integrate (4.11) for the whole interval [1;1]. Indeed, the corrections come from near the edges, x 2 [1; 1+][[1 p;1], and g(x) is estimated to be of the order 1= there. Then, the corrections are of the order _____, and thus can be neglected.

The case of equally charged disks (upper signs in (4.1) and (4.2)) has not attracted much interests. So far, no prescription has been proposed to solve (4.2) asymptotically. Our solution (2.31) gives $_7$

$$C = \frac{1}{4} \int_{-1}^{2} F(x) dx = \frac{1}{2} + \frac{1}{4^2} \log \frac{4}{2} + 0 (); \qquad (4.15)$$

To obtain the last expression, the regularization of the divergent sum b in the same manner as (4.10) is required:

$$b = \sum_{n=0}^{X^{1}} \frac{1}{2m+1} = \lim_{k^{2} \leqslant 1} \frac{1}{2m+1} k^{2m} = \frac{1}{2} \log \frac{4 e}{2m};$$
(4.16)

where (4.13) has been used to obtain the last expression. The rst term in (4.15) was also referred to in the Kirchho 's paper.²⁵ The second term agrees with the analysis by Gaudin⁷ except 4 in the logarithm ic function.

5 Concluding Remarks

W e have presented solutions of the G audin integral equation (1.5). W e have solved the integral equation in the form of power series. B ased on the solution, the strong coupling case is analyzed, in relation to the Lieb-Liniger integral equation for the bosons with repulsive interaction. Then, we have a uni ed view on -function attractive ferm ions and repulsive bosons in one-dimension. M aim results are sum m arized as follow s.

- i) As to the therm odynam ic quantities, the expressions in the strong coupling cases j j = 1 are identical through the relations (3.7).
- ii) The lim it 1 = 0 yields the G irandeau gas,²⁶ in the sense that the energy spectra coincide with that of free ferm ions.
- iii) A round the point 1 = 0, the system behaves like a hard-core bose gas with a diam eter a given by (3.11). This indicates the existence of dim ers of tightly bound ferm ions (1 = 0, a > 0), which can be regarded as bosons with strong repulsion (1 = +0, a < 0).

 iv) The above regime of can be smoothly connected to the weakly attractive ferm igas and the weakly repulsive bose gas at both ends with su ciently smallvalue of j j. This re ects the analyticity of the Love equation (4.2) in on the whole real axis excluding the point = 0.

In the weak coupling case, we have again met the di culty of divergences. O bviously, the solution (2.31) holds only for x 2 [1+ ;1] (see AppendixA). Near the edges jxj 1, a di erent method might be necessary to determ ine the precise behavior of the solution. Further, the relation between and again is in portant, since the regularization of by yields the nite expressions of the physical quantities. Up to now, our method is powerful enough to obtain the rst two terms of the asymptotic expressions. We believe that other physical quantities, such as the dressed charge and the dressed energy, and other quantum integrable systems can be studied by use of this method. Those subjects remain as future problem s.

Appendices

A Another D erivation of (2.24) and (2.28)

In this appendix we present the expansion of the integral,

$$J(x;) - \int_{-1}^{Z} \frac{1}{2} dy \frac{F(y)}{2 + (x - y)^{2}} = - \int_{-1}^{Z} \frac{1}{2} dy \frac{m = 0}{2 + (x - y)^{2}}; \quad (A.1)$$

in power series of for the case 1. Since

$$\frac{1}{x^{2} + (x - y)^{2}} = \underbrace{\begin{cases} \frac{1}{(x - y)^{2}} \frac{1}{1 + \frac{x - y^{2}}{1 + \frac{x - y^{2}}{$$

for a xed x 2 [1+ ;1], A 1) becomes

$$J(x;) = - \begin{array}{c} x \\ z \\ (1;x) \\ (1;x$$

In the above, $\int_{k=0}^{2^{n}} m$ eans that we exclude the k = 2n + 1 terms in the sum with respect to k.

Let us rewrite (A 3) so that we can easily see the dependences of and x. For the purpose, we change the variables and the orders of the multiple sum s. To begin with, we consider the rst term of the rhs. of (A 3). The order of the triple series can be changed as

W ith a new variable p = m linstead of m, the rst term of the r.h.s. of A 3) can be written as

$$\frac{2}{2} \frac{x^{1}}{2} \frac{x^{1}}{2} \frac{x^{n}}{21} \frac{x^{n}}{21} \frac{2m}{21} \frac{(1)^{n} a_{m} x^{2(m-1)}}{21 21 2n 1} \\
= \frac{2}{2} \frac{x^{1}}{21} \frac{x^{1}}{21} \frac{21+2p}{21} \frac{x^{1}}{a_{1+p}} \frac{(1)^{n}}{21 2n 1} x^{2p} : (A.5)$$

In exactly the same way, the fourth term becomes

The third term is rewritten as follows:

$$\frac{1}{2} \begin{bmatrix} X^{k} & ^{n}X^{-1} & 2n+1 & 2m \\ & & 2n+1 & 2n+1 & 2n+1 \\ & & & 2n+1 & 2n+1 \\ & & & & & & \\ \end{bmatrix} \begin{bmatrix} X^{k} & 2n+1 & 2n+1 & 2n+1 \\ & & & & & \\ \end{bmatrix} \begin{bmatrix} X^{k} & X^{k} & 2m & 1 \\ & & & & & \\ 2n+1 & 2n+1 & 2n+1 & 2n+1 \\ & & & & \\ \end{bmatrix} \begin{bmatrix} x^{k} & X^{q} & 21+2n \\ & & & & \\ \end{bmatrix} \begin{bmatrix} x^{k} & X^{q} & 21+2n \\ & & & & \\ x^{k} & X^{q} & 21+2n \\ & & & & \\ x^{n+1} & 2n+1 & 2q & 21+1 \\ \end{bmatrix} \begin{bmatrix} x^{2q} & x^{2q} & x^{2q} \\ x^{2q} & x^{2q} \\ x^{2q} & x^{2q} \end{bmatrix}$$

The second term is rather complicated. Since

we decompose the sum with respect to k into four parts:

$$\hat{X}^{m} \circ X^{n} \hat{X}^{m} X^{n} X^{n} X^{1} X^{n} X^{n} X^{1}$$

$$= + = + + + :$$

$$k=0 \quad k=2n+2 \quad j=0 \quad j=0 \quad j=n+1 \quad j=n+1 \\ k=2j \quad k=2j+1 \quad k=2j \quad k=2j+1$$
(A.9)

For a xed n 0, the sum with respect to m can be decomposed into two parts:

$$X^{\hat{d}} = X^{n} X^{\hat{d}}$$

= + : (A.10)
 $M = 0 \quad M = 0 \quad M = n + 1$

W ith (A.9) and (A.10), the double series with respect to m and k become

Now we proceed to evaluate the second term of the r.h.s. of (A.3) using (A.9) and (A.11):

To obtain the last expression from the middle, we have again changed the variables and the orders of the multiple sum s. For instance, in the rst square bracket [], we have replaced m; j

and p w ith q = m, l = m j, r = m j + p, and used the relation

to collect the term s of 21 and 21 1.

1 1

The combination of (A .5), (A .12), (A .7) and (A .6) yields the following form of (A .3):

$$J(x;) = \frac{2}{n} \sum_{n=0}^{N} [A_n(x) + B_n(x)] + \frac{2}{n} \sum_{n=0}^{N} (1)^n \sum_{n=0}^{2n+1} [C_n(x) + D_n(x)] + E_n(x) + F_n(x)];$$
(A.14)

where A_n (x), B_n (x), C_n (x), D_n (x), E_n (x) and F_n (x) are de ned as

$$A_{n}(\mathbf{x}) = \frac{\sum_{p=0}^{\infty} \sum_{q=0}^{\infty} 2n + 2p}{2n} \frac{(1)^{p} a_{n+p}}{2q - 2n + 1} \mathbf{x}^{2p};$$
(A.15)

$$B_{n}(\mathbf{x}) = \sum_{p=0 q=0}^{\mathbf{X}^{d}} \frac{\mathbf{X}^{d}}{2n} \frac{2n+2p}{2n} \frac{(1)^{p}a_{n+p}}{2q+2n+1} \mathbf{x}^{2p};$$
(A.16)

$$C_{n}(\mathbf{x}) = \begin{pmatrix} X^{n} & X^{p} & X^{2q} \\ & a_{q}A_{npqr}x^{2p} + \\ & p=0 & q=0 r=0 \\ & q=p+1r=0 \\ & p=n+1q=0 r=0 \\ & (A.17) \end{pmatrix}$$

$$D_{n}(\mathbf{x}) = \frac{\overset{1}{\mathbf{x}} \overset{1}{\mathbf{x}} \overset{1}{\mathbf{x}}^{2q}}{a_{q}A_{npqr}\mathbf{x}^{2p}} + \frac{\overset{n}{\mathbf{x}}^{p} \overset{1}{\mathbf{x}}^{2p}}{a_{q}A_{npqr}\mathbf{x}^{2p}} + \overset{1}{\overset{}} \overset{\mathbf{x}}{a_{q}A_{npqr}\mathbf{x}^{2p}} + \overset{1}{\overset{}} \overset{\mathbf{x}}{a_{n+p}A_{np;p+n;2p}\mathbf{x}^{2p}};$$

$$(A.18)$$

$$E_{n}(x) = a_{q}A_{npqr}x^{2p};$$
 (A.19)

$$F_{n}(x) = \frac{\overset{x}{k} \overset{x}{x^{p}} 2q + 2n}{\underset{p=1 q=1}{\overset{2n+1}{2}} 2n + 1} \frac{a_{n+q}}{2q} \overset{x^{2p}}{2p} :$$
(A 20)

Here, the sequence

$$A_{npqr}$$
 (1) $\begin{pmatrix} 2q & 2q & 2n & 1 & r \\ r & 2p & r & 2q & 2n & 1 & r \\ r & 2p & r & 2q & 2n & 1 & r \\ \end{pmatrix}$; n;p;q;r2Z₀; (A.21)

has been introduced to simplify the expressions. Note that sum s in the above whose lower bounds are greater than its upper bounds, such as $P^{p}_{q=p+1}$, are treated as zero.

From (A.14), we obtain the expansion of J(x;) with respect to as follows:

$$J(x;) = \int_{n=0}^{X^{d}} J_{n}(x); \qquad (A 22)$$

where the st few J_n are

$$J_{0}(x) = \frac{2}{-} [A_{0}(x) + B_{0}(x)]; \qquad (A 23)$$

$$J_{1}(x) = \frac{2}{2} [C_{0}(x) + D_{0}(x) + E_{0}(x) + F_{0}(x)]; \qquad (A 24)$$

$$J_{2}(x) = \frac{2}{-} [A_{1}(x) + B_{1}(x)]; \qquad (A 25)$$

$$J_{3}(x) = - [C_{1}(x) + D_{1}(x) + E_{1}(x) + F_{1}(x)]:$$
 (A 26)

From (A 23), (A 25) and

$$\sum_{n=0}^{X^{1}} \frac{(1)^{n}}{2n+1} = \frac{1}{4};$$
 (A 27)

we easily verify that

$$J_{0}(x) = \int_{p=0}^{X^{2}} a_{p} x^{2p} = F(x); \qquad (A 28)$$

$$J_{2}(\mathbf{x}) = \bigvee_{p=0}^{X^{1}} (p+1)(2p+1)a_{p+1}x^{2p}:$$
 (A.29)

U se of (A 28) in the G audin integral equation (2.3) leads to (2.24). To simplify the expressions of J_1 (x) and J_3 (x), we need a little calculation and some form ulae. The results are,

$$J_{1}(x) = \frac{2}{n} \frac{x^{d}}{\sum_{p=0}^{n=0} \frac{x^{d}}{2n}} \frac{2p+1}{(2p+1)} a_{n}^{"} x^{2p}; \qquad (A.30)$$

$$J_{3}(\mathbf{x}) = \frac{2}{p} \frac{\mathbf{x}}{2p} + 3 \qquad \mathbf{x}^{2} \frac{\mathbf{x}}{2p} + 3 \qquad \mathbf{x}^{2} \frac{\mathbf{a}_{n}}{2n} \frac{\mathbf{a}_{n}}{(2p+3)} \mathbf{x}^{2p} : \qquad (A 31)$$

Below, we shall explain these shortly. We observe that equation (A.30) is in accordance with (2.28).

Therefore, we conclude that our form alexpansion (2.7) is also valid for small in the interval [1 + ; 1].

Calculation of $J_1(x)$

From (A 24) with (A 17) { (A 21), we have

$$\frac{1}{2}J_{1}(\mathbf{x}) = \frac{x^{i}}{2} a_{0}A_{0p00}x^{2p} + \frac{x^{i}}{2}x^{p}}{a_{q}A_{0pq/2q}x^{2p} + x^{i}} \frac{x^{i}}{2}x^{k} \frac{x^{p}}{x^{p}} a_{q}A_{0pqr}x^{2p} + \frac{x^{i}}{2}x^{k} \frac{x^{p}}{2q} a_{q}A_{0pqr}x^{2p} + \frac{x^{i}}{2}x^{2p} \frac{x^{2p}}{2q} \frac{1}{2q} \frac$$

Here, to obtain the third expression from the second, the following formula has been used:

Proof (A.33)

Set

$$\frac{P_{p}(m)}{m p} = \begin{pmatrix} X^{p} \\ (1)^{l} & m+1 \\ 1 & p \\ l & n \end{pmatrix} = \begin{pmatrix} X^{p} \\ (1)^{l} & m+1 \\ 1 & p \\ l & m \end{pmatrix} = \begin{pmatrix} X^{p} \\ (X,34) \\ ($$

It can be seen that

$$P_p$$
 (m) is a polynom ial of m whose degree is equal to or less than p. (A 35)

It will be shown that $P_{\rm p}\ (\!m$) is a constant, and actually,

$$P_{p}(m) = (1)^{p}(p+1)$$
: (A.36)

W hen p = 0 or p = 1, we can verify it directly, so we assume that p = 2.

1

First, let us denote the 1-th term of the sum $p_{l=0}$ in the r.h.s. of (A.34) as $Q_1(m) = (m p)$, namely,

$$P_{p} (m) = \sum_{l=0}^{X^{p}} Q_{l} (m):$$
 (A.37)

Then, Q_1 (m) is explicitly written as

$$Q_{1}(m) = (1)^{1} \frac{m+1}{1} \frac{m}{p} \frac{1}{m} \frac{m}{p}$$

$$\stackrel{8}{\underbrace{(m \ 1)(m \ 2)}{p!}} \frac{m}{p!} \frac{m}{p!} \quad \text{if } l = 0,$$

$$= (1)^{1} \frac{(m+1)m}{p!} \frac{(m \ 1+1)m}{1!(p \ 1)!} \frac{m}{p!}; \quad \text{if } l = p,$$

$$\stackrel{(1)^{p} \frac{(m+1)m}{p!}}{p!} \frac{(m \ p+2)}{p!}; \quad \text{if } l = p.$$

Here, () denotes that such term is excluded from the sequence of the product. From (A.37) and (A.38), $P_p(0)$; $P_p(1)$; ::: $P_p(p)$ are calculated as follows:

$$P_{p}(k) = \begin{pmatrix} Q_{k}(k) + Q_{k+1}(k); & \text{if } 0 & k & p & 1, \\ Q_{p}(p); & \text{if } k = p, \\ = (1)^{p}(p+1): \quad (A.39)$$

Therefore, (A 39) with (A 35) assures that $P_p(m) = (1)^p(p+1)$ for all m, which completes the proof.

Calculation of $J_3(x)$

This could be done in almost the same, but more complicated way than $J_1(x)$. We sketch

the calculation. From (A 26) with (A .17) { (A 21), we have

$$\frac{1}{2}J_{3}(x) = a_{0} \bigwedge_{p=0}^{X} A_{1p00}x^{2p} + a_{1}A_{1010}x^{0} + a_{1} \bigwedge_{p=1r=0}^{X} A_{1p1r}x^{2p}$$

$$+ \bigwedge_{p=2q=2r=2q}^{X} A_{q}A_{1pqr}x^{2p} + \bigwedge_{p=1}^{X} 2p+2 \prod_{q=1}^{1} a_{p+1}x^{2p}$$

$$+ \bigwedge_{p=1q=2}^{X} X^{p} (1)^{p} 2q (2p+3)^{2p}$$

$$+ \bigwedge_{p=0q=p+2}^{X} X^{2} (1)^{p} 2q (2q)^{3} r^{r} \frac{1}{2q} 3 r^{2p}$$

$$= a_{0} \sum_{p=0}^{X} 2p+2 \prod_{r=0}^{X} x^{2p} + a_{1} \sum_{p=0}^{X} 2p+3 \prod_{q=0}^{1} \frac{1}{2(2p+1)}x^{2p}$$

$$+ \frac{\chi^{q} \chi^{q}}{p=1q=2} 3 \frac{2q}{2(2p+3)}x^{2p} + \chi^{q} \chi^{q} 2p+3 \prod_{q=0q=p+2}^{q} \frac{1}{2(2p+1)}x^{2p}$$

$$= \frac{\chi^{q} \chi^{q}}{p=1q=2} 3 \frac{2q}{2(2p+3)}x^{2p} + \chi^{q} \chi^{q} \chi^{q} 2p+3 \prod_{p=0q=p+2}^{q} \frac{1}{2(2p+3)}x^{2p}$$

$$= \frac{\chi^{q} \chi^{q}}{q} 2p+3 \prod_{q=0q=p+2}^{2} \frac{1}{3} \frac{2q}{2q} (2p+3)}{q} \chi^{2p} : \qquad (A.40)$$

In the above, the second expression is obtained from the st by applying the identity,

for the sum with respect to r in the last term . We note that (A .41) can be proved in a similar way as (A .33).

B Solutions of the Lieb-Liniger and the G audin integral equations around 1 = 0

In this appendix, we solve an integral equation,

H (x;)
$$-\frac{\sum_{1}^{2} \frac{H(y;) dy}{1 + \frac{2}{2} (x - y)^{2}} = \frac{1}{2};$$
 (B.1)

around = 0. We remark that, H (x;) gives the solutions F (x) of (2.3) and g(x) of (3.1) in the strong coupling case (i.e. j = j = 1),

$$F(x) = 4 H(x; 1=);$$
 (B.2)

$$g(x) = H(x; 1=):$$
 (B.3)

The physical quantities are expressed as follows, for 0,

$$F = 4 \qquad H (x;) dx; ; (B.4a)$$

$$Q = D_F 4 H(x;) dx;$$
 (B.4b)

$$\frac{E_{F}^{e}}{L} = \frac{D_{F}^{3}}{16} \begin{bmatrix} Z_{1} & & & & \\ & & x^{2}H (x;) dx & & H (x;) dx \end{bmatrix} (B.4c)$$

for 0,

$$\vec{Z}_{1} \qquad 1$$

$$K = D_{B} \qquad H(x;) dx ; \qquad (B.5b)$$

$$\frac{E_B}{L} = D_B^3 \xrightarrow{X^2H} (x;) dx \xrightarrow{I} H (x;) dx : (B.5c)$$

Hereafter, we denote H $(\!x\!\!\!;\,)$ as H $(\!x\!)$ for simplicity. W ith the expansions of H $(\!x\!)$ and the kernel,

H (x) =
$$\sum_{p=0}^{X^{4}} b_{p} x^{2p}$$
; (B.6)

$$\frac{1}{1+\frac{2}{3}} \left(x - y\right)^{2} = \int_{n=0}^{x^{2}} (-1)^{n} (x - y)^{2n-2n}; \qquad (B.7)$$

equation (B.1) becom es

which, as in x 2, reduces to the following set of algebraic equations for fb_pg :

$$b_{p} = \frac{2}{m} \sum_{m=0}^{M} \frac{2}{n-p} \left(1\right)^{n-2n+1} \frac{b_{m}}{2m+2n-2p+1} \frac{2n}{2n-2p} = \frac{1}{2} \sum_{p_{0}} (B.9)$$

From (B.9), we get

$$b_{0} = \frac{1}{2} + \frac{2}{2} + \frac{4}{2} + \frac{2}{2} + \frac{2}{4} + \frac{4}{2} + \frac{1}{3} + \frac{3}{3}$$

$$\frac{4}{2} + \frac{4}{2} + \frac{4}{2} + \frac{2}{3} + \frac{20}{3^{2}} + \frac{1}{5} + \frac{16}{4} + \frac{5}{5} + 0 + 0 + \frac{6}{3}; \quad (B.10)$$

$$b_1 = \frac{1}{2} \cdot 3 \cdot 1 + \frac{2}{2} \cdot 2 \cdot 1 \cdot \frac{2}{2} \cdot 2 + O(^6);$$
 (B.11)

$$b_2 = \frac{1}{2} + O(6); \qquad (B.12)$$

$$b_p = O(2^{p+1}); p 3:$$
 (B.13)

By use of the above solution, may be expressed as

$$= +\frac{4}{3} \frac{3}{4} \frac{32}{15} \frac{5}{5} + 0 (7); \qquad (B.14)$$

where

$$= \frac{1}{{}_{\rm B}+2} = \frac{1}{4_{\rm F}+2}:$$
 (B 15)

In term s of , we have

$$b_0 = \frac{1}{2(1 2)} 1 \frac{2}{3} + 0(^6); \qquad (B 16a)$$

$$b_{1} = \frac{3}{1 - 2} + O(6); \qquad (B.16b)$$

$$b_2 = {}^{3} {}^{5} + O({}^{6}):$$
 (B.16c)

It follows from the above relations that for $0_{,}$

$$K = D_{B} (1 \ 2 \) \ 1 + \frac{4}{3} \ ^{3} \ ^{3} + O(\ ^{5}); \tag{B.17}$$

$$\frac{E_B}{L} = D_B^3 \frac{2}{3} (1 \quad 2 \quad 2 \quad 1 + \frac{32}{15} \quad 2 \quad 3 \quad + O(6):$$
 (B.18)

O ther physical quantities are readily obtained, which reproduce (3.5) and (3.6).

C Proof of (4.14)

We begin with the following expression of the complete elliptic integral of the rst kind,

$$K (k) = \int_{0}^{\frac{2}{2}} \frac{d}{1 + k^{2} \sin^{2}} = \frac{1}{2} \int_{m=0}^{\frac{1}{2}} \frac{(2m - 1)!!}{(2m)!!} k^{2m}; \quad j \le 1:$$
(C.1)

If we consider the expansion such that

$$\frac{2}{\pi} \frac{dt}{dt} \frac{dt}{t^{2l+2}} \int_{0}^{\frac{\pi}{2}} \frac{d}{p \frac{d}{1 + t^{2} \sin^{2}}} = \int_{n=-1}^{\frac{M}{2}} c_{n} x^{n}; \text{ for } x > 0, \qquad (C 2)$$

we see from (C .1) that

$$c_{0} = \frac{X^{2}}{m} \frac{(2m \ 1)!!}{(2m \)!!} \frac{2}{2m} \frac{1}{2m} \frac{1}{2m} (C \ 3)$$

In the lhs. of (C 2), since the double de nite integral is convergent for x > 0, we can invert the order of the integrations. To integrate with respect to t, we change the variable from t to ', where ' is de ned as ' = arcsin(tsin); 0 ' =2. Then the integration in the lhs. of (C 2) becomes

$$\frac{2}{\pi} \frac{2}{x} \frac{dt}{t^{2l+2}} \frac{dt}{0} \frac{dt}{1 + \frac{2}{t} \sin^2} = \frac{2}{\pi} \frac{2}{t^2} \frac{d}{d} \frac{dt}{dt} \frac{dt}{dt} \frac{\sin^{2l+1}}{\sin^{2l+2} t}; \quad (C.4)$$

By use of the form ula,

$$\frac{Z}{\sin^{2l+2} \prime} = \frac{(2l)!!}{(2l+1)!!} \frac{\cos'}{\sin^{2l+1} \prime} \frac{X^{1}}{k=0} \frac{(2l-2k-1)!!}{(2l-2k)!!} \sin^{2k} \prime; \qquad (C.5)$$

the r.h.s. of (C.4) is rewritten as

Since the second term in the square bracket [] consists of odd powers of x, only the rst term contributes to c_0 , which gives

$$c_{0} = \frac{2}{(21)!!} \frac{(21)!!}{(21+1)!!} \frac{X^{\perp}}{k=0} \frac{(21-2k-1)!!}{(21-2k)!!} \frac{1}{2k+1} :$$
 (C.7)

Therefore, we obtain

$$\frac{X^{1}}{(2m)!!} \frac{(2m-1)!!}{(2m)!!}^{2} \frac{1}{2m} \frac{1}{(2l+1)} = \frac{2}{(2l)!!} \frac{(2l)!!}{(2l+1)!!} \frac{X^{1}}{(2l-2k)!!} \frac{(2l-2k-1)!!}{(2l-2k)!!} \frac{1}{2k+1}$$

$$= \frac{2}{(2l)!!} \frac{(2l)!!}{(2l+1)!!} \frac{X^{1}}{(2l-2k)!} \frac{1}{(2k+1)!!} \frac{(2k-1)!!}{(2k)!!};$$

l= 0;1;2;:::; (C .8)

which is nothing but (4.14).

References

- [1] E.Lieb and W.Linger: Phys. Rev. 130 (1963) 1605.
- [2] E.Lieb: Phys. Rev. 130 (1963) 1616.
- [3] M.Gaudin: Phys. Lett. 24A (1967) 55.
- [4] C.N.Yang: Phys.Rev.Lett.19 (1967) 1312.
- [5] M.Gaudin: Phys.Rev.A 4 (1971) 386.
- [6] D.C.M attis (ed.): The Many-Body Problem : An Encyclopedia of Exactly Solved Models in One D im ension (W orld Scientic, Singapore, 1993).
- [7] M.Gaudin: La Fonction d'Onde de Bethe (The Bethe W ave Function) (M asson S.A., Paris, 1983) [in French].
- [8] V.E.Korepin, N.M.Bogoliubov and A.G.Izergin: Quantum Inverse Scattering Method and Correlation Functions (Cambridge University Press, New York, 1993) Chap.1.
- [9] M. Takahashi: Therm odynam ics of One-D imensional Solvable Models (Cambridge University Press, Cambridge, 1999).
- [10] M.W adati: J.Phys.Soc.Jpn.71 (2002) 2657.
- [11] T. Lida and M. W adati: J. Phys. Soc. Jpn. 72 (2003) 1874.
- [12] M.T.Batchelor, X.W.Guan and J.B.McGuire: J.Phys.A 37 (2004) L497.
- [13] I.V. Tokatly: Phys. Rev. Lett. 93 (2004) 090405-1.
- [14] J.N.Fuchs, A.Recatiand W. Zwerger: Phys. Rev. Lett. 93 (2004) 090408-1.
- [15] L.Tonks: Phys.Rev.50 (1936) 955.
- [16] V.Ya.Krivnov and A.A.Ovchinnikov: Sov. Phys. JETP 40 (1975) 781.
- [17] P. K. Kythe and P. Puri: Computational Methods for Linear Integral Equations (Birkhauser, Boston, 2002).
- [18] M.W. Zwierlein, C.A. Stan, C.H. Schunck, S.M. F. Raupach, S. Gupta, Z. Hadzibabic and W. Ketterle: Phys. Rev. Lett. 91 (2003) 250401-1.
- [19] M.W adatiand G.Kato: Chaos, Solitons & Fractals 14 (2002) 23.
- [20] M. Wadati, G. Kato and T. Iida: Statistical Mechanics of Quantum Integrable Systems, in Nonlinear Waves: Classical And Quantum Aspects, eds. F. Kh. Abdullaev and V. V. Konotop (Kluwer Academic Publishers, 2004) pp. 193.
- [21] B.Pardes, A.W idera, V.Murg, O.M andel, S.Folling, I.Cirac, G.V.Shlyapnikov, T.W. Hansch and I.Bloch: Nature 429 (2004) 277.
- [22] E.R.Love: Quart.J.Mech.Appl.Math.2 (1949) 428.
- [23] V.Hutson: Proc.Camb.Phil.Soc.59 (1963) 211.

- [24] V.N.Popov: Theor.M ath.Phys.30 (1977) 222.
- [25] G.Kirchho : Monatsb.Acad.W iss.Berlin (1877) 144.
- [26] M.Girardeau: J.Math.Phys.1 (1960) 516.