

Effective interaction between molecules in the BEC regime of a superfluid Fermi gas

Y. Ohashi

Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan,

Department of Physics, University of Toronto,

Toronto, Ontario, Canada M5S 1A7

(Dated: March 23, 2022)

Abstract

We investigate the effective interaction between Cooper-pair molecules in the strong-coupling BEC regime of a superfluid Fermi gas with a Feshbach resonance. Our work uses a path integral formulation and a renormalization group (RG) analysis of fluctuations in a single-channel model. We show that a physical cutoff energy ω_c originating from the finite molecular binding energy is the key to understanding the interaction between molecules in the BEC regime. Our work thus clarifies recent results by showing that $a_M = 2a_F$ is a *bare* molecular scattering length while $a_M = (0.6 \sim 0.75)a_F$ is the low energy molecular scattering length renormalized to include high-energy scattering up to ω_c (here a_F is the scattering length between Fermi atoms). We also include many-body effects at finite temperatures. We find that a_M is strongly dependent on temperature, vanishing at T_c , consistent with the earlier Bose gas results of Bijlsma and Stoof.

PACS numbers: 03.75.Ss, 03.75.-b, 03.75.Kk

Recently, the interaction between bound molecules has attracted much attention in the strong-coupling BEC regime of a superfluid Fermi gas[1, 2, 3, 4]. In a trapped Fermi gas with a Feshbach resonance, one can tune the magnitude of a pairing interaction by varying the threshold energy of the Feshbach resonance[5, 6, 7]. Using this tunable interaction, the BCS-BEC crossover[7] has been observed in ^{40}K and ^6Li , where the character of superfluidity continuously changes from the weak-coupling BCS type to the BEC of tightly bound molecules, as one increases the pairing interaction[1, 8]. In the strong-coupling BEC regime, where the molecules already form above the superfluid phase transition temperature T_c [7], the system can be regarded as a molecular Bose gas.

The magnitude of the molecular interaction in the BEC regime was originally studied in the superconductivity literature[1, 2], where the result $a_M = 2a_F$ was obtained using a Ginzburg-Landau (GL) type expansion. Here a_M is the molecular scattering length and a_F is the atomic s -wave two-body scattering length. Later, Pieri and Strinati[3] obtained $a_M \simeq 0.75a_F$ within a two-body t -matrix approximation. More recently, Petrov and co-workers[4] obtained $a_M \simeq 0.6a_F$ by solving a four-fermion problem. This last result is confirmed by a direct Monte-Carlo simulation for the ground state[9]. In addition, Grimm and co-workers also obtained $a_M \simeq 0.6a_F$ in superfluid ^6Li [10], using a Thomas-Fermi fit to their molecular condensate density profile.

In this paper, we investigate this interaction between molecules in the strong-coupling BEC regime of a uniform Fermi gas, using a single-channel model. We show that the finite binding energy of a Cooper-pair “molecule” naturally leads to a *physical* cutoff energy ω_c , which is not present in the usual *atomic* BEC. In a two-molecule case, we show that ω_c is the key to understanding the physics behind the difference between earlier results, namely, $a_M = 2a_F$ [1, 2] and $a_M = (0.6 \sim 0.75)a_F$ [3, 4]. We also calculate the many-body t -matrix at finite temperatures, extending the renormalization group analysis[11] for an atomic BEC to the BEC regime of a superfluid Fermi gas. In a many-particle system, we show that a_M depends strongly on temperature, vanishing in the region near T_c .

We consider a two-component Fermi gas described by the single-channel BCS model. In the functional integral formalism, the action S in the partition function $Z \equiv \sum_{\sigma} \int \mathcal{D}\Psi_{\sigma}^{\dagger} \mathcal{D}\Psi_{\sigma} e^{-S}$ has the form[2], $S = \int_0^{\beta} d\tau \int d\mathbf{r} \left[\sum_{\sigma} \Psi_{\sigma}^{\dagger} \left(\frac{\partial}{\partial \tau} + \frac{\hat{\mathbf{p}}^2}{2m} - \mu \right) \Psi_{\sigma} - U \Psi_{\uparrow}^{\dagger} \Psi_{\downarrow}^{\dagger} \Psi_{\downarrow} \Psi_{\uparrow} \right]$. Here, $\Psi_{\sigma}(\mathbf{r}, \tau)$ and $\Psi_{\sigma}^{\dagger}(\mathbf{r}, \tau)$ are a Grassmann variable and its conjugate, describing Fermi atoms with pseudo-spin $\sigma = \uparrow, \downarrow$. μ is the atomic chemical potential. In ^{40}K and ^6Li Fermi

gases, the tunable pairing interaction U is associated with the Feshbach resonance[5, 6, 7]. We simply treat U as a tunable parameter in a single channel formulation. This is valid for a broad Feshbach resonance.

To discuss the strong-coupling BEC regime, it is convenient to introduce a Cooper-pair Bose field $\Delta(\mathbf{r}, \tau)$, using the usual Stratonovich-Hubbard transformation[2]. After functional integrations over Ψ_σ and Ψ_σ^\dagger , we obtain $Z = \int \mathcal{D}\Delta^\dagger \mathcal{D}\Delta e^{-S_\Delta}$, where $S_\Delta = \int_0^\beta d\tau \int d\mathbf{r} \frac{|\Delta|^2}{U} - \text{Tr}[\ln[-\hat{G}^{-1}]]$ [2]. The fermion single-particle Green's function is given by

$$\hat{G}^{-1} \equiv -\frac{\partial}{\partial \tau} - \left(\frac{\hat{\mathbf{p}}^2}{2m} - \mu\right)\tau_3 + \begin{pmatrix} 0 & \Delta(\mathbf{r}, \tau) \\ \Delta^\dagger(\mathbf{r}, \tau) & 0 \end{pmatrix}, \quad (1)$$

where τ_j ($j = 1, 2, 3$) is the Pauli matrices. The mean-field gap equation is obtained from the saddle point solution for Δ_{MF} determined by $\partial S_\Delta / \partial \Delta = 0$. Expanding the action S_Δ around this mean-field solution (Δ_{MF}), we obtain

$$S_\Delta = \int_0^\beta d\tau \int d\mathbf{r} \frac{|\delta\Delta|^2}{U} + \sum_{n=2}^{\infty} \frac{(-1)^n}{n} \text{Tr} \left[\left[\hat{G}^0 \begin{pmatrix} 0 & \delta\Delta \\ \delta\Delta^\dagger & 0 \end{pmatrix} \right]^n \right], \quad (2)$$

where $\delta\Delta(\mathbf{r}, \tau) \equiv \Delta(\mathbf{r}, \tau) - \Delta_{\text{MF}}$ describes fluctuations in the particle-particle channel. The mean-field 2×2 matrix Green's function \hat{G}^0 has the standard BCS form,

$$\hat{G}^0(\mathbf{p}, i\omega_m) = -\frac{i\omega_m + (\varepsilon_{\mathbf{p}} - \mu)\tau_3 - \Delta_{\text{MF}}\tau_1}{\omega_m^2 + E_{\mathbf{p}}^2}. \quad (3)$$

Here, $i\omega_m$ is the fermion Matsubara frequency. $E_{\mathbf{p}} \equiv \sqrt{(\varepsilon_{\mathbf{p}} - \mu)^2 + \Delta_{\text{MF}}^2}$ is the single-particle excitation spectrum, where $\varepsilon_{\mathbf{p}}$ is the kinetic energy of a Fermi atom.

In the BEC regime, the atomic chemical potential μ is related to the atomic scattering length a_F by the formula $\bar{\mu} \equiv -1/2ma_F^2$ [1, 2]. We note that $\bar{\mu}$ is large and negative as we approach the BEC limit ($a_F \rightarrow +0$). In this regime, since the BCS order parameter at $T = 0$ is given by $\Delta_{\text{MF}} = \sqrt{16/3\pi}|\bar{\mu}|^{1/4}\varepsilon_F^{3/4}$ [2], we find $|\bar{\mu}| \gg \Delta_{\text{MF}}$. Thus, we can use approximation $E_{\mathbf{p}} \simeq \varepsilon_{\mathbf{p}} + |\bar{\mu}|$ in (3). In addition, the energy gap E_g is given by $E_g = \sqrt{|\mu|^2 + \Delta_{\text{MF}}^2} \simeq |\bar{\mu}|$, so that the binding energy E_{bind} of a Cooper-pair is given by $E_{\text{bind}} = 2E_g = 2|\bar{\mu}|$.

In the BEC regime, we expand (2) in powers of $\delta\Delta$, retaining terms up to $n = 4$. We recall that the Gaussian approximation[1, 2, 7, 8] only keeps fluctuation terms to order




FIG. 1: (a) Interaction between molecules $V_{\text{eff}}^{(4)}$ mediated by Fermi atoms. The dotted lines describe the molecular Boson propagator. (b) Particle-non-conserving interaction $V_{\text{eff}}^{(3)}$.

$n = 2$. In the superfluid phase, we find

$$\begin{aligned}
S_{\Delta} = & \frac{1}{\beta} \sum_q \Phi_q^{\dagger} (-i\nu_n + \varepsilon_q^M + U_M n_c) \Phi_q \\
& + \frac{n_c U_M}{2} \frac{1}{\beta} \sum_q (\Phi_q \Phi_{-q} + \Phi_{-q}^{\dagger} \Phi_q^{\dagger}) \\
& + \frac{1}{\beta^2} \sum_{q, K} V_{\text{eff}}^{(3)}(q, K) (\Phi_{\frac{K}{2}+q}^{\dagger} \Phi_{\frac{K}{2}-q}^{\dagger} \Phi_K + \Phi_K^{\dagger} \Phi_{\frac{K}{2}-q} \Phi_{\frac{K}{2}+q}) \\
& + \frac{1}{2\beta^3} \sum_{q, q', K} V_{\text{eff}}^{(4)}(q, q', K) \Phi_{\frac{K}{2}+q'}^{\dagger} \Phi_{\frac{K}{2}-q'}^{\dagger} \Phi_{\frac{K}{2}-q} \Phi_{\frac{K}{2}+q}.
\end{aligned} \tag{4}$$

Here, $U_M \equiv 4\pi(2a_F)/M$ (where $M = 2m$) is the bare s -wave molecular interaction obtained in what amounts to the time-dependent GL theory[1, 2]; $\varepsilon_q^M \equiv q^2/2M$ is the kinetic energy of a free molecule, and $i\nu_n$ is the pair boson Matsubara frequency, with $p \equiv (\mathbf{p}, i\omega_m)$. In the BEC limit of interest, the fluctuations in the order parameter $\delta\Delta_q$ involve the molecular condensate. We introduce a renormalized molecular field $\Phi_q \equiv \delta\Delta_q/\eta$, where $\eta = [\sum_{\mathbf{p}} 1/4(\varepsilon_{\mathbf{p}} - \bar{\mu})^2]^{-1/2}$. Defining $\phi_M \equiv \Delta_{\text{MF}}/\eta$ as the equilibrium value of the Bose-condensate order parameter, at $T = 0$, one can show that $n_c \equiv \phi_M^2 = n_F/2$ (where n_F is the number of Fermi atoms) gives the number of Bose-condensed molecules.

$V_{\text{eff}}^{(4)}$ in (4) describes an interaction between molecules mediated by the Fermi gas [see Fig. 1(a)], given by

$$V_{\text{eff}}^{(4)}(q, q', K) = \frac{\eta^4}{\beta} \sum_p G_{11}^0(p + \frac{q - q'}{2} + \frac{K}{4}) G_{11}^0(-p + \frac{q + q'}{2} + \frac{K}{4})$$




FIG. 2: Calculated momentum dependence of the boson-boson interaction $V_{\text{eff}}^{(4)}(q, 0, 0)$ (solid line). The dashed line is the Lorentzian fit given by $U_M/(1 + (q/\Lambda)^2)$. The vertical dotted line shows the position of the cutoff momentum Λ .

$$\times G_{11}^0(-p - \frac{q+q'}{2} + \frac{K}{4})G_{11}^0(p - \frac{q-q'}{2} + \frac{K}{4}). \quad (5)$$

Here, G_{11}^0 is the diagonal (1,1)-component of the BCS mean-field Green's function defined in (3). $V_{\text{eff}}^{(3)}$ in (4) represents a particle-non-conserving interaction, given by [see also Fig. 1(b)]

$$V_{\text{eff}}^{(3)}(q, K) = -\frac{\eta^3}{\beta} \sum_p G_{11}^0(p + \frac{K}{2})G_{11}^0(-p + \frac{K}{2})G_{12}^0(p + q). \quad (6)$$

This interaction involves the off-diagonal (anomalous) Green's function G_{12}^0 defined in (3), so that $V_{\text{eff}}^{(3)}$ only exists in the superfluid phase.

For $q = q' = K = 0$, one can show that $V_{\text{eff}}^{(4)}(0, 0, 0) = U_M$ and $V_{\text{eff}}^{(3)}(0, 0) = \sqrt{n_c}U_M$, where $U_M = 4\pi(2a_F)/M$. Using these results in place of $V_{\text{eff}}^{(4)}(q, q', K)$ and $V_{\text{eff}}^{(3)}(q, K)$ in (4), we reproduce the previous results obtained in the literature, namely, $a_M = 2a_F$ [1, 2]. However, $V_{\text{eff}}^{(4)}$ and $V_{\text{eff}}^{(3)}$ actually depend on momentum, and go to zero with a characteristic cutoff momentum Λ . To calculate this cutoff momentum Λ , we expand (5) and (6) in terms of the relative momentum (taking the center-of-mass-momentum \mathbf{K} to be zero[12]), keeping terms of $O(q^2)$. We obtain $V_{\text{eff}}^{(4)}(q, q', 0) \simeq U_M F(q)F(q')$ and $V_{\text{eff}}^{(3)}(q, 0) \simeq \sqrt{n_c}U_M F(q)$, where $F(q) = 1 - (q/\Lambda)^2$. The characteristic momentum is found to be $\Lambda = 4/\sqrt{5}a_F$. For larger values of the momentum, we replace $F(q)$ by the Lorentzian $[1 + (q/\Lambda)^2]^{-1}$. This gives a very good fit to the momentum dependent interaction, as shown in Fig. 2. In terms of energy, we

can write $F(q) = [1 + \varepsilon_{\mathbf{q}}^M/\omega_c]^{-1}$, where the interaction or molecular cutoff energy is defined by

$$\omega_c \equiv \frac{\Lambda^2}{2M} = \frac{8}{5}|\bar{\mu}|. \quad (7)$$

We note that ω_c is comparable to the binding energy $E_{\text{bind}} = 2|\bar{\mu}|$ of a molecule in the BEC limit. Thus we conclude that ω_c is a *physical* cutoff determined by the dissociation energy of a molecule in the BEC phase.

To summarize, when we replace $V_{\text{eff}}^{(4)}(q, q', K)$ and $V_{\text{eff}}^{(3)}(q, K)$ in (4) with $V_{\text{eff}}^{(4)}(0, 0, 0)$ and $V_{\text{eff}}^{(3)}(0, 0)$, respectively, we have to include the *physical* momentum cutoff Λ , as discussed above. We recall that $\Phi_q = \delta\Delta_q/\eta$ describes *excitations* from the Bose-condensate. The molecular Bose field involving the condensate part is given by $\phi_q \equiv \Phi_q + \phi_M$. Using ϕ_q in (4), we obtain

$$\begin{aligned} S_{\Delta} = & \frac{1}{\beta} \sum_q \phi_q^\dagger (-i\nu_n + \varepsilon_q^M - \mu_M) \phi_q \\ & + \frac{1}{2\beta^3} \frac{4\pi(2a_F)}{M} \sum_{q, q', K}^{\Lambda} \phi_{\frac{K}{2}+q'}^\dagger \phi_{\frac{K}{2}-q'}^\dagger \phi_{\frac{K}{2}-q} \phi_{\frac{K}{2}+q}, \end{aligned} \quad (8)$$

where $\mu_M \equiv U_M n_c$ is the molecular chemical potential. We note that $V_{\text{eff}}^{(3)}$ has been absorbed into the last term of (8). The corresponding Hamiltonian has the form

$$\begin{aligned} H_{\text{BEC}} = & \sum_{\mathbf{q}} \phi_{\mathbf{q}}^\dagger (\varepsilon_{\mathbf{q}}^M - \mu_M) \phi_{\mathbf{q}} \\ & + \frac{4\pi(2a_F)}{M} \sum_{q, q', K}^{\Lambda} \phi_{\frac{K}{2}+q'}^\dagger \phi_{\frac{K}{2}-q'}^\dagger \phi_{\frac{K}{2}-q} \phi_{\frac{K}{2}+q}. \end{aligned} \quad (9)$$

This Hamiltonian is also obtained in the normal phase, but now with $\mu_M = 4(\mu + \sqrt{|\mu||\bar{\mu}|})$.

The effective Hamiltonian in (9) gives a clear physical picture of the two different molecular scattering lengths found in earlier literature, namely $a_M = 2a_F[1, 2]$ and $a_M = (0.6 \sim 0.75)a_F[3, 4]$. Although $U_M = 4\pi(2a_F)/M$ looks like the low-energy expression for *s*-wave interaction, (9) shows that U_M is actually a *bare* interaction involving a cutoff energy ω_c . In the case of two molecules in a vacuum, to obtain the low-energy interaction $U_M^{2b} \equiv 4\pi a_M^{2b}/2M$, we have to *renormalize* U_M to remove the need for a high-energy cutoff[13]. In the usual manner, this is given by

$$\begin{aligned} \frac{4\pi a_M^{2b}}{M} &= \frac{\frac{4\pi(2a_F)}{M}}{1 + \frac{4\pi(2a_F)}{M} \sum \omega_c \frac{1}{2\varepsilon_{\mathbf{q}}^M}} = \frac{4\pi}{M} \frac{2a_F}{1 + 2.28} \\ &= \frac{4\pi(0.61a_F)}{M}. \end{aligned} \quad (10)$$

Here we note that a_F in the denominator has been canceled out by the a_F involved in ω_c , defined in (7). The resulting renormalized Hamiltonian has the same form as (9), but now with $0.61a_F$ in place of $2a_F$ [14]. Eq. (10) agrees with the recent four-fermion (two-molecule) analysis, which gives $a_M = 0.6a_F$ [4]. Thus, we find that $a_M = 2a_F$ is the bare molecular scattering length before renormalization, while $a_M = (0.6 \sim 0.75)a_F$ is the *renormalized* two-molecule scattering length in which the high-energy processes up to ω_c have been incorporated.

The effective action in (8) is also useful to calculate the molecular scattering length a_M in a *many-particle* system. In this case, a_M is affected by many-body effects, as well as by temperature, which are not included in the $T = 0$ two-molecule result in (10). To include these effects, we apply the RG theory developed for an atomic Bose-condensed gas[11] to deal with (8), including the physical cutoff Λ . In the 1-loop level, the RG equations are given by[11], for $\mu_M < 0$,

$$\frac{d\mu_M}{dl} = 2\mu_M - \frac{\Lambda^3}{\pi^2} U_M N(\varepsilon_\Lambda^M - \mu_M), \quad (11)$$

$$\begin{aligned} \frac{dU_M}{dl} = & -U_M - \frac{\Lambda^3}{2\pi^2} U_M^2 \left[\frac{2N(\varepsilon_\Lambda^M - \mu_M) + 1}{2(\varepsilon_\Lambda^M - \mu_M)} \right. \\ & \left. + 4\beta N(\varepsilon_\Lambda^M - \mu_M)[N(\varepsilon_\Lambda^M - \mu_M) + 1] \right], \end{aligned} \quad (12)$$

and for $\mu_M \geq 0$ [15],

$$\begin{aligned} \frac{d\mu_M}{dl} = & 2\mu_M - \frac{\Lambda^3}{2\pi^2} U_M \left[\frac{\varepsilon_\Lambda^M + \mu_M}{E_\Lambda^M} [2N(E_\Lambda^M) + 1] - 1 \right. \\ & \left. + 4\beta N(E_\Lambda^M)[N(E_\Lambda^M) + 1] \right], \end{aligned} \quad (13)$$

$$\begin{aligned} \frac{dU_M}{dl} = & -U_M - \frac{\Lambda^3}{2\pi^2} U_M^2 \left[\frac{2N(E_\Lambda^M) + 1}{2E_\Lambda^M} \right. \\ & \left. + 4\beta N(E_\Lambda^M)[N(E_\Lambda^M) + 1] \right]. \end{aligned} \quad (14)$$

Here, $E_\Lambda^M = \sqrt{\varepsilon_\Lambda^M(\varepsilon_\Lambda^M + 2\mu_M)}$, and $N(\varepsilon)$ is the Bose distribution function. μ_M , U_M , and T involve the trivial scaling as $\mu_M(l) = \mu_M e^{2l}$, $U_M(l) = U_M e^{-l}$, $T(l) = T e^{2l}$. We solve the RG equations together with the equations for the number of molecules $n_M = n_F/2 = n_c + n_n$. Here, $n_c = \lim_{l \rightarrow \infty} \mu_M(l)/U_M(l)e^{-3l}$ ($\mu_M(l) > 0$) is the number of Bose-condensed molecules, and the number of non-condensate molecules n_n is determined by[11],

$$n_n = \frac{\Lambda^3}{2\pi^2} \int_0^\infty dl N(\varepsilon_\Lambda^M - \mu_M) e^{-3l} \quad (\mu_M < 0), \quad (15)$$




FIG. 3: Many-body molecular scattering length a_M as a function of temperature. p_F is the Fermi momentum for a free Fermi gas. We show results down to $T = 0$, although the RG equations (13) and (14) are not valid for temperatures such that $T/T_F \lesssim 0.1(p_F a_F)$. The inset shows T_c in the BEC regime determined from the fixed points of the RG equations (13) and (14). The dashed line shows $T_c^{\text{ideal}} = 0.218T_F$.

$$n_n = \frac{\Lambda^3}{2\pi^2} \int_0^\infty dl \left[\frac{\varepsilon_\Lambda^M + \mu_M}{2E_\Lambda^M} [N(\varepsilon_\Lambda^M - \mu_M) + 1] - \frac{1}{2} \right] e^{-3l} \quad (\mu_M \geq 0). \quad (16)$$

Figure 3 shows the *many-body* molecular scattering length a_M at finite temperatures. The scattering length a_M depends on temperature, and the temperature dependence is stronger as one approaches the crossover regime. At T_c , a_M vanishes. This is expected since if molecules are to be stable at T_c , the scattering between them must vanish at T_c . We note that a similar T -dependent scattering length was also obtained earlier in the study of Bose-condensed gases[11, 16]. However, in the present situation, one may have such large values of a_F near the resonance that the decrease in the effective value of a_M may have observable consequences at finite temperatures. In atomic Bose gases, the small size of the scattering length meant this region was restricted to temperatures very close to T_c and hence of less interest.

As shown in the inset in Fig.3, the many-body effects enhance T_c compared with $T_c^{\text{ideal}} = 0.218T_F$ for the transition temperature of a non-interacting Bose gas. In the extreme BEC limit, we find $\Delta T_c \equiv T_c - T_c^{\text{ideal}} \propto (p_F a_F) \propto (n_M^{1/3} a_M)$, consistent with the previous work on

the BEC in Bose gases[11, 17, 18, 19]. Although T_c in the inset initially increases with $p_F a_F$ from the BEC side, it will eventually decrease in the crossover region, smoothly going into the weak-coupling BCS result[7].

Our theory is only valid in so far as the effect of fluctuations involving terms higher order than $n = 4$ in (2) is small. Comparing the fourth order term with the sixth order term above T_c , we find that the fourth order term is dominant as long as $(p_F a_F)^{-1} \gtrsim 1$, i.e., we are not in the unitarity region.

To conclude, we have studied the effective interaction between bound molecules in the BEC regime of a superfluid Fermi gas. We have included the binding energy of a molecule, which naturally leads to a physical cutoff energy in the molecular Bose gas. For the two-molecule case, we showed that this cutoff gives a simple physical explanation of the difference between the two molecular scattering lengths, the bare value $a_M = 2a_F$ [1, 2] and the renormalized value $a_M = (0.6 \sim 0.75)a_F$ [3, 4]. We have also shown that the *many-body* scattering length deviates considerably from the two-body scattering length as the temperature approaches T_c if we are close to the unitarity limit. By employing the RG technique, the present paper includes fluctuations past the mean-field BCS theory in a more sophisticated way than the original crossover theory developed by Nozières and Schmitt-Rink[8].

I would like to thank Allan Griffin for stimulating discussions and comments, as well as a critical reading of this paper. This work was supported by funds from Ministry of Education of Japan and from NSERC Canada.

-
- [1] M. Randeria, in *Bose-Einstein Condensation*, edited by A. Griffin et al. (Cambridge University Press, N.Y., 1995), p.355. Earlier references are given here.
 - [2] C. Sa de Melo, et al., Phys. Rev. Lett. **71**, 3202 (1993).
 - [3] P. Pieri and G. Strinati, Phys. Rev. B **61**, 15370 (2000).
 - [4] D. Petrov, et al., Phys. Rev. Lett. **93**, 090404 (2004); Phys. Rev. A **71**, 012708 (2005).
 - [5] E. Timmermans et al., Phys. Lett. A **285**, 228 (2001).
 - [6] M. Holland et al., Phys. Rev. Lett. **87**, 120406 (2001).
 - [7] Y. Ohashi and A. Griffin, Phys. Rev. Lett. **89**, 130402 (2002). See also Y. Ohashi and A. Griffin, Phys. Rev. A **67**, 033603 (2003); **67**, 063612 (2003).

- [8] P. Nozières and S. Schmitt-Rink, J. Low. Temp. Phys. **59**, 195 (1985).
- [9] G. Astrakharchik et al., Phys. Rev. Lett. **93**, 200404 (2004).
- [10] M. Bartenstein et al., Phys. Rev. Lett. **92**, 120401 (2004).
- [11] M. Bijlsma and H. T. C. Stoof, Phys. Rev. A **54**, 5085 (1996).
- [12] Including the K -dependence of $V_{\text{eff}}^{(4)}$, we obtain another cutoff for the center-of-mass momentum as $\Lambda_K \equiv \sqrt{32/3}/a_F$, which corresponds to the energy of an incident molecule given by $\varepsilon_{\Lambda_K/2}^M = (4/3)|\mu|$. We note that Λ_K is not important in determining the low-energy value of a_M .
- [13] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University Press, 2002).
- [14] The numerical factor 0.61 in (10) weakly depends on the way of introducing the cutoff in the present analysis. When we use the smooth cutoff $F(q)F(q')$ for the molecular interaction in (9), we obtain $a_B^{2b} = 0.71a_F$.
- [15] The RG equations in the 1-loop level has an infrared divergence below T_c [11]. We have eliminated this divergence, using the prescription[11] that one replaces the coefficients in the Bogoliubov transformation as $(u_{\mathbf{p}}, v_{\mathbf{p}}) = (1, 0)$. This method is valid when we can approximate $E_{\mathbf{p}} = \sqrt{\varepsilon_{\mathbf{p}}^M(\varepsilon_{\mathbf{p}}^M + 2\mu_M)} \simeq \varepsilon_{\mathbf{p}}^M + \mu_M$. Since the temperature works as a low energy cutoff, this condition can be written as $T \gtrsim \mu_M$, or $T/T_F \gtrsim 0.1p_F a_F$ (assuming the $T = 0$ result $a_M = 0.6a_F$). Because $(p_F a_F)^{-1} \gg 1$ in the BEC region, this condition is largely satisfied, except at very low temperatures.
- [16] H. Shi and A. Griffin, Phys. Rep. **304**, 2 (1998).
- [17] P. Grüter et. al., Phys. Rev. Lett. **79**, 3549 (1997).
- [18] G. Baym et al., Phys. Rev. Lett. **83**, 1703 (1999).
- [19] P. Arnold et al., Phys. Rev. Lett. **65**, 013606 (2001).