

## Critical Dynamics of Three-Dimensional Spin Systems with Long-Range Interactions

S.V. Belim

Omsk State University, 55-a, pr. Mira, Omsk, Russia, 644077 nn

(dated: March 23, 2024)

A field-theoretic description of critical behavior of Ising systems with long-range interactions is obtained by using the Padé-Borel summation technique in the two-loop approximation directly in the three-dimensional space. It is shown that long-range interactions affect the relaxation time of the system.

PACS numbers: 64.60.-i

It was shown in [1] that effects due to long-range interaction are essential for the critical behavior of Ising systems. The renormalization-group approach to spin systems with long-range interactions developed in [2] directly in the three-dimensional space made it possible to calculate the static critical exponents in the two-loop approximation. However, analogous calculations of critical dynamics have never been performed for these systems.

In this paper, a field-theoretic description of critical behavior of homogeneous systems with long-range interactions is developed directly for  $D = 3$  in the two-loop approximation. The model under analysis is the classical spin system with exchange integral depending on the distance between spins. The corresponding Hamiltonian is

$$H = \frac{1}{2} \sum_{ij} J(\mathbf{r}_i - \mathbf{r}_j) S_i S_j; \quad (1)$$

where  $S_i$  is a spin variable and  $J(\mathbf{r}_i - \mathbf{r}_j)$  is the exchange integral. This model is thermodynamically equivalent to the  $O(n)$ -symmetric Ginzburg-Landau-Wilson model defined by the effective Hamiltonian

$$H = \int d^D x \left[ \frac{1}{2} (\nabla \phi)^2 + u_0 \phi^4 \right]; \quad (2)$$

where  $\phi$  is a fluctuating order parameter,  $D$  is the space dimension,  $\phi_0 = T/T_c$  ( $T_c$  is the critical temperature), and  $u_0$  is a positive constant. Critical behavior strongly depends on the parameter characterizing the interaction as a function of distance. It was shown in [3] that long-range interaction is essential when  $0 < \alpha < 2$ , whereas systems with  $\alpha = 2$  exhibit critical behavior characteristic of short-range interactions. For this reason, the analysis that follows is restricted to the case of  $0 < \alpha < 2$ .

Relaxational dynamics of spin systems near the critical temperature can be described by a Langevin-type equation for the order parameter:

$$\frac{\partial \phi}{\partial t} = -\phi \left[ \frac{H}{\phi} + \gamma_0 h \right]; \quad (3)$$

where  $\gamma_0$  is a kinetic coefficient,  $\phi(\mathbf{x};t)$  is a Gaussian random force (representing the effect of a heat reservoir) defined by the probability distribution

$$P = A \exp \left[ -\frac{1}{2} \int d^D x dt \phi^2(\mathbf{x};t) \right] \quad (4)$$

with a normalization factor  $A$ , and  $h(t)$  is an external field thermodynamically conjugate to the order parameter. The temporal correlation function  $G(\mathbf{x};t)$  of the order-parameter field can be found by solving Eq. (3) with  $H[\phi]$  given by (2) for  $\phi[\mathbf{x};h]$  averaging the result over  $P$ , and retaining the component linear in  $h(0)$

$$G(\mathbf{x};t) = \frac{1}{h(0)} \langle \phi'(\mathbf{x};t) \phi(\mathbf{0};0) \rangle; \quad (5)$$

where

$$\langle \phi'(\mathbf{x};t) \phi(\mathbf{0};0) \rangle = B^{-1} \int d^D x' \int_0^\infty dt' g'(\mathbf{x};t) P; \quad (6)$$

$$B = \int d^D x \int_0^\infty dt g P; \quad (7)$$

When applying the standard renormalization-group procedure to this dynamical model, one has to deal with substantial difficulties. However, it was shown in [4] that the model of critical dynamics in homogeneous systems without long-range interaction based on a Langevin-type equation is equivalent to that described by the standard Lagrangian [5]

$$L = \int d^D x dt \left[ \frac{1}{2} \dot{\phi}^2 + i \phi \left( \frac{\partial \phi'}{\partial t} + \frac{H}{\phi} \right) \right]; \quad (8)$$

where  $\phi'$  denotes an auxiliary field. The corresponding correlation function  $G(\mathbf{x};t)$  of the order parameter is defined for a homogeneous system as

$$G(\mathbf{x};t) = \frac{1}{Z} \langle \phi'(\mathbf{0};0) \phi(\mathbf{x};t) \rangle = \frac{1}{Z} \int d^D x' \int_0^\infty dt' g(\mathbf{x};t) \exp(-L[\phi';\phi]);$$

where

$$Z = \int d^D x \int_0^\infty dt \int d\phi \exp(-L[\phi';\phi]); \quad (10)$$

Instead of dealing with the correlation function, it is reasonable to invoke the Feynman diagram technique and represent the corresponding vertex in the two-loop approximation as

$$^{(2)}(k; i; 0; u_0; 0) = 0 + k \frac{i!}{0} 96u_0^2 D_0; \quad (11)$$

$$D_0 = \frac{3}{4} \frac{\int \frac{d^D q d^D p}{(1 + j^2)(1 + p^2)}}{1}$$

$$(3 + j^2 + p^2 + p + q \quad i! = )$$

The next step in the field-theoretic approach is the calculation of the scaling functions  $J_1$ ,  $J_0$ ,  $G$ , and  $D_0$  in the renormalization-group differential equation for vertexes

$$\frac{\partial}{\partial \epsilon} + \frac{\partial}{\partial u} \quad \frac{\partial}{\partial \epsilon} + \frac{\partial}{\partial \epsilon} \frac{m}{2}, \quad (12)$$

$$^{(m)}(k; i; ; u; ; ) = 0;$$

where the scaling parameter  $\epsilon$  is introduced to change to dimensionless variables.

Further analysis requires the use of the function  $\Phi$  and the dynamic scaling function  $\Psi$ .

An expression for  $\Phi$  in the two-loop approximation were obtained in [2]:

$$= (4 - D) \frac{1}{1} 36uJ_0 + 1728 - 2J_1 - J_0^2 \frac{2}{9} G u^2; \quad (13)$$

$$J_1 = \int \frac{d^D q d^D p}{(1 + j^2)^2 (1 + p^2) (1 + j^2 + p^2 + 2pq)^2};$$

$$J_0 = \int \frac{d^D q}{(1 + j^2)^2};$$

$$G = \frac{\int \frac{d^D q d^D p}{(1 + j^2 + k^2 + 2kq)^2 (1 + p^2)}}{1}$$

$$(1 + j^2 + p^2 + 2pq)^2)$$

The function  $\Psi$  calculated in the two-loop approximation is

$$= (2 - D) 2 \Phi^0 G u^2; \quad (13)$$

$$D^0 = \frac{\partial D_0}{\partial (i! = )} \quad i=0, i=0:$$

Defining the effective interaction vertex  $v = u_0$  one obtains the following expressions for  $\Phi$  and  $\Psi$ :

$$= (2 - D) \frac{1}{1} 36v + 1728 - 2\bar{J}_1 - \frac{2}{9} \bar{G} v^2; \quad (14)$$

$$= (2 - D) 96 (\bar{\Phi}^0 - \bar{\Phi}) v^2;$$

$$\bar{J}_1 = \frac{J_1}{J_0^2} \quad \bar{G} = \frac{G}{J_0^2} \quad \bar{\Phi} = \frac{D^0}{J_0^2};$$

This redefinition is meaningful for  $D=2$ . In this case,  $J_0, J_1, G$  and  $D^0$  are divergent functions. Introducing the

cuto parameter  $\Lambda$ , we obtain finite expressions for the ratios

$$\frac{J_1}{J_0^2} = \frac{\int \frac{d^D q d^D p}{(1 + j^2)^2 (1 + p^2)}}{\int \frac{d^D q}{(1 + j^2)^2}} \quad i. h^Z \quad d^D q = (1 + j^2)^2 i_2;$$

$$\frac{G}{J_0^2} = \frac{\int \frac{d^D q d^D p}{(1 + j^2 + k^2 + 2kq)^2 (1 + p^2)}}{\int \frac{d^D q d^D p}{(1 + j^2)^2 (1 + p^2)}} \quad i. h^Z \quad d^D q = (1 + j^2)^2 i_2;$$

$$\frac{D^0}{J_0^2} = \frac{3}{4} \frac{\int \frac{d^D q d^D p}{(1 + j^2)^2 (1 + p^2)}}{\int \frac{d^D q d^D p}{(1 + j^2)^2 (1 + p^2)}} \quad i. h^Z \quad d^D q = (1 + j^2)^2 i_2;$$

as  $\epsilon \rightarrow 1$ .

The integrals are performed numerically. For  $D=2$ , a sequence of  $J_1=J_0^2$  and  $G=J_0^2$  corresponding to various values of  $\epsilon$  is calculated and extrapolated to infinity.

Critical behavior is completely determined by the stable fixed points of the renormalization group (RG) transformation. These points can be found from the condition

$$(v) = 0: \quad (16)$$

The effective interaction vertexes were evaluated at the stable fixed points of the RG transformation in [2].

The dynamic critical exponent  $z$  characterizing critical slowing-down of relaxation is determined by substituting the effective charges at a fixed point into the scaling function:

$$z = 2 + \quad : \quad (17)$$

The table shows the stable fixed points of the RG transformation and the values of the dynamic critical exponent for  $1.5 \leq \epsilon \leq 1.9$ . When  $0 < \epsilon < 1.5$ , the only fixed point is the unstable gaussian one,  $v = 0$ .

A comparison of the present results with the value  $z = 2.017$  of the dynamic critical exponent for three-dimensional systems with short-range interactions obtained in [6] demonstrates the essential role played by long-range interactions in critical dynamics of spin systems. In particular, the system's relaxation time increases according to the scaling  $t \sim T_c j^z$ , where  $z$  is the critical exponent characterizing the increase in the correlation radius near a critical point. In three-dimensional systems with long-range interaction, both critical dynamics and static behavior become increasingly gaussian as the long-range interaction parameter decreases. When  $\epsilon \rightarrow 1.8$ , critical behavior is virtually gaussian.

The work is supported by Russian Foundation for Basic Research N 04-02-16002.

Table 1.

a	v	z
1.5	0.015151	2.000072
1.6	0.015974	2.000180
1.7	0.020485	2.000777
1.8	0.023230	2.001529
1.9	0.042067	2.006628

- 
- [1] E. Luijten and H. M. Ebbingfeldt, Phys. Rev. Lett. 86, 5305 (2001).
- [2] S. V. Belin, Pis'ma Zh. Eksp. Teor. Fiz. 77, 118 (2003) [JETP Lett. 77, 112 (2003)].
- [3] M. E. Fisher, S.-K. Ma, and B. G. Nickel, Phys. Rev. Lett. 29, 917 (1972).
- [4] C. De Dominicis, Lett. Nuovo Cimento 12, 567 (1975).
- [5] E. Brezin, J. C. LeGuillon, and J. Zinn-Justin, Phys. Rev. D 8, 434 (1973); Phys. Rev. D 8, 2418 (1973).
- [6] V. V. Prudnikov, S. V. Belin, A. V. Ivanov, et al., Zh. Eksp. Teor. Fiz. 114, 972 (1998) [JETP 87, 527 (1998)].