

Slave boson model for two-dimensional trapped Bose-Einstein condensate

K. K. Rajagopal^a

^a*Scuola Normale Superiore, I-56126 Pisa, Italy*

Abstract

A system of N bosons in a two-dimensional harmonic trap is considered. The system is treated in term of the slave boson representation for hard-core bosons which is valid in the arbitrary density regimes. I discuss the consequences of higher order interactions on the density profiles by mapping the slave boson equation to the known Kohn-Sham type equation within the Density functional scheme.

Key words: Bose-Einstein condensation; Slave-boson model.

PACS: 03.75.Fi, 05.30.Jp, 32.80.Pj

1 Introduction

The possibility of Bose-Einstein condensation (BEC) phase transition occurring in a 2D geometry has a remarkable history and breakthroughs. Recent experiments have realized a quasi-two-dimensional (Q2D) cold Bose gas by tuning the anisotropy of the trapping potential [1,2]. This achievement has attracted vast attention on the new physics appearing in the currently studied low-dimensional trapped gases and the nature of finite-size and finite-temperature effects. The effect of dimensionality on the existence and character of BEC or superfluid phase transition has seen a spurt of interest in recent years. A peculiar feature of the low-dimensionality (1D or 2D geometry) at finite temperature is the absence of a 'true condensate' following the Bogoliubov k^{-2} theory, believed to be originated from the long wave-length phase fluctuations [3]. This has been proved in the seminal work of Mermin and Wagner [4] and Hohenberg [5]. A Kosterlitz-Thouless transition to a superfluid state still occurs on cooling through the binding of vortices of opposite vorticity [6] and

* Corresponding author, e-mail: k.rajagopal@sns.it

leads to an algebraic decay of the one-body density matrix. However, in a trapped 2D fluid the modification of the density of states caused by the confining potential allow a true condensate to exist even at finite temperature. Another important consequence lower dimensionality is that the T-matrix for two-body collisions *in vacuo* at low momenta and energy in the dilute regime vanishes [3,7]. It is then necessary to evaluate the scattering processes between pairs of Bose particles by taking into account the presence of surrounding condensate particles [8–11].

The dilute limit ($\rho a^2 \ll 1$), where ρ is the uniform density of particles and a is the s-wave scattering length of a Bose gas can be described quite satisfactorily by traditional approaches, such as the Gross-Pitaevskii (GP) approach in the weakly interacting regime [12]. However, it is intuitively obvious that as the density of particles in the condensate or the scattering length increases, the nature of interactions will become more and more complex, and one will need to take even three-body and higher order effects into account. It is a crucial point since in the strictly two-dimensional trap, the scattering length of the bosons become larger than the axial thickness of the condensate cloud. Thus we need to go beyond the mean-field approach to describe the system of dense Bose gas .

These limitations motivates me to address the problem particularly the 2D dense condensate by introducing the slave boson model (SB) which is valid for arbitrary density regimes. The slave boson technique was first introduced by Kotliar and Ruckenstein [13], who used it to deal with the fermionic Hubbard model. A functional integral approach to the problem of hard-core bosons hopping on a lattice has been previously put forward by Ziegler [15] and Frésard [16]. Ziegler [15] has demonstrated that slave boson (SB) representation allows one to describe the dynamics of Bose gas taking into account three-body and higher effects at arbitrary densities for the three-dimensional (3D) case. The advantage of using slave boson representation is by the fact that there are only two states per site in a hard-core system: a lattice site is either empty or occupied by a single boson. A hopping process of a boson appears as an exchange of an empty site with singly occupied site.

The paper is outlined as follows. In section 2 the Bose gas with hard-core interaction defined as an effective functional integral form is summarized in the slave boson formalism. The SB equation is evaluated in section 3. Some concluding remarks are presented in section 4.

2 The Slave Boson model

A Bose gas, defined as a grand canonical ensemble of bosons, can be described in a second quantization formalism, for instance using a functional integral representation [18] as following:

$$Z = \int \exp(-S[\phi^*, \phi]/\hbar) \prod d[\phi^*]d[\phi] \quad (1)$$

where the Euclidean action $S[\phi^*, \phi]$ of the complex field ϕ needs to be defined accordingly to the model system of interest (*e.g.* GP, SB, ect.). In this work I will resort to the slave boson representation of hard-core bosons which has been exhaustively defined in the literatures [13,15,17,19]. The representation relies on the Bose-Hubbard model where the particle trades its position with an empty site on the lattice. Both particle as well as the empty site are described by the corresponding creation and annihilation operators. In the slave boson representation for instance, a lattice site x that is occupied by boson is represented by complex field b_x and a site that is empty by e_x . Thus the slave boson action S_{sb} can be translated into the following picture [13,14]

$$S_{sb} = \sum_{x,x'} b_x^* e_x t_{x,x'} b_{x'} e_{x'}^* + V_x b_x^* b_x + i\lambda_x (e_x^* e_x + b_x^* b_x - 1) \quad (2)$$

where the first term describes the exchange of bosons and empty sites in a hopping process at sites x and x' with rate $t_{x,x'}$. In the second term I have introduced an external confining potential V_x . The field λ_x in the last term of the above equation enforces a constraint $e_x^* e_x + b_x^* b_x = 1$ which guarantees that a site is either empty or singly occupied. It is indeed the core condition of the slave boson model. With this choice of action, the functional integral in Eq. (1) can be written explicitly describing the partition function of the grand canonical ensembles of bosons as:

$$Z = \int e^{-S_{sb}} \prod_x d\lambda_x db_x db_x^* de_x de_x^*. \quad (3)$$

The slave boson field e_x and b_x can be combined to form a collective field $b_x^* e_x \rightarrow \Phi_x$. Then the constraint field λ_x and the slave bosons fields can be integrated out which finally leads to the action for the collective field Φ_x . This was demonstrated in detail in [15,17]. Here, I only present the results:

$$Z = \int e^{-S_b - S_1} \prod_x d\Phi_x d\Phi_x^* \quad (4)$$

with the hopping("kinetic") term

$$S_b = \sum_{x,x'} \Phi_x (1 - t)_{x,x'}^{-1} \Phi_{x'}^* \approx \sum_x (1 - \tau)^{-1} |\Phi_x|^2 + \frac{\tau}{6a^2(1 - \tau)^2} \Phi_x (\nabla^2 \Phi_x^*) \quad (5)$$

and the potential term

$$S_1 = - \sum_x \log \left(e^{-V_x - 1/4} \int d\varphi \exp[-\varphi^2] \frac{\sinh \sqrt{(\varphi - V_x^2)^2 + |\Phi_x|^2}}{\sqrt{(\varphi - V_x^2)^2 + |\Phi_x|^2}} \right). \quad (6)$$

$V_x = x^2$ is the planar harmonic confinement where motion of atoms are in planar direction with dimensionless radial distance x . The density of the condensate in the trap can be calculated by the saddle-point approximation. In this work, we neglect the thermal fluctuations which is not important at zero temperatures for Bose-Einstein condensation to occurs in the system. However, if one needed to account for the thermal fluctuations, one could do so by studying the deviations around the saddle point in the functional integral of Eq. (4) [18]. That means, we have to look for the solution order parameter Φ_x which minimizes $S_b + S_1$ taking a saddle-point approximation on the functional integral Eq. (4) to obtain the generalized non-linear Schrödinger equation [17]

$$\left(\frac{\tau}{6a^2} \nabla^2 + \alpha + \alpha^2 \frac{\partial S_1}{\partial |\Phi_x|^2} \right) \Phi_x = 0, \quad (7)$$

where $\alpha = 1 - \tau$. The term $\partial S_1 / \partial |\Phi_x|^2$ contains higher-order interaction (three-body interactions, *etc.*). However this equation is difficult to solve as it is because the term $\partial S_1 / \partial |\Phi_x|^2$ can lead to heavy non-linearity.

3 Density functional solution to the slave boson equation

Taking into account the possible effects of higher order interactions and to avert strong non-linearity in Eq. (7), I find a much feasible way to solve it by mapping to the known Kohn-Sham [20] type equation *via* Density functional theory (DFT) scheme. The main idea of the DFT formalism is in finding an equilibrium ground state energy by minimizing the following energy functional

$$E[n(x)] = T[n(x)] + E_{ex}[n(x)] + \int dx n(x) V_x \quad (8)$$

which is a unique functional of density $n(x) = |\Phi_x|^2$ for a given external potential V_x [21]. The first term in Eq. (8) is the kinetic energy of the non-interacting system while the second term represent the excess contribution of free energy $E_{ex}[n(x)]$. The remainder term is the interaction energy of the particles in an external field. Minimizing $E[n(x)]$, the ground state energy of the system can be determined by imposing the stationary equilibrium condition

$$\frac{\delta E[n(x)]}{\delta n(x)} = \mu \quad (9)$$

where the Lagrange multiplier μ can be identified as the chemical potential. This is equivalent to the condition of equilibrium for a non-interacting particles under the influence of an effective external potential [22]

$$v_{eff}[n(x)] = \frac{\delta E_{ex}[n(x)]}{\delta n(x)} + V_x. \quad (10)$$

It is evident that the equilibrium density profile $n(x)$ can be obtained for a given $v_{eff}[n(x)]$ by solving the following Kohn-Sham type equation [20],

$$\left[\frac{\delta T[n(x)]}{\delta n(x)} + v_{eff}[n(x)] \right] \Phi_x = \mu \Phi_x. \quad (11)$$

By setting $\alpha = -\mu$ and $\tau/6a^2 = -1$, Eq.(7) can be mapped to Eq. (11) yielding ,

$$\mu^2 S_1[n(x)] = \hat{V}_H[n(x)] + E_{xc}[n(x)] + \int dx n(x) V_x. \quad (12)$$

In this relation, I have introduced an exchange-correlation energy $E_{xc}[n(x)]$ by explicitly splitting from the excess free energy $E_{ex}[n(x)]$ the following Hartree energy [22],

$$\hat{V}_H[n(x)] = \frac{1}{2} \int \int dx dx' V(x-x') n(x) n(x') \quad (13)$$

with a generic inter-particle potential $V(x-x')$. In general, the exchange correlation functional energy $E_{xc}[n(x)]$ in a DFT calculation is not known exactly. Despite, one can resort to approximations such as the Local density approximation (LDA) which reads,

$$E_{xc}[n(x)] \approx \int dx E_{xc}^{hom}[\rho]|_{\rho \rightarrow n(x)} \quad (14)$$

where $E_{xc}^{hom}[\rho] = \rho \epsilon_{xc}[\rho]$ and $\epsilon_{xc}[\rho]$ is the exchange correlation energy of a homogeneous system with uniform density ρ . The Functional derivatives of the above relation can be written as [23]

$$V_{xc}[n(x)] = \frac{\delta E_{xc}[n(x)]}{\delta n(x)} = \frac{\partial(\rho \epsilon_{xc}[\rho])}{\partial \rho}|_{\rho \rightarrow n(x)}. \quad (15)$$

Hence Eq. (7) can be written explicitly as the following Kohn-Sham type equation:

$$[-\nabla^2 + r^2 + V_H[n(x)] + V_{xc}[n(x)]] \Phi_x = \mu \Phi_x, \quad (16)$$

where $V_H[n(x)] = 4n(x)/|\ln(n(x)a^2)|$ is the Hartree potential calculated from Eq. (13) using a hard-core potential [7,26]. The homogeneous exchange correlation energy $\epsilon_{xc}[\rho]$ can be approximated to the following expression [25]:

$$\epsilon_{xc}^{kolo}[\rho] \approx \frac{2\rho}{\ln(1/\rho a^2)} \left(-\frac{\ln(\ln(1/\rho a^2))}{\ln(1/\rho a^2)} \right). \quad (17)$$

Very recently Pilati et. al [31] did a variational Monte Carlo (vmc) calculation in obtaining the ground state energy of 2D bose gas using various inter-atomic potentials. The following is the fit of their beyond mean-field ground state energy :

$$\epsilon_{xc}^{vmc}[\rho] \approx \frac{2\rho}{[\ln(1/\rho a^2)]} \left(-\frac{0.86 \ln(\ln(1/\rho a^2))}{\ln(1/\rho a^2)} \right). \quad (18)$$

Fig (1) illustrate these homogeneous exchange correlation energies for various values of gas parameter ρa^2 . Setting $E_{xc}[n(x)] = 0$ one recovers the GP equation.

There are three cases to consider in solving Eq.(16), namely using information $\epsilon_{xc}^{kolo}[\rho]$, $\epsilon_{vmc}^{kolo}[\rho]$ or $E_{xc}[n(x)] = 0$. The KS equation Eq.(16) need to be solved self-consistently with the condition that the areal integral of the condensate density $n(x) = |\Phi_x|^2$ is equal to the total number N of particles. For a numerical illustration, we have taken the radial trap frequency, and the scattering length as appropriate for ^{23}Na atoms in the experiment of Görlitz *et al.* [1], namely $\omega_{\perp} = 188.4 \text{ Hz}$, s-wave scattering length $\tilde{a} = 2.8 \text{ nm}$. I am implicitly assuming however, that the trap has been axially squeezed to reach the strictly 2D scattering regime. In this work all the distances and energies have been scaled in the harmonic oscillator unit ($a_{ho} = \sqrt{\hbar/m\omega_{\perp}}$) and energies by $\hbar\omega_{\perp}/2$. Numerical iteration method by discretization using a two-step Crank-Nicholson has been used in performing this calculation for 5×10^3 and 5×10^4 23 sodium atoms.

The results obtained from Eq. (16) using information of the exchange correlation energies $\epsilon_{xc}^{kol}[\rho]$ [$n_{kol}(x)$] and $\epsilon_{xc}^{vmc}[\rho]$ [$n_{vmc}(x)$] comparing to the case with $E_{ex}[n(x)] = 0$ [$n_{GP}(x)$] are depicted in Fig. (2) and Fig. (3) for 5×10^3 and 5×10^4 atoms respectively. The main effect of the correlation potentials can be observed where the condensate density reaches the peak at the trap center. So it is feasible to explain the observation by measuring the differences $\Delta n^* = n_{kol}(0) - n_{GP}(0)$ and $\Delta \tilde{n} = n_{kol}(0) - n_{vmc}(0)$ at the trap centre as done by Nunes [24]. For a fixed number of atoms N the conditions $\int dx \Delta n^* = 0$ and $\int dx \Delta \tilde{n} = 0$ are imposed so that the rest of curves $\Delta n^*, \Delta \tilde{n}$ are well determined at $x = 0$. The differences $\Delta n^*, \Delta \tilde{n}$ increases with $\Delta n^*/n_{kol}, \Delta \tilde{n}/n_{kol}$ varying from 9% and 1.33% for $N = 5 \times 10^3$ reaching about 13% and 2.27% for $N = 5 \times 10^4$ respectively. The increment in $\Delta n^*, \Delta \tilde{n}$ corresponds to the magnitude of reduction in the effective interaction energy of the system. The small increment in $\Delta \tilde{n}$ shows that the density profile predicted using the correlation energies ϵ_{xc}^{vmc} agrees well with the one obtained using ϵ_{xc}^{kol} . However what is more spectacular is the large values of Δn^* and the increment of $\Delta \tilde{n}/n_{kol}$ from 9% to 13% as the number the number of atom increases from $N = 5 \times 10^3$ to $N = 5 \times 10^4$. The effect of exchange-correlation clearly shows a reduction in effective interaction energy thus the profiles are more localized at the centre of the trap in comparison to the GP case. This argument can also be supported by the observation in Fig (1) where the magnitude of homogeneous

exchange correlation energies decreases as the density increases. On the other hand $\epsilon_{xc}[\rho]$ is negligible for the very dilute regimes $\rho a^2 \ll 0.01$ justifying GP approximation is only a good model in this limit

4 Conclusion

In summary, the harmonically trapped 2D Bose gas at zero temperature described by the slave boson model has been studied for both the dilute and dense limits. The central issue of this work is in how to treat the slave boson term $\partial S_1 / \partial |\Phi_x|^2$ in the highly non-linear partial differential equation (7). The importance of the higher order interactions *via* the exchange-correlation energy has been revealed by mapping Eq. (7) to the Kohn-Sham scheme of DFT in the dense limit. This latter term is responsible for the correction of up to 9% and 13% of the peak density for $N = 5 \times 10^3$ and $N = 5 \times 10^4$ atoms at the center of trap. In other words these exchange correlation has restored the correct high density behaviour of the inhomogeneous Bose condensate imposed by the strict hard-core condition of the slave-boson representation. On the opposite dilute regime I have demonstrated that the slave boson equation (7) can be approximated to a Gross-Pitaevskii type equation (*via* DFT).

At absolute zero temperature where the thermal fluctuation is frozen out and quantum fluctuation is prominent. This quantum fluctuation plays an important role in the phase transition from superfluid to mott insulator [27–29] in the three-dimensional case. Recently Wu et.al. [30] have demonstrated that a vortex in 2D optical lattice can induce the SF-MI transition. As a future direction a practical realization of this transition will be investigated using the slave-boson model of the 2D condensate accomodating a single or many vortices.

Acknowledgement

I would like to thanks Professor Ziegler, Professor S. K. Adhikari and Ch. Moseley for useful comments. Also acknowledging Inst. für Physik, Augsburg for the hospitality and financial support during early part of this work.




Fig. 1. Homogeneous exchange correlation energy $\epsilon_{xc}[\rho]$ versus the gas parameter ρa^2 for the expression ϵ_{xc}^{kol} (solid line) compared to ϵ_{xc}^{vmc} (dashed lines)




Fig. 2. Density profiles $n(x)$ for the condensate (in units $a_{ho}^{-2}N^{-1}$, with $a_{ho} = \sqrt{\hbar/m\omega_{\perp}}$) versus the radial distance x (in units of a_{ho}) obtained by solving Eq.(16) using information of ϵ_{xc}^{kol} (solid line), ϵ_{xc}^{vmc} (long-dashed lines) and setting $E_{xc}[n(x)] = 0$ (short-dashed lines).




Fig. 3. caption as in Fig. (2) for $N = 5 \times 10^4$ atoms.

References

- [1] A. Görlitz, J. M. Vogels, A. E. Leanhardt, C. Raman, T. L. Gustavson, J. R. Abo-Shaeer, A. P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband, and W. Ketterle. *Phys. Rev. Lett.*, 87:130402, 2001.
- [2] A. I. Safanov, S. A. Vasilyev, I. S. Yasnikov, I. I. Lukashevich and S. Jaakkola. *Phys. Rev. Lett.*, 81:4545, 1998; S. Burger, F. S. Cataliotti, C. Fort, P. Maddaloni, F. Minardi and M. Inguscio *Europhys. Lett.*, 57:1, 2002; V. Schweikhard, I. Coddington, P. Engels, V. P. Mogendorff and E. A. Cornell. *Phys. Rev. Lett.*, 92:040404, 2004.
- [3] V. N. Popov. *Functional Integral in Quantum Field Theory and Statistical Physics*. Reidl, Dordrecht, 1983.
- [4] N. D. Mermin and H. Wagner. *Phys. Rev. Lett.*, 17:1133, 1966.
- [5] P. C. Hohenberg. *Phys. Rev.*, 158:383, 1967.
- [6] J. M. Kosterlitz and D. J. Thouless. *J. Phys. C.*, 6:1181, 1973.
- [7] M. Schick. *Phys. Rev. A.*, 3:1067, 1971.
- [8] H.T.C. Stoof and M.Bijlsma. *Phys. Rev. E*, 47:939, 1993.
- [9] M.Bijlsma and H.T.C. Stoof. *Phys. Rev. A.*, 55:498, 1997.
- [10] M. D. Lee, S. A. Morgan, M. J. Davis, and K. Burnett. *Phys. Rev. A.*, 65:043617, 2002.
- [11] K. K. Rajagopal, P. Vignolo, and M. P. Tosi. *Physica B*, 344:157, 2004.
- [12] E.P. Gross. *Nuovo Cimento*, 20:454, 1961; *J.Math.Phys.(N.Y.)*, 4:195, 1963; L.P. Pitaevskii. *Zh.Eksp.Teor.Fiz*, 40:646, 1961.
- [13] G. Kotliar and A. E. Ruckenstein. *Phys. Rev. Lett.*, 57:1362, 1986.
- [14] S. E. Barnes. *J. Phys. F.*, 6:1375, 1976; P. Wölffe and T. Li *Z. Phys. B.*, 78:45, 1990.
- [15] K.Ziegler. *Europhys. Lett.*, 23:463, 1993.
- [16] R. Frésard. *cond-mat/9405053*, 1994.
- [17] K. Ziegler and A. Shukla. *Phys. Rev. A.*, 56:1438, 1997.
- [18] J. W. Negele and H.Orland. *Quantum Many-Particle Systems*. Addison-Wesley, New York, 1988.
- [19] D. B. M. Dickerscheid, D. van. Oosten, P. J. H. Denteneer and H. T. C. Stoof *Phys. Rev. A.*, 68:043623, 2003.
- [20] W. Kohn and L. J. Sham. *Phys. Rev.*, 140:1133, 1965.

- [21] P. Hohenberg and W. Kohn. *Phys. Rev.*, B136:864, 1964.
- [22] S. Moroni and G. Senatore. *Phys. Rev. B.*, 44:9864, 1991.
- [23] A. P. Albus, F. Illuminati and M. Wilkins. *Phys. Rev. A.*, 67:063606, 2003.
- [24] G. S. Nunes. *J. Phys. B: At. Mol. Opt. Phys.*, 32:4293, 1999.
- [25] E. B. Kolomeisky, J. P. Straley. *Phys. Rev. B.*, 46:11749, 1992.
- [26] B. Tanatar. *J. Phys. B: At. Mol. Opt. Phys.*, 35:2719, 2002.
- [27] S. Sachdev. *Quantum Phase Transition*. Cambridge University Press, Cambridge, 2001.
- [28] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch. *Nature*, 415:39, 2002.
- [29] W. Zwerger. *J. Opt. B*, 5:s9, 2003.
- [30] C. Wu, H. Chen, J. Hu, and S.-C. Zhang. *Phys. Rev. A.*, 69:43609, 2004.
- [31] S. Pilati, J. Boronat, J. Casulleras, and S. Giorgini. *cond-mat/0411049*, 2004.