

Local energy statistics in disordered systems: a proof of the local REM conjecture¹

Anton Bovier²

Weierstraß-Institut
für Angewandte Analysis und Stochastik
Mohrenstrasse 39, 10117 Berlin, Germany

and

Institut für Mathematik
Technische Universität Berlin
Strasse des 17. Juni 136, 12623 Berlin, Germany

Irina Kurkova³

Laboratoire de Probabilités et Modèles Aléatoires
Université Paris 6
4, place Jussieu, B.C. 188
75252 Paris, Cedex 5, France

Abstract: Recently, Bauke and Mertens conjectured that the local statistics of energies in random spin systems with discrete spin space should in most circumstances be the same as in the random energy model. Here we give necessary conditions for this hypothesis to be true, which we show to hold in wide classes of examples: short range spin glasses and mean field spin glasses of the SK type. We also show that, under certain conditions, the conjecture holds even if energy levels that grow moderately with the volume of the system are considered.

Keywords: universality, level statistics, random energy model, extreme value theory, disordered systems, spin glasses

AMS Subject Classification: 60G70, 82B45

¹Research supported in part by the DFG in the Dutch-German Bilateral Research Group "Mathematics of Random Spatial Models from Physics and Biology" and by the European Science Foundation in the Programme ERDSES.

²e-mail: bovier@wias-berlin.de

³e-mail: kourkova@ccr.jussieu.fr

1. Introduction.

In a recent paper [BaMe], Bauke and Mertens have formulated an interesting conjecture regarding the behaviour of local energy level statistics in disordered systems. Roughly speaking, their conjecture can be formulated as follows. Consider a random Hamiltonian, $H_N(\omega)$, i.e., a real-valued random function on some product space, S^N , where S is a finite space, typically $\{0, 1\}$, of the form

$$H_N(\omega) = \sum_{A \subset [N]} \alpha_A(\omega); \quad (1.1)$$

where $[N]$ are finite subsets of \mathbb{Z}^d of cardinality, say, N . The sum runs over subsets, A , of $[N]$ and α_A are random local functions, typically of the form

$$\alpha_A(\omega) = J_A \prod_{x \in A} \omega_x \quad (1.2)$$

where J_A , $A \subset [N]$, is a family of (typically independent) random variables, defined on some probability space, $(\Omega; \mathcal{F}; P)$, whose distribution is not too singular. In such a situation, for typical ω , $H_N(\omega) \sim \sqrt{N}$, while $\sup_{\omega} H_N(\omega) \sim N$. Bauke and Mertens then ask the following question: Given a fixed number, E , what is the statistics of the values $N^{-1/2} H_N(\omega)$ that are closest to this number, and how are configurations, ω , for which these good approximations of E are realised, distributed on S^N ? Their conjectured answer, which at first glance seems rather surprising, is quite simple: find $N_{\pm E}$ such that $P(|N^{-1/2} H_N(\omega) - E| \leq N_{\pm E}) \sim 2N_{\pm E}$; then, the collection of points, $\{N^{-1/2} H_N(\omega) - E\}$ over all $\omega \in S^N$, converges to a Poisson point process on \mathbb{R}_+ . Furthermore, for any finite k , the k -tuple of configurations, $\omega^1, \omega^2, \dots, \omega^k$, where the k -best approximations are realised, is such that all of its elements have maximal Hamming distance between each other. In other words, the asymptotic behavior of these best approximations of E is the same, as if the random variables $H_N(\omega)$ were all independent Gaussian random variables with variance N , i.e., as if we were dealing with the random energy model (REM) [Dor1]. Bauke and Mertens call this "universal REM-like behaviour".

A comparable result had previously been conjectured by Mertens [Mer1] in the particular case of the number partitioning problem. In that case, the function H_N is simply given by

$$H_N(\omega) = \sum_{i=1}^N X_i \omega_i; \quad (1.3)$$

with X_i iid. random variables uniformly distributed on $[0, 1]$, $i = 1, 2, \dots, N$, and one is interested in the distribution of energies near the value zero (which corresponds to an optimal

partitioning of the N random variables, X_i , into two groups such that their sum in each group is as similar as possible). This conjecture was later proven by Borgs, Chayes, and Pittel [BCP, BCM P]. It should be noted that in this problem, one needs, of course, take care of the obvious symmetry of the Hamiltonian under the transformation $\sigma_i \rightarrow -\sigma_i$. An extension of these results in the spirit of the REM conjecture was proven recently in [BCM N], i.e., when the value zero is replaced by an arbitrary value, E .

In [BK 2] we generalised this result to the case of the k -partitioning problem, where the random function to be considered is actually vector-valued (consisting of the vector of differences between the sums of the random variables in each of the k subsets of the partition). To be precise, we considered the special case where the subsets of the partition are required to have the same cardinality, N/k (restricted k -partitioning problem). The general approach to the proof we developed in that paper sets the path towards the proof of the conjecture by Bauke and Mertens that we will present here.

The universality conjecture suggests that correlations are irrelevant for the properties of the local energy statistics of disordered systems for energies near "typical energies". On the other hand, we know that correlations must play a rôle for the extremal energies near the maximum of $H_N(\cdot)$. Thus, there are two questions beyond the original conjecture that naturally pose themselves: (i) assume we consider instead of fixed E, N -dependent energy levels, say, $E_N = N C$. How fast can we allow E_N to grow for the REM-like behaviour to hold? and (ii) what type of behaviour can we expect once E_N grows faster than this value? We will see that the answer to the first question depends on the properties of H_N , and we will give an answer in models with Gaussian couplings. The answer to question (ii) requires a detailed understanding of $H_N(\cdot)$ as a random process, and we will be able to give a complete answer only in the case of the GREM, when H_N is a hierarchically correlated Gaussian process. This will be discussed in a separate paper [BK 05].

Our paper will be organized as follows. In Chapter 2, we prove an abstract theorem, that implies the REM-like-conjecture under three hypothesis. This will give us some heuristic understanding why and when such a conjecture should be true. In Chapter 3 we then show that the hypothesis of the theorem are fulfilled in two classes of examples: p -spin Sherrington-Kirkpatrick like models and short range Ising models on the lattice. In both cases we establish conditions on how fast E_N can be allowed to grow, in the case when the couplings are Gaussian.

Acknowledgements: We would like to thank Stephan Mertens for interesting discussions.

2. Abstract theorems.

In this section we will formulate a general result that implies the REM property under some concise conditions, that can be verified in concrete examples. This will also allow us to present the broad outline of the structure of the proof without having to bother with technical details. Note that our approach is rather different from that of [BCMN] that involves computations of moments.

Our approach to the proof of the Mertens conjecture is based on the following theorem, which provides a criterion for Poisson convergence in a rather general setting.

Theorem 2.1: Let $V_{i,M} \geq 0$, $i \in \mathbb{N}$, be a family of non-negative random variables satisfying the following assumptions: for any $\ell \in \mathbb{N}$ and all sets of constants $b_j > 0$, $j = 1, \dots, \ell$,

$$\lim_{M \rightarrow \infty} \sum_{(i_1, \dots, i_\ell) \in \mathbb{N}^\ell} P \left(\sum_{j=1}^{\ell} V_{i_j, M} < b_j \right) = \prod_{j=1}^{\ell} b_j \quad (2.1)$$

where the sum is taken over all possible sequences of different indices (i_1, \dots, i_ℓ) . Then the point process

$$P_M = \sum_{i=1}^M \delta_{V_{i,M}}; \quad (2.2)$$

on \mathbb{R}_+ , converges weakly in distribution, as $M \rightarrow \infty$, to the standard Poisson point process, P on \mathbb{R}_+ (i.e., the Poisson point process whose intensity measure is the Lebesgue measure).

Remark: Theorem 2.1 was proven (in a more general form, involving vector valued random variables) in [BK2]. It is very similar in its spirit to an analogous theorem for the case of exchangeable variables proven in [BM] in an application to the Hopfield model. The rather simple proof in the scalar setting can be found in Chapter 13 of [B].

Naturally, we want to apply this theorem with $V_{i,M}$ given by $\frac{1}{M} \sum_{j=1}^M H_N(i, j) E_N(j)$ properly normalised.

We will now introduce a setting in which the assumptions of Theorem 2.1 are verified. Consider a product space $S^{\mathbb{N}}$ where S is a finite set. We define on $S^{\mathbb{N}}$ a real-valued random process, $Y_N(i)$. Assume for simplicity that

$$E Y_N(i) = 0; \quad E (Y_N(i))^2 = 1; \quad (2.3)$$

Define on $S^{\mathbb{N}}$

$$b_N(i; i_0) = \text{cov}(Y_N(i); Y_N(i_0)); \quad (2.4)$$

Let us also introduce the Gaussian process, Z_N , on S^N , that has the same mean and the same covariance matrix as $Y_N(\cdot)$.

Let G be the group of automorphisms on S_N , such that, for $g \in G$, $Y_N(g) = Y_N(\cdot)$, and let F be the larger group, such that, for $g \in F$, $\tilde{Y}_N(g) = \tilde{Y}_N(\cdot)$. Let

$$E_N = cN^\alpha; \quad c \in \mathbb{R}; \quad 0 < \alpha < 1/2; \quad (2.5)$$

be a sequence of real numbers, that is either a constant, $c \in \mathbb{R}$, if $\alpha = 0$, or converges to plus or minus infinity, if $\alpha > 0$. We will define sets \mathcal{N}_N as follows: If $c \neq 0$, we denote by \mathcal{N}_N be the set of residual classes of S^N modulo G ; if $c = 0$, we let \mathcal{N}_N be the set of residual classes modulo F . We will assume throughout that $|\mathcal{N}_N| \geq N^\alpha$, for some $\alpha > 1$. Define the sequence of numbers

$$\eta_N = \frac{c}{2} e^{E_N^2/2} |\mathcal{N}_N|^{-1}; \quad (2.6)$$

Note that η_N is exponentially small in N^{-1} , since $\alpha < 1/2$. This sequence is chosen such that, for any $b > 0$,

$$\lim_{N \rightarrow \infty} |\mathcal{N}_N|^{-1} \mathbb{P}(\tilde{Y}_N(\cdot) - E_N \leq b \eta_N) = b; \quad (2.7)$$

For $\ell \in \mathcal{N}_N$, and any collection, $\ell^1; \dots; \ell^r \in \mathcal{N}_N$, we denote by $B_N(\ell^1; \dots; \ell^r)$ the covariance matrix of $Y_N(\cdot)$ with elements

$$b_{i,j}(\ell^1; \dots; \ell^r) = b_N(\ell^i, \ell^j); \quad (2.8)$$

Assumptions A.

(i) Let $R_{N,\ell}$ denote the set

$$R_{N,\ell} = \{f(\ell^1; \dots; \ell^r) \in \mathcal{N}_N : \exists 1 \leq i < j \leq r, \mathbb{P}_N(\ell^i, \ell^j) \neq 0\}; \quad (2.9)$$

Then there exists a continuous decreasing function, $\phi(\cdot) > 0$, on $]0; \gamma_0[$ (for some $\gamma_0 > 0$), and $\delta > 0$, such that

$$|R_{N,\ell}| \geq 1 - \exp(-\phi(\gamma) N^{\delta/2}) |\mathcal{N}_N|^{-1}; \quad (2.10)$$

(ii) Let $\ell^r \in \mathcal{N}_N$, $r = 1; \dots; r$. Let

$$L_{N,r}^{\ell^r} = \{(\ell^1; \dots; \ell^r) \in \mathcal{N}_N : \exists 1 \leq i < j \leq r, \mathbb{P}_N(\ell^i, \ell^j) \neq 0\} \quad (2.11)$$

$$\text{rank}(B_N(\ell^1; \dots; \ell^r)) = r$$

Then there exists $d_{r,\lambda} > 0$, such that, for all N large enough,

$$\mathbb{P}_N^\lambda(j_N \leq j_N) \leq e^{-d_{r,\lambda} N} : \quad (2.12)$$

(iii) For any $\lambda \geq 1$, any $r = 1, 2, \dots$, and any $b_1, \dots, b_r \geq 0$, there exist constants, $p_{r,\lambda} \geq 0$ and $Q < 1$, such that, for any $\lambda_1, \dots, \lambda_r \geq \lambda$ for which $\text{rank}(B_N(\lambda_1, \dots, \lambda_r)) = r$,

$$\mathbb{P} \left(\sum_{i=1}^r \mathbb{Y}_N(\lambda_i) \leq \sum_{i=1}^r b_i \right) \leq Q \sum_{i=1}^r N^{-p_{r,\lambda}} : \quad (2.13)$$

Theorem 2.2: Assume the Assumptions A hold. Assume that $\lambda \in [0, 1[$ is such that, for some $\epsilon \in]0, 2[$, we have:

$$\lambda < 2; \quad (2.14)$$

$$\lambda < 2 + \epsilon \quad (2.15)$$

and

$$\lambda < 1; \quad (2.16)$$

Furthermore, assume that, for any $\lambda \geq 1$, any $b_1, \dots, b_r \geq 0$, and $(\lambda_1, \dots, \lambda_r) \in \mathbb{R}_+^r$,

$$\mathbb{P} \left(\sum_{i=1}^r \mathbb{Y}_N(\lambda_i) \leq \sum_{i=1}^r b_i \right) = \mathbb{P} \left(\sum_{i=1}^r \mathbb{Y}_N(\lambda_i) \leq \sum_{i=1}^r b_i + o(j_N) \right) : \quad (2.17)$$

Then, the point process,

$$\sum_{i=1}^r \mathbb{Y}_N(\lambda_i) \xrightarrow{d} \sum_{i=1}^r \mathbb{Y}(\lambda_i) : \quad (2.18)$$

converges weakly to the standard Poisson point process P on \mathbb{R}_+ .

Moreover, for any $\lambda > 0$ and any $b \in \mathbb{R}_+$,

$$\mathbb{P} \left(\sum_{i=1}^r \mathbb{Y}_N(\lambda_i) \leq \sum_{i=1}^r b_i \right) = \mathbb{P} \left(\sum_{i=1}^r \mathbb{Y}(\lambda_i) \leq \sum_{i=1}^r b_i \right) = 0 : \quad (2.19)$$

Remark: Before giving the proof of the theorem, let us comment on the various assumptions.

(i) Assumption A (i) holds with some ϵ in any reasonable model, but the function $\lambda \mapsto \epsilon(\lambda)$ is model dependent.

- (ii) Assumptions A (ii) and (iii) is also apparently valid in most cases, but can be tricky sometimes. An example where (ii) proved difficult is the k -partitioning problem, with $k > 2$.
- (iii) Condition (2.19) is essentially a local central limit theorem. In the case $\beta = 0$ it holds, if the Hamiltonian is a sum over independent random interactions, under mild decay assumptions on the characteristic function of the distributions of the interactions. Note that some such assumptions are obviously necessary, since if the random interactions take on only finitely many values, then also the Hamiltonian will take values on a lattice, whose spacings are not exponentially small, as would be necessary for the theorem to hold. Otherwise, if $\beta > 0$, this will require further assumptions on the interactions. We will leave this problem open in the present paper. It is of course trivially verified, if the interactions are Gaussian.

Proof: We just have to verify the hypothesis of Theorem 2.1, for V_{iM} given by $\sum_{N=1}^{\infty} \beta_N \mathcal{J}_N(\cdot)$ E_N i.e., we must show that

$$\sum_{(\beta_1, \dots, \beta_N) \in \mathbb{R}_+^{N-1}} \mathbb{P} \left(\sum_{i=1}^N \beta_i \mathcal{J}_N(\cdot) \in E_N \mid \beta_1, \dots, \beta_N \right) > b_1 N^{-1} \quad (2.20)$$

We split this sum into the sums over the set R_N^1 , and its complement. First, by the assumption (2.17)

$$\begin{aligned} & \sum_{(\beta_1, \dots, \beta_N) \in \mathbb{R}_+^{N-1}} \mathbb{P} \left(\sum_{i=1}^N \beta_i \mathcal{J}_N(\cdot) \in E_N \mid \beta_1, \dots, \beta_N \right) \\ &= \sum_{(\beta_1, \dots, \beta_N) \in \mathbb{R}_+^{N-1}} \mathbb{P} \left(\sum_{i=1}^N \beta_i \mathcal{J}_N(\cdot) \in E_N \mid \beta_1, \dots, \beta_N \right) \mathbf{1}_{R_N^1} + o(1) \end{aligned} \quad (2.21)$$

But, with $C(E_N) = \mathbf{f}_N = (x_1, \dots, x_N) \in \mathbb{R}^N : \sum_{i=1}^N \beta_i \mathcal{J}_N(x_i) \in E_N$,

$$\mathbb{P} \left(\sum_{i=1}^N \beta_i \mathcal{J}_N(\cdot) \in E_N \mid \beta_1, \dots, \beta_N \right) = \frac{\int_{C(E_N)} \frac{e^{-(\mathbf{z}; B_N^{-1}(\beta_1, \dots, \beta_N)\mathbf{z})/2}}{\det(B_N(\beta_1, \dots, \beta_N))} d\mathbf{z}}{(2\pi)^{N/2}} \quad (2.22)$$

where $B_N(\beta_1, \dots, \beta_N)$ is the covariance matrix defined in (2.8). Since β_N is exponentially small in N , we see that, uniformly for $(\beta_1, \dots, \beta_N) \in \mathbb{R}_+^{N-1}$, the integral (2.22) equals

$$\frac{1}{(2\pi)^{N/2}} \int_{C(E_N)} \frac{e^{-(\mathbf{z}; B_N^{-1}(\beta_1, \dots, \beta_N)\mathbf{z})/2}}{\det(B_N(\beta_1, \dots, \beta_N))} d\mathbf{z} (1 + o(1)); \quad (2.23)$$

where we denote by E_N the vector $(E_N; \dots; E_N)$.

We treat separately the sum (2.21) taken over the smaller set, R_N^2 , R_N^1 , and the one over R_N^1 , nR_N^2 .

Since, by (2.14), ϵ_2 is chosen such that $E_N^2 N^{-2} \neq 0$, by (2.17), (2.22), and (2.23), each term in the sum over $R_{N;l}^2$, equals

$$(2_N = \frac{P}{2}) \cdot (b_1 \dots) \cdot \exp\left(\frac{1}{2} k E_N k^2 (1 + o(N^{-2}))\right) (1 + o(1)) = (b_1 \dots) \cdot \mathfrak{P}_{N;l} \cdot (1 + o(1)); \quad (2.24)$$

uniformly for $(\epsilon_1; \dots; \epsilon_n) \in R_{N;l}^2$. Hence by Assumption A (i)

$$\begin{aligned} & \sum_{(\epsilon_1; \dots; \epsilon_n) \in R_{N;l}^2} \prod_{i=1}^n \mathfrak{P}_{N;l}(\epsilon_i) \cdot E_N j < b_{i,N} = \mathfrak{P}_{N;l} \prod_{i=1}^n \mathfrak{P}_{N;l}(\epsilon_i) (1 + o(1)) \\ & \quad \neq b_1 \dots b_n; \end{aligned} \quad (2.25)$$

Now let us consider the remaining set $R_{N;l}^1, nR_{N;l}^2$, (if it is non-empty, i.e., if strictly $\epsilon_1 < \epsilon_2$), and let us find $\epsilon_1 = 0 < \epsilon_1 < \epsilon_2 = \epsilon_2$, such that

$$\epsilon_1 = \epsilon_2 + (\epsilon_1 + 1) = 2 \quad \forall i = 0; 1; \dots; n-1; \quad (2.26)$$

which is possible due to the assumption (2.15). Then let us split the sum over $R_{N;l}^1, nR_{N;l}^2$, into n sums, each over $R_{N;l}^i, nR_{N;l}^{i+1}$, $i = 0; 1; \dots; n-1$. By (2.17), (2.22), and (2.23), we have, uniformly for $(\epsilon_1; \dots; \epsilon_n) \in R_{N;l}^i$,

$$\begin{aligned} \prod_{i=1}^n \mathfrak{P}_{N;l}(\epsilon_i) \cdot E_N j < b_{i,N} &= (2_N = \frac{P}{2}) \cdot (b_1 \dots) \cdot \exp\left(\frac{1}{2} k E_N k^2 (1 + o(N^{-i}))\right) (1 + o(1)) \\ &= C \cdot \mathfrak{P}_{N;l} \cdot e^{N^2 \epsilon_i}; \end{aligned} \quad (2.27)$$

for some constant $C < 1$. Thus by Assumption A (i),

$$\begin{aligned} & \sum_{R_{N;l}^i, nR_{N;l}^{i+1}} \prod_{i=1}^n \mathfrak{P}_{N;l}(\epsilon_i) \cdot E_N j < b_{i,N} = C \cdot \mathfrak{P}_{N;l} \cdot nR_{N;l}^{i+1} \prod_{i=1}^n \mathfrak{P}_{N;l} \cdot e^{N^2 \epsilon_i} \\ & \quad = C \exp\left((\epsilon_i + 1)N^{-(i+1)} + N^2 \epsilon_i\right); \end{aligned} \quad (2.28)$$

that, by (2.26), converges to zero, as $N \rightarrow \infty$, for any $i = 0; 1; \dots; n-1$. So the sum (2.21) over $R_{N;l}^1, nR_{N;l}^2$ vanishes.

Now we turn to the sum over collections, $(\epsilon_1; \dots; \epsilon_n) \in R_{N;l}^1$. We distinguish the cases when $\det(B_N(\epsilon_1; \dots; \epsilon_n)) = 0$ and $\det(B_N(\epsilon_1; \dots; \epsilon_n)) \neq 0$. For the contributions from the latter case, using Assumptions A (i) and (iii), we get readily that,

$$\begin{aligned} & \sum_{(\epsilon_1; \dots; \epsilon_n) \in R_{N;l}^1, \text{rank}(B_N(\epsilon_1; \dots; \epsilon_n)) = n} \prod_{i=1}^n \mathfrak{P}_{N;l}(\epsilon_i) \cdot E_N j < b_{i,N} = \mathfrak{P}_{N;l} \cdot e^{-(1)N^{-(1)}} \cdot Q_{j,N} \cdot N^P \\ & \quad = C N^P \cdot \exp\left((1)N^{-(1)} + E_N^2 = 2\right); \end{aligned} \quad (2.29)$$

The right-hand side of (2.29) tends to zero exponentially fast, if condition (2.16) is verified.

Finally, we must deal with the contributions from the cases when the covariance matrix is degenerate, namely

$$\sum_{\substack{(\mathbf{y}^1, \dots, \mathbf{y}^l) \in \mathbb{R}^{N \times l} \\ \text{rank}(B_N(\mathbf{y}^1, \dots, \mathbf{y}^l)) = r}} P(\mathbf{y}_N^1 : \mathbf{y}_N^l : E_N \mathbf{y}^j < b_{1N}) ; \quad (2.30)$$

for $r = 1, \dots, l-1$. In the case $c = 0$, this sum is taken over the set L_N^r , since \mathbf{y}^1 and \mathbf{y}^0 in \mathbb{R}^N are different, i.e. $\mathbf{y}_N^1 \notin \mathbf{y}_N^0$ by definition of L_N^r . In the case $c \neq 0$, this sum is taken over l -tuples $(\mathbf{y}^1, \dots, \mathbf{y}^l)$ of different elements of \mathbb{R}^N , i.e., such that $\mathbf{y}_N^i \notin \mathbf{y}_N^j$, for any $1 \leq i < j \leq l$. But for all N large enough, all terms in this sum over l -tuples, $(\mathbf{y}^1, \dots, \mathbf{y}^l)$, such that $\mathbf{y}_N^i = \mathbf{y}_N^j$, for some $1 \leq i < j \leq l$, equal zero, since the events $\{\mathbf{y}_N^i : E_N \mathbf{y}^j < b_{1N}\}$ and $\{\mathbf{y}_N^i : E_N \mathbf{y}^j < b_{jN}\}$, with $E_N = cN$, $c \neq 0$, are disjoint. Therefore (2.30) is reduced to the sum over L_N^r , in the case $c \neq 0$ as well. Then, by Assumptions A (ii) and (iii), it is bounded from above by

$$L_N^r \cdot Q(N)^{rN^{Pr}} \cdot \sum_{\mathbf{y}^j \in L_N^r} e^{d_{r,N}} Q(N)^{rN^{Pr}} \cdot C e^{d_{r,N}} e^{E_N^2 = 2N^{Pr}} ; \quad (2.31)$$

This bound converges to zero exponentially fast, since $E_N^2 = c^2 N^2$, with $c < 1/2$. This concludes the proof of the first part of the theorem.

The second assertion (2.19) is elementary: by (2.29) and (2.31), the sum of terms $P(\mathbf{y}_N^1 : \mathbf{y}_N^l : E_N \mathbf{y}^j < b_{1N})$ over all pairs, $(\mathbf{y}^1, \mathbf{y}^2) \in \mathbb{R}^{N \times 2}$, such that $\mathbf{y}^1 \notin \mathbf{y}^2$, converges to zero exponentially fast. Thus (2.19) follows from the Borel-Cantelli lemma.

Finally, we remark that the results of Theorem 2.2 can be extended to the case when $EY_N(\mathbf{y}) \neq 0$, if $\mathbf{y} = 0$, i.e., $E_N = c$. Note that, e.g. the unrestricted number partitioning problem falls into this class. Let now $Z_N(\mathbf{y})$ be the Gaussian process with the same mean and covariances as $Y_N(\mathbf{y})$. Let us consider both the covariance matrix, B_N , and the mean of Y_N , $EY_N(\mathbf{y})$, as random variables on the probability space $(\mathbb{R}^N; B_N; E)$, where E is the uniform law on \mathbb{R}^N . Assume that, for any $l \geq 1$,

$$B_N(\mathbf{y}^1, \dots, \mathbf{y}^l) \leq I_d; N^{1-l} ; \quad (2.32)$$

where I_d denotes the identity matrix, and

$$EY_N(\mathbf{y}) \leq D; N^{1-l} ; \quad (2.33)$$

where D is some random variable D . Let

$$e_N = \frac{q}{2} K^{-1} j_N j^1: \quad (2.34)$$

where

$$K = E e^{(c \cdot D)^2} = 2. \quad (2.35)$$

Theorem 2.3: Assume that, for some $R > 0$, $\|Y_N(\cdot)\| \leq N^R$, for all $N \geq 1$. Assume that (2.10) holds for some $\epsilon > 0$ and that (ii) and (iii) of Assumptions A are valid. Assume that there exists a set, $Q_N \subset R_N$, such that (2.17) is valid for any $(\cdot^1; \dots; \cdot^N) \in Q_N$, and that $R_N \setminus Q_N \subset j_N j^N$, with some $\epsilon > 0$. Then, the point process

$$P_N = \sum_{j_N \in R_N} e_N^{-1} Y_N(\cdot) e_N j^1: P \quad (2.36)$$

converges weakly to the standard Poisson point process P on R_+ .

Proof. We must prove again the convergence of the sum (2.20), that we split into three sums: the first over Q_N , the second over $R_N \setminus Q_N$, and the third over the complement of the set R_N . By assumption, (2.17) is valid on Q_N , and thus the terms of the first sum are reduced to

$$\begin{aligned} & \sum_{i=1}^N \int_{\mathbb{R}^N} \frac{e^{-(z \cdot E Y_N(\cdot)) B_N^{-1}(\cdot^1; \dots; \cdot^N) (z \cdot E Y_N(\cdot)) = 2}}{(2)^{N-2} \det(B_N(\cdot^1; \dots; \cdot^N))} dz \\ & = (2^{N-2})^{-1} (b_1, \dots, b_N) \otimes (e^{-(z \cdot E Y_N(\cdot)) B_N^{-1}(\cdot^1; \dots; \cdot^N) (z \cdot E Y_N(\cdot)) = 2} (1 + o(1))); \end{aligned} \quad (2.37)$$

with $e = (c; \dots; c)$, and $E Y_N(\cdot) = (E Y_N(\cdot^1); \dots; E Y_N(\cdot^N))$, since N is exponentially small and $\|Y_N(\cdot)\| \leq N^R$. By definition of Q_N , the quantities (2.37) are at most $O(j_N j^N)$, while, by the estimate (2.10) and by the assumption on the cardinality of $R_N \setminus Q_N$, the number of N -tuples of configurations in $R_N \setminus Q_N$ is exponentially smaller than $j_N j^N$. Hence

$$\begin{aligned} & \sum_{j_N \in R_N} P(\delta_{i=1}^N : Y_N(\cdot^i) \in E_N j^i < b_{i,N}) \\ & = \sum_{j_N \in Q_N} (2^{N-2})^{-1} (b_1, \dots, b_N) \otimes (e^{-(z \cdot E Y_N(\cdot)) B_N^{-1}(\cdot^1; \dots; \cdot^N) (z \cdot E Y_N(\cdot)) = 2} (1 + o(1)) + o(1)) \\ & = \sum_{j_N \in R_N \setminus Q_N} (2^{N-2})^{-1} (b_1, \dots, b_N) \otimes (e^{-(z \cdot E Y_N(\cdot)) B_N^{-1}(\cdot^1; \dots; \cdot^N) (z \cdot E Y_N(\cdot)) = 2} (1 + o(1)) + o(1)) \\ & = \frac{b_1}{j_N j^N} \cdot b_N \sum_{j_N \in R_N} e^{-(z \cdot E Y_N(\cdot)) B_N^{-1}(\cdot^1; \dots; \cdot^N) (z \cdot E Y_N(\cdot)) = 2} (1 + o(1)) + o(1): \end{aligned} \quad (2.38)$$

The last quantity converges to b_1 , by the assumptions (2.32), (2.33) and (2.35).

The sum of the probabilities, $P(\{s_{i=1}^N : \sum_{j=1}^N E_N j < \sum_{j=1}^N b_1\})$, over all N -tuples of R_N , $\sum_{j=1}^N Q_N$, contains at most $\sum_{j=1}^N j e^{-N}$ terms, while, by Assumption A (iii), (and since, for any $(s_1, \dots, s_N) \in R_N$, the rank of $B_N(s_1, \dots, s_N)$ equals N) each term is at most of order $\sum_{j=1}^N j e^{-N}$, up to a polynomial factor. Thus this sum converges to zero.

Finally, the sum of the same probabilities over the collections $(s_1, \dots, s_N) \in R_N$, converges to zero, exponentially fast, by the same arguments as those leading to (2.29) and (2.31), with $\beta_1 = 0$.

3. Examples

We will now show that the assumptions of our theorem are verified in a wide class of physically relevant models: 1) the Gaussian p -spin SK models, 2) SK-models with non-Gaussian couplings, and 3) short-range spin-glasses. In the last two examples we consider only the case $\beta_1 = 0$.

3.1 p -spin Sherrington-Kirkpatrick models, $0 < p \leq 2$.

In this subsection we illustrate our general theorem in the class of Sherrington-Kirkpatrick models. Consider $S = \{-1, 1\}^N$.

$$H_N(s) = \frac{1}{N} \sum_{1 \leq i_1 < i_2 < \dots < i_p \leq N} J_{i_1, \dots, i_p} s_{i_1} s_{i_2} \dots s_{i_p} \quad (3.1)$$

is the Hamiltonian of the p -spin Sherrington-Kirkpatrick model, where J_{i_1, \dots, i_p} are independent standard Gaussian random variables.

The following elementary proposition concerns the symmetries to the Hamiltonian.

Proposition 3.1 Assume that, for any $0 < i_1 < \dots < i_p \leq N$, $s_{i_1} \dots s_{i_p} = 0$. Then, if p is pair, either $s_i = 0$, for all $i = 1, \dots, N$, or $s_i = 0$, for all $i = 1, \dots, N$, and, if p is odd, then $s_i = 0$, for all $i = 1, \dots, N$. Assume that, for any $0 < i_1 < \dots < i_p \leq N$, $s_{i_1} \dots s_{i_p} = 0$. This is impossible, if p is pair and $s_i = 0$, for all $i = 1, \dots, N$, if p is odd.

This proposition allows us to construct the space Σ_N : If p is odd and $c \neq 0$, $\Sigma_N = S^N$, thus $\sum_{j=1}^N j = 2^N$. If p is even, or $c = 0$, Σ_N consists of equivalence classes where configurations and $-s$ are identified, thus $\sum_{j=1}^N j = 2^{N-1}$.

Theorem 3.2: Let $p \geq 1$ be odd. Let $\mu_N = S^N$. If $p = 1$ and $\alpha \in [0; 1/4[$, and, if $p = 3; 5; \dots$, and $\alpha \in [0; 1/2[$, for any constant $c \in \mathbb{R} \setminus \{0\}$ the point process

$$P_N^X = \frac{1}{2^N} \sum_{\mathbf{x} \in S^N} \exp(-c \sum_{i=1}^N x_i^2) \exp(-\alpha \sum_{i=1}^N x_i^4) \quad (3.2)$$

where $\mu_N = 2^{-N} \exp(-c \sum_{i=1}^N x_i^2) \exp(-\alpha \sum_{i=1}^N x_i^4)$, converges weakly to the standard Poisson point process, P , on \mathbb{R}_+ .

Let p be even. Let μ_N be the space of equivalence classes of S^N where \mathbf{x} and \mathbf{y} are identified. For any $\alpha \in [0; 1/2[$ and any constant, $c \in \mathbb{R}$, the point process

$$P_N^X = \frac{1}{2^N} \sum_{\mathbf{x} \in S^N} \exp(-c \sum_{i=1}^N x_i^2) \exp(-\alpha \sum_{i=1}^N x_i^4) \quad (3.3)$$

converges weakly to the standard Poisson point process, P , on \mathbb{R}_+ . The result (3.3) holds true as well in the case of $c = 0$, for p odd.

Proof of Theorem 3.2. We have to verify the assumptions of Theorem 2.2 for the process $\mu_N^{\alpha, c} = \mu_N^{\alpha, c}$. The elements of the covariance matrix (2.8) are:

$$b_{j,j}(\alpha; \dots; \alpha) = 1; \quad \alpha_{j=1} \quad (3.4)$$

$$b_{j,m}(\alpha; \dots; \alpha) = \frac{1}{2^N} \sum_{\mathbf{x} \in S^N} x_j^2 x_m^2 \exp(-c \sum_{i=1}^N x_i^2) \exp(-\alpha \sum_{i=1}^N x_i^4); \quad \alpha_{j < m} \quad (3.5)$$

It has been observed in [BKL] that its non-diagonal elements can be written as

$$b_{j,m}(\alpha; \dots; \alpha) = \frac{1}{2^N} \sum_{\mathbf{x} \in S^N} x_j^2 x_m^2 \exp(-c \sum_{i=1}^N x_i^2) \exp(-\alpha \sum_{i=1}^N x_i^4) = \frac{1}{2^N} \sum_{\mathbf{x} \in S^N} x_j^2 x_m^2 \exp(-c \sum_{i=1}^N x_i^2) \exp(-\alpha \sum_{i=1}^N x_i^4) \quad (3.6)$$

Now let us verify the Assumption A (i). Let

$$Q_{N, \alpha} = \frac{1}{2^N} \sum_{\mathbf{x} \in S^N} \exp(-c \sum_{i=1}^N x_i^2) \exp(-\alpha \sum_{i=1}^N x_i^4) \quad (3.7)$$

The α -tuples of this set satisfy the following property: for any $\alpha_1, \dots, \alpha_p \in [0; 1/2[$, the sets of sites $A_{\alpha_1, \dots, \alpha_p} = \{i : \alpha_1 = 1; \alpha_2 = 1; \dots; \alpha_p = 1\}$ has the cardinality $N^{2(\alpha_1 + \dots + \alpha_p)}$. Then it is an easy combinatorial computation to check that there exists $h > 0$, such that, for any $q \in \mathbb{R}_+$, and any $\alpha \in [0; 1/2[$,

$$Q_{N, \alpha}^q \leq \exp(-h N^{1/2}) \quad (3.8)$$

for all N large enough. By the representation (3.6), we have $\prod_{k=0}^T Q_N^{(k)} = (p-2k) R_N$, with $Q_N^{(k)} = (p-2k) \prod_{k=0}^{2k} (k-1)!!$. But, for any $2 \leq p=2l$, and any $k = 0, 1, \dots, [p/2]$, $Q_N^{(k)} = (p-2k) Q_N^{(k+1)} = (p-2(k+1))$. Therefore,

$$Q_N^{(p)} = R_N. \quad (3.9)$$

Thus, due to (3.8), Assumption A (i) is verified with $(\cdot) = 1 \leq p$, for $2 \leq p=2l$.

Let us now check the Assumption A (ii). To estimate the cardinality of L_N^r , we need to introduce an ℓ -by- N matrix, $C_p(\cdot; \cdot)$, as follows. For any given $\cdot; \cdot$, the j th column is composed of all $\prod_{i_1 \dots i_\ell}^j$, over all subsets $1 \leq i_1 < i_2 < \dots < i_\ell \leq N$. Then we have

$$C_p^T(\cdot; \cdot) C_p(\cdot; \cdot) = \prod_{i_1 \dots i_\ell}^1 B_N(\cdot; \cdot). \quad (3.10)$$

Let $\cdot; \cdot$ be such that $\text{rank}(B_N(\cdot; \cdot)) = r < \ell$. Then, r columns of the matrix $C_p(\cdot; \cdot)$ form a basis of its ℓ columns. Assume that these are, e.g., the first r columns. The matrix $C_p(\cdot; \cdot^r)$ can contain at most 2^r different rows. We will show that, for any $(\cdot; \cdot) \in L_N^r$, it can in fact not contain all 2^r rows, due to the following proposition.

Proposition 3.3: Assume that an $2^r \times r$ matrix, A , with elements, 1 or -1 , consists of all 2^r different rows. Assume that a column n of length 2^r with elements 1 or -1 is a linear combination of the columns of A . Then this column n is a multiple (with coefficient $+1$ or -1) of one of the columns of the matrix A .

Proof. The proof can be carried out by induction over r . A generalisation of this fact is proven in [BK-npp].

Now, if the matrix $C_p(\cdot; \cdot^r)$ contained all 2^r rows, then, by Proposition 3.3, for any $j = r+1, \dots, \ell$, there would exist $m = 1, \dots, r$, such that, either, for any $0 < i_1 < \dots < i_m \leq N$, $\prod_{i_1 \dots i_m}^j = \prod_{i_1 \dots i_m}^m$, or, for any $0 < i_1 < \dots < i_m \leq N$, $\prod_{i_1 \dots i_m}^j = -\prod_{i_1 \dots i_m}^m$, which would imply $\mathcal{Y}_N(\cdot^j) = \mathcal{Y}_N(\cdot^m)$. But this is excluded by the definition of L_N^r .

Thus, for any $(\cdot; \cdot) \in L_N^r$, the matrix $C_p(\cdot; \cdot^r)$ contains at most $2^r - 1$ different rows. There are $O((2^r - 1)^N)$ possibilities to construct such a matrix. Furthermore, there is only an N -independent number of possibilities to complete it by adding linear combinations of its columns to $C_p(\cdot; \cdot)$. To see this, consider the restriction of $C_p(\cdot; \cdot^r)$ to any r linearly independent rows. There are not more than $2^{r(\ell-r)}$ ways to complete it by

(r) columns of B_N of length r , that are linear combinations of its r columns. But each such choice determines uniquely linear coefficients in these linear combinations and hence the completion of the whole $C_p(\mathbf{b}_N^{(1)}; \dots; \mathbf{b}_N^{(r)})$ up to $C_p(\mathbf{b}_N^{(1)}; \dots; \mathbf{b}_N^{(r)})$. Thus $\mathbb{P}_N^r, j=0, \dots, (2^r-1)^N$.

It remains to verify the Assumption A (iii). This is easy: if $\text{rank}(B_N(\mathbf{b}_N^{(1)}; \dots; \mathbf{b}_N^{(r)})) = r$, then r of the random variables $Y_N(\mathbf{b}_N^{(1)}); \dots; Y_N(\mathbf{b}_N^{(r)})$ are linearly independent. Assume that these are, e.g., $Y_N(\mathbf{b}_N^{(1)}); \dots; Y_N(\mathbf{b}_N^{(r)})$. Then the covariance matrix $B_N(\mathbf{b}_N^{(1)}; \dots; \mathbf{b}_N^{(r)})$ is non-degenerate, and the corresponding probability is bounded from above by

$$\mathbb{P}(8_{j=1}^r Y_N(\mathbf{b}_N^{(j)}) \leq E_N j < N b_{i_j}) \leq \frac{(2^r)^r \det(B_N(\mathbf{b}_N^{(1)}; \dots; \mathbf{b}_N^{(r)}))}{(2^r)^r \det(B_N(\mathbf{b}_N^{(1)}; \dots; \mathbf{b}_N^{(r)}))} : \quad (3.11)$$

From the representation of the matrix elements of $B_N(\mathbf{b}_N^{(1)}; \dots; \mathbf{b}_N^{(r)})$, (3.5), one sees that the determinant, $(\det B_N(\mathbf{b}_N^{(1)}; \dots; \mathbf{b}_N^{(r)}))$, is a finite polynomial in the variables N^{-1} , and thus its inverse can grow at most polynomially.

Thus, we have established that Assumption A is verified. We now turn to conditions (2.14), (2.15), and (2.16) on \mathbf{b}_N . Since $\mathbf{b}_N = 1 - 2^{-p}$, for $2 \leq p \leq 2$, we should find $\mathbf{b}_N \geq 2 \leq p \leq 2$ such that $\mathbf{b}_N \leq 2^{-p}$, $\mathbf{b}_N = 2^{-p+1} = 2^{-p}$ for $2 \leq p \leq 2$, and $\mathbf{b}_N \leq 2^{-p+1} = 2^{-p}$. We see that, for any $p \geq 2$ and $2 \leq p \leq 2$, it is possible to choose $\mathbf{b}_N > 0$ small enough, and $2 \leq p \leq 2$ close enough to 1, such that these assumptions are satisfied. If $p = 1$, then such a choice is possible only for $2 \leq p \leq 4$. The assumption (2.17) need not be verified here as $Y_N(\mathbf{b}_N)$ is a Gaussian process. }

Remark: Values $p = 1$; $\mathbf{b}_N = 1/4$. The value $\mathbf{b}_N = 1/4$ is likely to be the true critical value in the case $p = 1$. In this case, one can check that the principle part of our sum gives a contribution of the form

$$\frac{\text{const}(1 + o(1))}{(2^r)^{k(k-1)/2}} \sum_{\substack{m_1, m_2, \dots, m_k \\ 1 \leq j \leq N}} \exp(-c^2 N^2) \sum_{1 \leq i < j \leq N} m_{i,j} (1 + o(1)) \frac{N}{2} \sum_{1 \leq i < j \leq N} m_{i,j}^2 (1 + o(1)) ; \quad (3.12)$$

which in turn is easily seen to be of order $(e^{c^2=2})^{k(k-1)/2}$, that it does not behave like a constant to the power k . Note that the term proportional to $\frac{N}{2}$ in the exponents arises from the off-diagonal part of the covariance matrix B_N .

If $\mathbf{b}_N > 1/4$, the contribution from the (3.12) is already of order $(e^{N^4 - 1} c^2=2)^{k(k-1)/2}$, which cannot be compensated by any normalisation of the form $\frac{k}{N}$. Thus at least the conditions of Theorem 2.1 cannot hold in this case.

3.2. Generalized p-spin SK models at level $\beta = 0$.

In this subsection we generalize Theorem 3.2 to the case of non-Gaussian process in the case of non-zero mean and $\beta = 0$. Let $p \geq 1$, U_{i_1, i_2, \dots, i_p} be any $\frac{N}{p}$ iid. random variables with $\mathbb{E}U = a$ and $\text{Var}U = 1$. Let

$$H_N(\mathbf{x}) = \frac{1}{N} \sum_{1 \leq i_1 < i_2 < \dots < i_p \leq N} \prod_{j=1}^p U_{i_1, \dots, i_p} x_{i_j} \quad (3.13)$$

Let $\psi(s) = \mathbb{E}e^{is(U-a)}$ be the generating function of $(U-a)$.

Assumption B. We will assume in this section that $\mathbb{E}|j|^3 < \infty$ and $j(s) = O(|s|^{-1})$, as $|s| \rightarrow \infty$.

Remark: The decay assumption on the Fourier transform is not optimal, but some condition of this type is needed, as the result cannot be expected for discrete distributions, where the number of possible values the Hamiltonian takes on would be finite.

We consider $Y_N(\mathbf{x}) = N^{-1/2} H_N(\mathbf{x})$. The state space \mathcal{N} is defined as in the previous example. The covariance matrix, given by (3.6), converges in law to the identity matrix by the law of large numbers. Furthermore, analogously to (3.6), we see that $\mathbb{E}Y_N(\mathbf{x}) = Q_p(N^{-1/2} \sum_{i=1}^N x_i)$, where

$$Q_p(x) = \sum_{k=0}^{\lfloor p/2 \rfloor} \frac{(-1)^k}{p} \frac{p!}{(p-2k)!} x^{p-2k} \quad (3.14)$$

By the central limit theorem, $\mathbb{E}Y_N(\mathbf{x}) \xrightarrow{d} Q_p(J)$ where J is a standard Gaussian random variable. Hence, (2.32) and (2.33) are verified and we may define the constant

$$K_p = \mathbb{E} \exp \left(-c \left(\sum_{i=1}^N Y_i \right)^2 \right) \quad (3.15)$$

Then, $e_N = K_p^{-1} j_N j^{-1} \left(\frac{p}{2} \right)$, with $j_N j = 2^N$ for p odd and $j_N j = 2^{N-1}$ for p even.

Theorem 3.4:

(i) Let p be odd. Let $\mathcal{N} = S^N$. For any $c \neq 0$, the point process

$$P_N = \sum_{i=1}^N \delta_{\frac{1}{2} K_p \left(\frac{p}{2} \right) Y_i(\mathbf{x}) - c j} \quad (3.16)$$

converges weakly to the standard Poisson point process on \mathbb{R}_+ .

(ii) Let p be odd and $c = 0$, or let p be even and $c \neq 0$. Denote by \mathcal{Y}_N the space of the 2^{N-1} equivalence classes in S^N where \sim and \sim' are identified. Then the point process

$$P_N^X = \sum_{2 \leq N} f_{2^{N-1}K_p(2^{p-1})} \mathcal{Y}_N(\cdot) \quad (3.17)$$

converges weakly to the standard Poisson point process on R_+ .

Proof of Theorem 3.4. We should check the assumptions of Theorem 2.3. The Assumptions A (i), for any $2 \leq p \leq 2^l$, and (ii) have been already verified in the proof of Theorem 3.2. We must check (iii) and also the assertion (2.17) on an appropriate subset Q_N .

We will use the construction of the matrix $C_p(\cdot; \cdot; \cdot)$ explained in the proof of Theorem 3.2, see (3.10). Let us introduce the Fourier transform

$$f^{(1); \dots; (p)}(t_1; \dots; t_p) = E \exp \{ i [t_1 (Y_N(\cdot) - E Y_N(\cdot)) + \dots + t_p (Y_N(\cdot) - E Y_N(\cdot))] \} \quad (3.18)$$

A simple computation shows that

$$f^{(1); \dots; (p)}(t_1; \dots; t_p) = \sum_{m=1}^N \frac{1}{p} f_{C_p(\cdot; \cdot; \cdot)} \mathbf{t}_m; \quad (3.19)$$

where $f_{C_p(\cdot; \cdot; \cdot)} \mathbf{t}_m$ is the m th coordinate of the product of the matrix $C_p(\cdot; \cdot; \cdot)$ with the vector $\mathbf{t} = (t_1; \dots; t_p)$.

Assumption A (iii) is valid due to the following proposition.

Proposition 3.5: There exists a constant, $Q = Q(r; b_1; \dots; b_r) > 0$, such that, for any $(\cdot; \cdot; \cdot) \in \mathcal{Y}_N$, any $r \leq N$, if $\text{rank } B_N(\cdot; \cdot; \cdot) = r$,

$$P_{\otimes_{i=1}^r \mathcal{Y}_N(\cdot)} \leq c_j \prod_{i=1}^r b_i \prod_{i=1}^r Q_N^{pr=2+1}; \quad (3.20)$$

Proof. Recall that it follows from the hypothesis that the rank of the matrix $C_p(\cdot; \cdot; \cdot)$ equals r . Let us remove from this matrix r columns such that the remaining r columns are linearly independent. They correspond to a certain subset of r configurations. Without loss of generality, we may assume that they are $(\cdot; \cdot; \cdot)^r$, i.e., we obtain the matrix $C_p(\cdot; \cdot; \cdot)^r$. Obviously,

$$P_{\otimes_{i=1}^r \mathcal{Y}_N(\cdot)} \leq c_j \prod_{i=1}^r b_i \prod_{j=1}^r \mathcal{Y}_N(\cdot) \leq c_j \prod_{i=1}^r b_i \quad (3.21)$$

Then

$$P_{8_{i=1}^N} : \mathcal{Y}_N(\cdot) \rightarrow \mathbb{C}^{j \times N} b_i$$

$$\frac{1}{(2^{-N})^r} \lim_{D \rightarrow 1} \int_{[D, D]^r} f_N^{(1, \dots, r)}(t_1, \dots, t_r) \prod_{j=1}^r \frac{e^{it_j b_j \Theta_i} - e^{-it_j b_j \Theta_i}}{2it_j} dt_j : \quad (3.22)$$

As $e_N = O(2^{-N})$, the integrand in (3.22) is bounded by

$$\frac{e^{it_j b_j \Theta_i} - e^{-it_j b_j \Theta_i}}{2it_j} \leq \min\{Q_0 2^{-N}; 2|t_j|^{-1}\} ; \quad (3.23)$$

with a constant, $Q_0 = Q_0(b_j)$. Next, let us choose in the matrix $C_p(1, \dots, r)$ any r linearly independent rows and construct from them an $r \times r$ matrix, $C^{r \times r}$. Then, by (3.19) and by Assumption B on (s)

$$f_N^{(1, \dots, r)}(t) \prod_{j=1}^r \frac{e^{it_j b_j \Theta_i} - e^{-it_j b_j \Theta_i}}{2it_j} \leq \min\{1; Q_0 N^{p=2} |C^{r \times r}|^{-1}\} ; \quad (3.24)$$

with $Q_0 > 0$. Hence, the absolute value of the integral (3.22) is bounded by the sum of two terms,

$$Q_0(b_1) \int_{|t| \leq 2^{-N}} \prod_{j=1}^r \frac{e^{it_j b_j \Theta_i} - e^{-it_j b_j \Theta_i}}{2it_j} dt_j \leq \min\{1; Q_0 N^{p=2} |C^{r \times r}|^{-1}\} \int_{|t| \leq 2^{-N}} \prod_{j=1}^r dt_j$$

$$+ \int_{|t| > 2^{-N}} \prod_{j=1}^r \frac{e^{it_j b_j \Theta_i} - e^{-it_j b_j \Theta_i}}{(2t_j)^1} \leq \min\{1; Q_0 N^{p=2} |C^{r \times r}|^{-1}\} \int_{|t| > 2^{-N}} \prod_{j=1}^r dt_j : \quad (3.25)$$

Recall that the matrix $C^{r \times r}$ has matrix elements ≤ 1 and rank r . Since the total number of such matrices is at most 2^{r^2} , the smallest absolute value of the determinant of all such matrices is some positive number that does not depend on N , but only on r . Therefore, the change of variables, $\tilde{t} = C^{r \times r} t$, in the first term shows that the integral over $|t| \leq 2^{-N}$ is of order at most $N^{pr=2} \ln 2^{rN} \leq N^{pr=2+1}$. Thus the first term of (3.25) is bounded by $Q_1 2^{-N} N^{pr=2+1}$, with some constant $Q_1 < 1$. Using the change of variables $\tilde{t} = 2^{-rN} t$ in the second term of (3.25), one can see that the integral over $|t| > 2^{-N}$ is bounded by $Q_2 2^{-N} N^{pr=2}$, with some constant $Q_2 < 1$. This concludes the proof. }

Finally, let us fix any $2 \leq l \leq 2^k$ and introduce $Q_N = Q_N^{(p)}(q)$ (defined in (3.7)) with $q = (\max_{k=0, \dots, [p/2]} \binom{2k}{p} (k-1)!)^{-1}$. By (3.9) and (3.8), it is a subset of R_N , and $j_N \leq n_{Q_N} j$ is smaller than $2^N e^{-hN^{1/2}}$, with some $h > 0$. We need to verify (2.17) for Q_N . We abbreviate

$$W_N = v^{-1} (C - EY_N(1)); \dots; (C - EY_N(\cdot)) : \quad (3.26)$$

For any $(^1; \dots; ^N) \in \mathcal{Q}_N$, we split

$$P(8_{i=1}^N \mathcal{Y}_N(^i) \leq c_j < b_1 e_N) = \sum_{m=1}^{X^4} I_N^m(^1; \dots; ^N); \quad (3.27)$$

where

$$I_N^1(^1; \dots; ^N) = \sum_{R, j=1}^Z \frac{Y' e^{it_j b_j e_N} e^{it_j b_j e_N}}{2it_j} e^{it W_N} e^{t B_N(^1; \dots; ^N)} t=2 dt; \quad (3.28)$$

$$\sum_{k \leq N^{p=6}} \sum_{j=1}^Z \frac{Y' e^{it_j b_j e_N} e^{it_j b_j e_N}}{2it_j} e^{it W_N} e^{t B_N(^1; \dots; ^N)} t=2 dt;$$

$$I_N^2(^1; \dots; ^N) = \sum_{k \leq N^{p=6}} \sum_{j=1}^Z \frac{Y' e^{it_j b_j e_N} e^{it_j b_j e_N}}{2it_j} e^{it W_N} f_N^{1; \dots; ^N}(t) e^{t B_N(^1; \dots; ^N)} t=2 dt; \quad (3.29)$$

$$I_N^3(^1; \dots; ^N) = \sum_{N^{p=6} < k \leq N^p} \sum_{j=1}^Z \frac{Y' e^{it_j b_j e_N} e^{it_j b_j e_N}}{2it_j} e^{it W_N} f_N^{1; \dots; ^N}(t) dt; \quad (3.30)$$

and

$$I_N^4(^1; \dots; ^N) = (2)^{-N} \lim_{D \rightarrow 1} \sum_{[D, D] \setminus k \leq N^p} \sum_{j=1}^Z \frac{Y' e^{it_j b_j e_N} e^{it_j b_j e_N}}{2it_j} e^{it W_N} f_N^{1; \dots; ^N}(t) dt; \quad (3.31)$$

with some $\epsilon_j > 0$ to be chosen later.

The first part of $I_N^1(^1; \dots; ^N)$ is exactly the quantity $P(8_{i=1}^N \mathcal{Y}_N(^i) \leq c_j < b_1 e_N)$. Note that

$$\sum_{j=1}^Z \frac{Y' e^{it_j b_j e_N} e^{it_j b_j e_N}}{2it_j} \leq Q 2^{-N}; \quad (3.32)$$

with some $Q < 1$. Then the second part of I_N^1 is exponentially smaller than 2^{-N} , for all $(^1; \dots; ^N) \in \mathcal{Q}_N$. We must show that I_N^2, I_N^3, I_N^4 are $o(2^{-N})$, for all $(^1; \dots; ^N) \in \mathcal{Q}_N$. This is easy due to the following proposition.

Proposition 3.6: There exist constants, $C < 1$, $\epsilon_j > 0$, such that, for all $(^1; \dots; ^N) \in \mathcal{Q}_N$, the following estimates hold:

(i) For all $k \leq N^{p=6}$,

$$f_N^{1; \dots; ^N}(t) e^{t B_N(^1; \dots; ^N)} t=2 \leq \frac{C k t^3}{N^p} e^{t B_N(^1; \dots; ^N)} t=2. \quad (3.33)$$

(ii) For all $k \neq 0$, $k \leq \frac{N^p}{2}$,

$$f_N^{(1, \dots, \ell)}(t) = e^{-k|t|^2} : \quad (3.34)$$

Proof. The proof is elementary and completely analogous to the corresponding estimate in the proof of the Berry-Essen inequality. All details are completely analogous to those in the proof of Lemma 3.5 in [BK2] and therefore are omitted. \square

Using (3.33) and (3.32), we see that $I_N^2(\ell; \dots; \ell) = O(N^{-p/2})2^{-N^\ell}$. The third term, $I_N^3(\ell; \dots; \ell)$, is exponentially smaller than 2^{-N^ℓ} by (3.34).

Finally, by (3.32) we may estimate $I_N^4(\ell; \dots; \ell)$ roughly as

$$|I_N^4(\ell; \dots; \ell)| \leq Q 2^{-N^\ell} \int_{|k| \leq \frac{N^p}{2}} |f_N^{(1, \dots, k)}(t)|^2 dt, \quad (3.35)$$

with some constant $Q < 1$. By the construction of the set Q_N (3.7), for any $(\ell; \dots; \ell) \in Q_N$, the matrix $C_1(\ell; \dots; \ell)$, (i.e., the matrix with N rows, the k th row being $(\frac{1}{k}; \frac{2}{k}; \dots; \frac{\ell}{k})$, contains at least $2^{\ell-1}$ possible different rows, each row being present at least $2^{-N^\ell}(1+o(1))$ times. Consequently, each of these rows is present in the matrix $C_p(\ell; \dots; \ell)$ at least $2^{-N^p}(1+o(1))$ times, for any $p \geq 2$. Then, by (3.19), $f_N^{(1, \dots, \ell)}(t)$ is the product of at least $2^{\ell-1}$ different characteristic functions, each is taken to the power at least $2^{-N^p}(1+o(1))$. Let us choose from a set of different rows of $C_p(\ell; \dots; \ell)$ m linearly independent ones, and denote by C the square matrix composed of them. Then there exists $\delta > 0$, such that $t^T C^T C t \geq \delta^2$, for all t , with $|k| \geq \frac{N^p}{2}$. Changing variables $s = \frac{N^p}{2}^{-1/2} C t$ in (3.35), one gets the bound

$$|I_N^4(\ell; \dots; \ell)| \leq Q 2^{-N^\ell} N^{p/2} \int_{|s| \geq \frac{N^p}{2}^{-1/2}} |Y(s)|^2 ds, \quad (3.36)$$

Assumption B made on (s) implies that (s) is aperiodic, and thus $|Y(s)| < 1$, for any $s \neq 0$. Moreover, for any $\epsilon > 0$, there exists $h(\epsilon) > 0$, such that $|Y(s)| < 1 - h(\epsilon)$, for all s with $|s| \geq \epsilon$. Therefore, the right-hand side of (3.36) does not exceed

$$Q 2^{-N^\ell} N^{p/2} (1 - h(\epsilon))^2 \int_{|s| \geq \frac{N^p}{2}^{-1/2}} |Y(s)|^2 ds, \quad (3.37)$$

where the integral is finite again due to Assumption B. Therefore, $I_N^4(\ell; \dots; \ell)$ is exponentially smaller than 2^{-N^ℓ} . This concludes the proof of (2.17) on Q_N and of the theorem. \square

3.3. Short range spin glasses.

As a final example, we consider short-range spin glass models. To avoid unnecessary complications, we will look at models on the d -dimensional torus, Λ_N , of length N . We consider Hamiltonians of the form

$$H_N(\sigma) = \sum_{A \in \Lambda_N} \sum_{x \in \Lambda_N} r_A J_A \sigma_x \quad (3.38)$$

where $e_A = \sum_{x \in \Lambda_N} \sigma_x$, r_A are given constants, and J_A are random variables. We will introduce some notation:

- (a) Let Λ_N denote the set of all $A \in \Lambda_N$, such that $r_A \neq 0$.
- (b) For any two subsets, $A, B \in \Lambda_N$, we say that $A \sim B$, if there exists $x \in \Lambda_N$ such that $B = A + x$. We denote by \mathcal{A} the set of equivalence classes of Λ_N under this relation.

We will assume that the constants, r_A , and the random variables, J_A , satisfy the following conditions:

- (i) $r_A = r_{A+x}$, for any $x \in \Lambda_N$;
- (ii) there exists $k \in \mathbb{N}$, such that any equivalence class in \mathcal{A} has a representative A_k ; we will identify the set \mathcal{A} with a uniquely chosen set of representatives contained in Λ_k .
- (iii) $\sum_{A \in \Lambda_N} r_A^2 = N^d$.
- (iv) $J_A, A \in \mathbb{Z}^d$, are a family of independent random variables, such that
- (v) J_A and J_{A+x} are identically distributed for any $x \in \mathbb{Z}^d$;
- (vi) $EJ_A = 0$ and $EJ_A^2 = 1$, and $EJ_A^3 < 1$;
- (vii) For any $A \in \mathcal{A}$, the Fourier transform $\hat{J}_A(s) = E \exp(isJ_A)$, of J_A satisfies $|\hat{J}_A(s)| = O(|s|^{-1})$ as $|s| \rightarrow \infty$.

Observe that $EH_N(\sigma) = 0$,

$$b(\sigma; 0) = N^d EH_N(\sigma) H_N(\sigma) = N^d \sum_{A \in \Lambda_N} r_A^2 J_A^2 = 1 \quad (3.39)$$

where equality holds, if $\sigma = 0$.

Note that $Y_N(\sigma) = Y_N(\sigma_0)$ (resp. $Y_N(\sigma) = Y_N(\sigma_0)$), if and only if, for all $A \in \Lambda_N$, $\sigma_A = \sigma_A^0$ (resp. $\sigma_A = \sigma_A^0$). E.g., in the standard Edwards-Anderson model, with nearest

Let us assume for simplicity that $N = nk$ is a multiple of k , and introduce the sub-lattice,

$N/k = \{0; k; \dots; (n-1)k; nk\}^d$. Write

$$\sum_{A \in N/k} r_A^2 = \sum_{A \in N/k} \sum_{y \in N/k} \sum_{x \in N/k} r_{A+y+x}^2 Z_x(\cdot) \quad (3.45)$$

where

$$Z_x(\cdot) = \sum_{y \in N/k} Y_{y;x}(\cdot) \quad (3.46)$$

has the nice feature that, for fixed x , the summands

$$Y_{x;y}(\cdot) = \sum_{A \in N/k} r_{A+y+x}^2$$

are independent for different $y; y \in N/k$ (since the sets $A+y+x$ and A^0+y^0+x are disjoint for any $A; A^0 \in N/k$). Using the Holder inequality repeatedly,

$$\begin{aligned} E \exp \sum_{x \in N/k} Z_x(\cdot) &= E \exp \sum_{x \in N/k} \sum_{y \in N/k} Y_{y;x}(\cdot) \\ &= E \exp \sum_{y \in N/k} \sum_{x \in N/k} Y_{y;x}(\cdot) \\ &= E \exp \sum_{y \in N/k} Y_{y;0}(\cdot) \end{aligned} \quad (3.47)$$

It remains to estimate the Laplace transform of $Y_{0;0}(\cdot)$,

$$E \exp t k^d Y_{0;0}(\cdot) = E \exp t k^d \sum_{A \in N/k} r_A^2; \quad (3.48)$$

and, since $E r_A = 0$, using that $e^x = 1 + x + \frac{x^2}{2} e^{x_j}$,

$$E \exp t k^d \sum_{A \in N/k} r_A^2 = E \exp \frac{t^2}{2} k^{2d} \sum_{A \in N/k} r_A^2 e^{t k^d P_{A \in N/k} r_A^2} = E \exp \frac{t^2}{2} C e^{tD}; \quad (3.49)$$

so that

$$E \exp t N^d \sum_{x \in N/k} Z_x(\cdot) \leq \exp N^d \frac{t^2}{2} C e^{tN^d D}; \quad (3.50)$$

with constants, $C; C^0; D$, that do not depend on N . To conclude the proof of the lemma, the exponential Chebyshev inequality gives,

$$P \{b(\cdot; 0) > N\} \leq \exp -N t + N^d \frac{t^2}{2} C e^{tN^d D}; \quad (3.51)$$

Choosing $t = N^d$, this gives

$$P(|b(\cdot; \cdot^0)| > N) \leq \exp\left(-\frac{2}{N} N^d\right) = 1 - C^0 e^{-N^d} = 2 \quad (3.52)$$

Choosing ϵ small enough, but independent of N , we obtain the assertion of the lemma. \square

To verify Assumptions A (ii) and (iii), we need to introduce the matrix $C = C(\cdot^1; \cdot; \cdot^0)$ with ℓ columns and \mathfrak{A}_N rows, indexed by the subsets $A \in \mathcal{A}_N$: the elements of each of its column are $r_A^1; r_A^2; \dots; r_A^{\ell}$, so that $C^T C$ is the covariance matrix, $B_N(\cdot^1; \cdot; \cdot^0)$, up to a multiplicative factor N^d .

The assumption (ii) is verified due to Proposition 3.3. In fact, let us reduce C to the matrix $\tilde{C} = \tilde{C}(\cdot^1; \cdot; \cdot^0)$ with columns $\frac{1}{A}; \frac{2}{A}; \dots; \frac{\ell}{A}$, without the constants r_A . Then, exactly as in the case of p-spin SK models, by Proposition 3.3, for any $(\cdot^1; \cdot; \cdot^0) \in L_{N^d, r}$ the matrix $\tilde{C}(\cdot^1; \cdot; \cdot^0)$ can contain at most $2^r - 1$ different columns. Hence, $\sum_{N^d, r} \tilde{C}(\cdot^1; \cdot; \cdot^0) = O((2^r - 1)^{N^d})$ while $j_N \sim (2^{N^d} =)^r$.

The assumption (iii) is verified as well, and its proof is completely analogous to that of Proposition 3.5. The key observation is that, again, the number of possible non-degenerate matrices $C^{r \times r}$ that can be obtained from $C_p(\cdot^1; \cdot; \cdot^0)$ is independent of N . But this is true since, by assumption, the number of different constants r_A is N -independent.

Finally, we define Q_N as follows. For any $A \in \mathcal{A}$, let

$$Q_N^A = \{(\cdot^1; \cdot; \cdot^0) : \exists_{1 \leq j \leq \ell} \cdot^j \in \mathcal{A}_N \text{ such that } \sum_{x \in \mathbb{Z}^d: x+A \in \mathcal{A}_N} \frac{1}{A} \cdot^j < \mathfrak{A} j^{-1} N^d\} \quad (3.53)$$

Let us define $Q_N = \bigcup_{A \in \mathcal{A}} Q_N^A \subset R_N$. By Proposition 3.8, applied to a model where $\mathfrak{A} j = 1$, for any $A \in \mathcal{A}$, we have $\# Q_N^A \leq 2^{N^d} \exp(-h_A N^d(1 - 2^{-1}))$, with some $h_A > 0$. Hence, $\# Q_N$ has cardinality smaller than $j_N \exp(-h N^d(1 - 2^{-1}))$, with some $h > 0$. The verification of (2.17) on Q_N is analogous to the one in Theorem 3.4, using the analogue of Proposition 3.6. We only note a small difference in the analysis of the term I_N^4 where we use the explicit construction of Q_N . We represent the corresponding generating function as the product of $\mathfrak{A} j$ terms over different equivalence classes of A , with representatives A_k , each term being $\sum_{x \in \mathbb{Z}^d: x+A_k \in \mathcal{A}_N} (N^{d/2} r_A(t_{x+A_k} + \dots + t_{x+A_k}))$. Next, we use the fact that for any $(\cdot^1; \cdot; \cdot^0) \in Q_N$ each of these $\mathfrak{A} j$ terms is a product of at least 2^{-1} (and of course at most 2^{ℓ}) different terms, each is taken to the power $\mathfrak{A} j^{-1} N^d 2^{-1}(1 + o(1))$. This proves the first assertion of the theorem.

The proof of the second assertion, i.e., the case $\beta > 0$ with Gaussian variables J_A is immediate from the estimates above and the abstract Theorem 2.2, in view of the fact that the condition (2.17) is trivially verified. }

References

- [BFM] H. Bauke, S. Franz, and St. Mertens. Number partitioning as random energy model. *Journal of Statistical Mechanics: Theory and Experiment*, page P04003, 2004.
- [BaMe] H. Bauke and St. Mertens. Universality in the level statistics of disordered systems. *Phys. Rev. E*, 70:025102(R), 2004.
- [BCP] C. Borgs, J. Chayes, and B. Pittel. Phase transition and finite-size scaling for the integer partitioning problem. *Random Structures & Algorithms*, 19(3-4):247-288, 2001.
- [BCMP] C. Borgs, J. T. Chayes, S. Mertens, and B. Pittel. Phase diagram for the constrained integer partitioning problem. *Random Structures & Algorithms*, 24(3):315-380, 2004.
- [BCM N] C. Borgs, J. T. Chayes, S. Mertens, and Ch. Nair. Proof of the local REM conjecture for number partitioning. preprint 2005.
- [Bo] A. Bovier. *Statistical mechanics of disordered systems*. Cambridge University Press, to appear (2005).
- [BK1] A. Bovier and I. Kurkova. Derrida's generalised random energy models. I. Models with finitely many hierarchies. *Ann. Inst. H. Poincaré Probab. Statist.*, 40(4):439-480, 2004.
- [BK2] A. Bovier and I. Kurkova. Poisson convergence in the restricted k-partitioning problem. preprint 964, *WIAS*, 2004.
- [BKL] A. Bovier, I. Kurkov, and M. Lowe. Fluctuations of the free energy in the REM and the p-spin SK models. *Ann. Probab.* 30:605-651, 2002.
- [Der1] B. Derrida. Random-energy model: an exactly solvable model of disordered systems. *Phys. Rev. B* (3), 24(5):2613-2626, 1981.
- [Der2] B. Derrida. A generalisation of the random energy model that includes correlations between the energies. *J. Phys. Lett.*, 46:401-407, 1985.
- [Mer1] St. Mertens. Phase transition in the number partitioning problem. *Phys. Rev. Lett.*, 81(20):4281-4284, 1998.
- [Mer2] St. Mertens. A physicist's approach to number partitioning. *Theoret. Comput. Sci.*, 265(1-2):79-108, 2001.