

Absence of ferromagnetism in Ising model on directed Barabasi-Albert network

Muneer A. Sumour, Physics Department, Al-Aqsa University, P.O. B. 4051, Gaza, Gaza Strip, Palestinian Authority

M. M. Shabat, Physics Department, Islamic University, P.O. B. 108, Gaza, Gaza Strip, Palestinian Authority

D. Staufer, Institute for Theoretical Physics, Cologne University, D-50923 Köln, Euroland

Email:

Shabat@mailugaza.edu, msumoor@yahoo.com, staufer@thp.unik-koeln.de

Abstract:

With up to 7 million spins, the existence of spontaneous magnetization of Ising spins on directed Barabasi-Albert networks is investigated by Monte Carlo simulations. We confirm our earlier result that the magnetization for different temperatures T decays after a characteristic time $\tau(T)$, which we extrapolate to diverge at zero temperature by a modified Arrhenius law, or perhaps a power law.

Keywords: Monte Carlo simulations, Directed Barabasi-Albert networks

Introduction:

The Ising magnet is since decades a standard tool of computational physics [1]. We apply it here to scale-free networks [2], where previous simulations [3] indicated a Curie temperature increasing logarithmically with increasing system size N . In contrast to that work we use here directed [4] as opposed to undirected networks and then apply the standard Glauber kinetic Ising model [1] to the fixed network. We try to improve our previous note [5] by making the system an order of magnitude larger, correcting a programming error, and comparing networks with two and seven neighbours.

Directed scale-free network:

Putting Ising spins onto the sites (vertices, nodes) of a network, we simulate our Ising magnetic model on directed Barabasi-Albert networks. The Barabasi-Albert network is grown such that the probability of a new site to be connected to one of the already existing sites is proportional to the number of previous connections to this already existing site: The rich get richer. In this way each new site selects exactly m old sites as neighbours.




Figure 1: Characteristic time for $M(\beta) = 3/4$ using 7 million spins for $m = 2$ neighbours and 7 million spins for $m = 7$ neighbours (+). Ten, hundred, and thousand times smaller systems are denoted by x, stars, and squares. We plot the median over nine samples in this log-log plot. The two straight lines have negative slopes 8 (left) and 12 (right).

Then each spin is influenced by the fixed number m of neighbours which it had selected when joining the network. It is not influenced by other spins which selected it as neighbour after it joined the network.

At each step, a new spin is added which builds m new connections, randomly to already existing spins. The probability for an existing spin to be chosen as neighbour is proportional to the number of its neighbours.

Ising simulations:

The Ising interaction energy is

$$E = -J \sum_i \sum_k S_i S_k \quad (S_i = \pm 1)$$

where the sum over k goes only over the m neighbours which site i had




Figure 2: Same data (except for short times) plotted as $1/\ln(\tau)$ versus T .

selected when it joined the network. We measure the temperature in units of the usual Curie temperature of the square-lattice Ising model.

First we initialize a directed Barabasi-Albert network of N sites with m neighbours (all m initial spins are connected with each other and them selves), here $m = 7$. We put an Ising spins onto every site, with all spins up, because we test here for ferromagnetism. Then with the standard Glauber (heat bath) Monte Carlo algorithm spins we search for thermal equilibrium at positive temperature. Time is measured in Monte Carlo steps MCS per spin.

As shown earlier [5] the magnetization $M(t)$ as a function of time t shows strong fluctuations and sometimes very rapid changes in a very short interval. These problems became worse when we corrected a programming error. Now the magnetization reduces to a temperature-dependent metastable value after the first time step, stays there for a long time, and then flips or tries to flip. We follow [5] and define a characteristic time τ as that time where the magnetization has decreased to $3/4$ of its initial value, Fig.1. This double-logarithmic plot first suggests a power-law divergence $(T \rightarrow 0) \rightarrow 1$. But

we see slight curvature, unusually high exponents near 10, and an exponent varying with the number $m = 2$ or 7 of neighbours. More plausible thus is the modified Arrhenius plot of Fig 2 which suggests $\propto \exp(\text{const}/T)$ for low temperatures, or $1/\ln(\) / T + \dots$: This const in the exponential is about 1.4 for $m = 2$ and about 11 for $m = 7$.

Conclusion:

In this way we confirmed the asymptotic Arrhenius extrapolation $1 = \ln \ / T$ of [5], meaning that at all finite temperatures the magnetization eventually vanishes: No ferromagnetism.

References:

- [1] David P. Landau, Kurt Binder, A guide to Monte Carlo simulation in statistical physics; Cambridge University Press (2002).
- [2] R. Albert and A.L. Barabasi, Rev. Mod. Phys. 74, 47 (2002).
- [3] A. Aleksiejuk, J.A. Holyst, D. Stauffer, Physica A 310, 260 (2002); J.O. Indekeu, Physica A 333, 461 (2004).
- [4] D. Stauffer and H. Meyer-Ottmanns, Int. J. Mod. Phys.C 15, 241 (2004).
- [5] M.A. Sumour and M.M. Shabat, Int. J. Mod. Phys. C 16, no. 4 (2005) = e-print cond-mat/0411055 at www.arXiv.org