

Super uid Phases of ^3He in Aerogel

I.A. Fomin

P. L. Kapitza Institute for Physical Problems,
Kosygina 2, 119334 Moscow, Russia

Formal derivation of criterion for selection of super uid phases of ^3He in aerogel is presented. At the strength of the derived criterion variation of the order parameter of B-like phase in magnetic field differs from that of B-phase of pure ^3He . Possible observable consequence of this difference is discussed.

PACS numbers: 67.57.Pq, 67.57.Bc, 67.57.De

1. INTRODUCTION

In pure ^3He two super uid phases (A and B) are observed. Existence of more than one super uid phase is a manifestation of unconventional character of Cooper pairing in this liquid. In both phases Cooper pairs have angular momentum $l = 1$ and spin $s = 1$. Their order parameters are 3×3 complex matrices A_j , index refers to spin and j to the orbital projection. The particular form of order parameter for the A-phase is

$$A_j^{ABM} = \frac{1}{\sqrt{2}} \hat{d} (\hat{m}_j + i \hat{n}_j); \quad (1)$$

where \hat{m} , \hat{n} are mutually orthogonal unit vectors in orbital space and \hat{d} – a unit vector in spin space, \hat{d} is an overall amplitude. For B-phase the order parameter is

$$A_j^{BW} = \frac{1}{\sqrt{3}} e^{i\varphi} R_j; \quad (2)$$

where R_j is a real orthogonal matrix. In a presence of aerogel the property of super uidity survives¹ and two super uid phases (called A-like and B-like) are observed^{2,3} as well. In a simplified form aerogel can be represented as an array of randomly distributed and randomly oriented strands of diameter $d \approx 30 \text{ \AA}$. For 98% aerogel the average distance between the

strands is $l \approx 200\text{\AA}$, i.e. comparable with the correlation length of the condensate ξ_0 , which varies depending on a pressure in the interval 160–500 Å. Aerogel makes condensate of Cooper pairs spatially nonuniform. Scattering of quasiparticles on the strands has depairing effect on components of the order parameter which extends over a distance of the order of ξ_0^4 . An array of strands induces random variation of the order parameter in space. It is convenient to represent the order parameter $A_j(r)$ as a sum of its average value and a fluctuating increment: $A_j(r) = A_j + a_j(r)$. By the definition the average $a_j(r) = \langle a_j(r) \rangle = 0$. Averaging is assumed over a volume with a characteristic size exceeding a length for which the distribution of the strands is correlated. Onset of the long-range order is formally expressed by the condition $A_j \neq 0$. The form of A_j describes the change of symmetry at the phase transition and symmetry properties of the emerging phase. Identification of A_j for A-like and B-like phases is a key question for theoretical interpretation of their properties. Fluctuating part $a_j(r)$ influences averaged properties of the superfluid phases as well and because of nonlinear dependence of free energy of these phases on the order parameter fluctuations can influence a choice of A_j . This influence has been discussed recently^{5,6,7}. In the present paper some of the previous discussion is summarised and new application of the developed approach is presented.

2. RANDOM FIELD

Since the average distance between the strands is of the order of ξ_0 perturbation of the order parameter at a given point appears as effect of several neighbouring strands. In a vicinity of T_c correlation length $(T) \approx \xi_0$ and effect of aerogel can be represented by a local random field. Its form is determined by symmetry argument and natural assumptions about properties of aerogel. Most important assumption is that the scattered quasiparticles do not exchange their spins with the strands. In practice this condition is secured by a coverage of the strands by one-two monolayers of atoms of ^4He . With this precaution aerogel can influence directly only orbital part of the order parameter. A principal contribution of interaction with aerogel to the free energy functional is of the second order in the order parameter and it can be written in a form

$$F = N(0) \int_{j1}^Z (r) A_{j1} A_{j1} d^3r; \quad (3)$$

where $N(0)$ is the density of states on Fermi level and $_{j1}(r)$ is a random static tensor field. Further properties of $_{j1}(r)$ can be found on a strength of general argument. Since F must be real $_{j1}(r)$ is Hermitian matrix and to

secure that the invariance of the energy it must be symmetrical. So $\chi_{jl}(r)$ is real and symmetrical tensor field. Characteristic amplitude of the random field can be estimated with the use of results of the paper⁴. By the order of magnitude it is $|\chi_{jl}| \sim (d=l^2)$. For 98% aerogel this ratio is $l=10$ and effect of aerogel can be treated as a perturbation. With the account of interaction term (3) Ginzburg-Landau (GL) functional takes the following form :

$$F_{GL} = N(0) \int d^3r [A_j A_j + \chi_{jl}(r) A_j A_l + \frac{1}{2} \sum_{s=1}^5 I_s + \frac{1}{2} K_1 \frac{\partial A_j}{\partial x_j} \frac{\partial A_l}{\partial x_l} + K_2 \frac{\partial A_j}{\partial x_j} \frac{\partial A_l}{\partial x_l} + K_3 \frac{\partial A_j}{\partial x_j} \frac{\partial A_l}{\partial x_l}]; \quad (4)$$

where $\chi = (T - T_c) = T_c$, I_s - 4-th order invariants in the expansion of the free energy over A_j (cf.⁸). Coefficients $K_1; \dots; K_5; K_1; K_2; K_3$ are phenomenological constants. In what follows for their evaluation the weak coupling values are assumed. Possible random corrections to the gradient terms contain extra power of the wave-vector and will not influence the following discussion. Comparison of two first terms in the functional indicates a physical meaning of the tensor $\chi_{jl}(r)$ { its isotropic part $\frac{1}{3} \delta_{jl} \chi(r)$ describes local variations of $T_c = T_c(r)$ caused by fluctuations of the density of scatterers. Anisotropic part $\chi_{jl}(r) - \frac{1}{3} \delta_{jl} \chi(r)$ describes the local splitting of T_c for different projections of angular momenta caused by the local breaking of spherical symmetry by the aerogel strands.

3. CRITERION FOR SELECTION OF THE PHASES

The order parameters of stable phases are found as minima of the functional (4). Because of a presence of the field $\chi_{jl}(r)$ the gradient terms can not be omitted. According to the procedure developed for a scalar order parameter in a scalar random field⁹ the order parameter $A_j(r) = A_j + a_j(r)$ is substituted in the GL equation, corresponding to the functional (4) and terms up to the second order in a_j and χ_{jl} are kept. After averaging over coordinate r one arrives at the equation

$$A_j + \frac{1}{2} \sum_{s=1}^5 I_s \frac{\partial I_s}{\partial A_j} + \frac{1}{2} \frac{\partial^3 I_s}{\partial A_j \partial A_n \partial A_l} \langle a_n a_l \rangle + 2 \frac{\partial^3 I_s}{\partial A_j \partial A_n \partial A_l} \langle a_n a_l \rangle = \langle a_{1lj} \rangle; \quad (5)$$

Along with A_j the obtained equation contains averages of binary products of fluctuations $\langle a_n a_1 \rangle$ etc.. Equations for a_j are obtained by separation of fast varying terms in the GL equation

$$a_j + \frac{1}{2} \sum_{s=1}^5 X_s^5 \frac{\partial^2 I_s}{\partial A_j \partial A_n} a_n + \frac{\partial^2 I_s}{\partial A_j \partial A_n} a_n - \frac{1}{2} K \frac{\partial^2 a_j}{\partial x_1^2} + 2 \frac{\partial^2 a_1}{\partial x_1 \partial x_j} = A_{1lj} : \quad (6)$$

To proceed further one needs to introduce a nontrivial modification of the procedure of Ref.⁹ caused by a more complicated form of the order parameter and its higher degeneracy. Eq.(6) and its complex conjugate form a linear inhomogeneous system. The corresponding homogeneous system has nontrivial solutions as a consequence of the degeneracy of the unperturbed GL functional (functional (4) without the random field) with respect to orbital rotations of the order parameter. These solutions are the increments of A_j and A_j at the infinitesimal rotation n :

$$\delta_j = \epsilon_n e^{jn_r} A_r; \quad \delta_j = \epsilon_n e^{jn_r} A_r; \quad (7)$$

where e^{jn_r} is the absolutely antisymmetric tensor. Eq. (6) is solved by Fourier transformation. For further argument only a character of singularity of $a_j(k)$ at $k \rightarrow 0$ is important. To find it one can neglect anisotropy of the gradient terms in Eq. (6) and to substitute $\frac{1}{2} K \frac{\partial^2 a_j}{\partial x_1^2}$ instead of $\frac{1}{2} K \frac{\partial^2 a_j}{\partial x_1^2} + 2 \frac{\partial^2 a_1}{\partial x_1 \partial x_j}$. Then Eq.(6) is multiplied by δ_j . The result is summed with its complex conjugate and the obtained equation is solved with respect to the projection $a^l(k) = a_j(k) \delta_j + a_j(k) \delta_j$:

$$a^l(k) = \frac{2}{k^2 K} \epsilon_n e^{jn_r} Q_{rl} \delta_j^{(a)}(k); \quad (8)$$

where $Q_{rl} = A_r A_l + A_l A_r$. This projection gives a contribution to the averages $\langle a_n a_1 \rangle$ which is proportional to

$$\int \langle a^l(k) a^l(-k) \rangle d^3 k = \int \epsilon_p e^{jp_r} Q_{rl} \epsilon_q e^{mq_s} Q_{sn} \langle \delta_j^{(a)}(k) \delta_n^{(a)}(-k) \rangle \frac{d^3 k}{k^4}; \quad (9)$$

Form of the correlation function in the integrand after averaging over direction of k is determined by the symmetry properties of $\delta_j^{(a)}$:

$$\int \langle \delta_j^{(a)}(k) \delta_n^{(a)}(-k) \rangle \frac{d\Omega}{4} = \langle \delta_j^{(a)}(k) \delta_n^{(a)}(-k) \rangle \frac{1}{4} [3(\delta_{jm} \delta_{jn} + \delta_{ln} \delta_{jm})]; \quad (10)$$

Finally

$$\int \langle a^\dagger(\mathbf{k}) a^\dagger(-\mathbf{k}) \rangle d^3k$$

$$\sum_{p,q} [e^{npr} e^{lqs} Q_{rl} Q_{sn} + (\delta_{pqrs} - \delta_{ps} \delta_{qr}) Q_{rn} Q_{sn}] \frac{\epsilon(\mathbf{k})}{k^2} d\mathbf{k} \quad (11)$$

The integral in the right-hand side diverges at the lower limit and the averaged products $\langle a^\dagger a^\dagger \rangle$ give diverging contributions to Eq. (5). To make the iteration scheme Eqs. (5), (6) consistent one has to require the coefficient in front of the diverging integral to be zero. Equating to zero the expression in square brackets in Eq. (11) one arrives at $Q_{rn} = q_{rn}$, where q is a real number. In terms of the order parameter this condition reads as:

$$A_r A_j + A_j A_r = \text{const} \quad (12)$$

This is a necessary condition for existence of a superfluid phase with the order parameter A_r . The phases meeting this criterion are "quasi-isotropic" i.e. the energy of their interaction with aerogel does not change at the arbitrary orbital rotation:

$$A_{j1}^{(a)} A_{j1} = 0 \quad (13)$$

Tensor of superfluid densities for these phases is isotropic and as a consequence their order parameter is not oriented by a superflow. It has been shown before⁷ that when condition (13) is met the general argument of Imry and Ma¹⁰ does not apply and the long-range order is not destroyed by the random field $\epsilon_{j1}(r)$. It might be well to point out here the difference between the argument of Imry and Ma and the argument presented in this section. In the former case the order parameter is assumed to be fixed by minimization of free energy without the random field. Possible disordering is caused by a random walk of the order parameter over its space of degeneracy. No other change of the order parameter is admitted. In the latter case all variations of the order parameter A_j are considered. That makes possible an adjustment of the order parameter to the random field. To emphasize a property of phases, which satisfy criterion (12) or (13) to resist disordering action of the random field a term "robust" was suggested⁶.

Now let us discuss a region of applicability of the above argument. Dimensionless distance from T_c : $j = (T_c - T)/T_c$ is limited from above by the condition of applicability of the GL-expansion $j \ll 1$. There exist a limitation from below. Even when criterion (12) is met average fluctuation of the order parameter diverge when $j \rightarrow 0$ as $1 = \frac{1}{j}$. As in conventional superconductors⁹ fluctuations are small when $j \gg \frac{3}{\text{corr}} = \frac{1}{T_{tr}^2} T_c^2$. Here

ℓ_{corr} is the correlation length for the random field $j_l(r)$, ℓ_{tr} a transport mean free path for the Fermi excitations. Assuming $\ell_{\text{corr}} = 500\text{\AA}$, $\ell_{\text{tr}} = 200\text{\AA}$, $\rho_0 = 20\text{\AA}$, one arrives at inequality $j \leq 1/30$. This estimation is not very accurate since the poorly known value of ℓ_{corr} enters it in the sixth power. More reliable estimation comes from the experimentally observed smearing of the specific heat jump¹¹, it renders $j \leq 1/25$ for the lower limit.

The order parameter of ABM-phase Eq. (1) does not satisfy the criterion (12). According to the above argument it can not be the order parameter of the A-like phase. Possible forms of the order parameters suitable for a description of that phase were discussed elsewhere^{6,7}. For the BW-order parameter criterion (12) is met. Identification of the B-like phase as BW-phase agrees with the recent pulse-NMR experiments and with the observation of homogeneously precessing domain (HPD) in that phase¹². Restriction imposed by criterion (12) becomes essential when B-phase is perturbed by magnetic field. Joint effect of aerogel and magnetic field on the BW-order parameter is discussed in the next section.

4. B PHASE IN MAGNETIC FIELD

In magnetic field two leading terms in the strength of the field have to be added to the free energy Eq. (4): linear in magnetic field and proportional to imaginary part of the tensor A_{ij} :

$$f_H^{(1)} = i e \sum_{ij} A_{ij} A_j H_i; \quad (14)$$

and quadratic in the field and proportional to real part of the same tensor

$$f_H^{(2)} = \frac{1}{2} (\sum_{ij} A_{ij} A_j + \sum_{ij} A_{ji} A_j) H_i H_j; \quad (15)$$

Derivation of criterion (12) is based on the invariance of GL free energy with respect to orbital rotations of the order parameter. The added terms preserve this property and as a consequence the criterion (12). This criterion imposes restriction on possible changes of the order parameter of B-phase. As in the case of pure ^3He ⁸ let us search the order parameter in magnetic field in a form of B_2 -phase (or oblate state):

$$A_{ij} = \frac{e^{i\phi}}{2} \left[(d + i e) (\hat{m}_j - i \hat{n}_j) + \sum_i (d - i e) (\hat{m}_j + i \hat{n}_j) + 2 \sum_i f_i \hat{l}_j \right] \quad (16)$$

Amplitudes u ; $u_{\#}$; $u_{\# \#}$ are found by minimization of the free energy. The order parameter (16) satisfies criterion (12) if

$$u_{\#}^2 = \frac{1}{2} (u^2 + u_{\# \#}^2) : \quad (17)$$

Separate analysis¹³ shows that the ansatz (16) does not impose additional restrictions on possible forms of the order parameter, provided a deviation from the bulk B-phase is small. Let us introduce following parametrization: $u + u_{\# \#} = 2$, $u - u_{\# \#} = 2$, then from eq. (17) $u_{\#}^2 = u^2 + u_{\# \#}^2$. Minimization of the free energy in a principal order on a small deviation renders:

$$= \frac{H}{2 [\beta_1 + \beta_3 + \beta_4 + \beta_5]} : \quad (18)$$

Coefficient β is proportional to the derivative of the density of states over energy, this makes β really small. Keeping only principal order correction we arrive for the order parameter to the expression:

$$A_j = R_j + i(e d - d e) R_j : \quad (19)$$

Eq. (19) shows that in a presence of a magnetic field produces a change of the BW-order parameter which is different from that in bulk ^3He . In the bulk the order parameter remains real and principal effect is suppression of $u_{\#}$ pairing amplitude while $u - u_{\# \#}$. In aerogel the increment of the order parameter is imaginary and principal change is appearance of a difference between u and $u_{\# \#}$. In both cases the increments are small and hardly observable in a moderate field, but there is an indirect manifestation of the induced change. Rotation matrix R_j entering Eq. (2) is often parametrized by an angle of rotation θ and direction of the rotation axis n . The angle θ is found by minimization of the dipole energy⁸:

$$f_D = \frac{3}{5} g_D (A - A_{jj} + A_j A_j) : \quad (20)$$

while direction of n remains arbitrary. This degeneracy is lifted when B_2 -order parameter (16) is substituted in the dipole energy (20). For the order parameter (19) the relevant part of the dipole energy is

$$f_n = (9/2) g_D n_z^2 : \quad (21)$$

It has a minimum when $n_z = 0$, i.e. when $n \perp H$ in a contrast to the pure ^3He where the minimum configuration is $n \parallel H$. A strength of the orientational effect is also different. Convenient characteristics of this strength is a healing length l_H i.e. a distance over which orientation of n recovers to its bulk

value being perturbed by for example a boundary of a sample. The healing length is found by comparison of the orientational energy with the gradient energy. For the orientational energy in aerogel eq. (21) one arrives at: $\ell_H = \frac{10^3}{5} \frac{D}{H}$. Dimensionless dipole constant (cf.⁸) $D = 5 \cdot 10^{-10}$. Coefficient can be evaluated from the splitting of $A_1 - A_2$ transition in pure ^3He as 10^{-3} 1/koe. For $H = 1$ koe and not too close to T_c healing length ℓ_H is of a few millimeters. This distance is comparable with dimensions of samples used in most experiments. It means that orientation of n is determined by geometry of samples and not by the field.

5. CONCLUSIONS

Properties of superfluid phases of ^3He in aerogel to a great extent are determined by the spatial fluctuations of the order parameter induced by aerogel. This contrasts with the pure superfluid ^3He where only average value of the order parameter is important. In a presence of aerogel contribution of fluctuations to physical quantities remains small only for a special class of order parameters. Formal criterion for selection of such phases in a vicinity of T_c follows from very general assumptions about properties of aerogel. The derived criterion rules out ABM-order parameter as a possible candidate for A-like phase and imposes restriction on a possible variation of BW-order parameter in magnetic field. Because of generality of the proposed scheme it may be applied for description of the influence of a quenched disorder on the other ordered systems with a continuous degeneracy.

6. ACKNOWLEDGMENTS

The author is grateful to V.V. Dmitriev, L.A. Melnikov and S.B. Savin for useful discussions and comments. This research was supported in part by RFBR grant (no. 01-02-16714) by Ministry of Education and Science of Russian Federation and by NATO grant SA (PST CLG .979379)6993/FP.

REFERENCES

1. J.V. Porto, J.M. Parpia, Phys. Rev. Lett., 74, 4667, (1995).
2. B.I. Barker, Y. Lee, L. Polikhina et al. Phys. Rev. Lett., 85, 2148 (2000).
3. G. Gervais, T.M. Haard, R. Nomura, N. Mulders, and W.W. Halperin, Phys. Rev. Lett., 85, 035701-1 (2001).
4. D. Rainer and M. Vuorio, J. Phys. C: Solid State Phys., 10, 3093 (1977).

5. I.A. Fomin, *Pisma Zh. Eksp. Teor. Fiz.*, 77, 285 (2003) [*JETP Lett.* 77, 240 (2003)].
6. I.A. Fomin, *J. Low Temp. Phys.*, 134, 769 (2004).
7. I.A. Fomin, *Zh. Eksp. Teor. Fiz.*, 125, 1115 (2004) [*JETP*, 98, 974 (2004)].
8. D. Vollhardt and P. Wolfe, *The Superfluid Phases of Helium 3*, Taylor and Francis (1990).
9. A.I. Larkin and Yu.N. Ovchinnikov, *Zh. Eksp. Teor. Fiz.*, 61, 1221 (1971) [*Sov. Phys. JETP*, 34, 651 (1971)].
10. Y. Imry and S.M. Ma, *Phys. Rev. Lett.*, 35, 1399 (1975)
11. J. He, A.D. Corvin, J.M. Parpia and J.D. Reppy, *Phys. Rev. Lett.*, 89, 115301 (2002)
12. V.V. Dmitriev, V.V. Zavjalov, D.E. Zmeev, I.V. Kosarev, and N. Mulders, *Pisma Zh. Eksp. Teor. Fiz.*, 76, 371 (2002) [*JETP Lett.* 76, 312 (2002)].
13. S.B. Savin, *Diploma thesis*, P.L. Kapitza Institute, Moscow, 2004.