

Morphological transition between diffusion-limited and ballistic aggregation growth patterns

S. C. Ferreira Jr.

Departamento de Física, Universidade Federal Viçosa, 36571-000, Viçosa, MG, Brazil

S. G. Alves, A. Faissal Brito, and J. G. Moreira

Departamento de Física, Universidade Federal de Minas Gerais,

CP 702, 30161-970, Belo Horizonte, MG, Brazil

In this work, the transition between diffusion-limited and ballistic aggregation models was revisited using a model in which biased random walks simulate the particle trajectories. The bias is controlled by a parameter β , which assumes the value $\beta = 0$ (1) for ballistic (diffusion-limited) aggregation model. Patterns growing from a single seed were considered. In order to simulate large clusters, a new efficient algorithm was developed. For $\beta \neq 0$, the patterns are fractal on the small length scales, but homogeneous on the large ones. We evaluated the mean density of particles $\bar{\rho}$ in the region defined by a circle of radius r centered at the initial seed. As a function of r , $\bar{\rho}$ reaches the asymptotic value $\rho_0(\beta)$ following a power law $\bar{\rho} = \rho_0 + A r^{-\alpha}$ with a universal exponent $\alpha = 0.46(2)$, independent of β . The asymptotic value has the behavior $\rho_0 \propto |\beta|^{-\gamma}$, where $\gamma = 0.26(1)$. The characteristic crossover length that determines the transition from DLA- to BA-like scaling regimes is given by $l_c \propto |\beta|^{-\delta}$, where $\delta = 0.61(1)$, while the cluster mass at the crossover follows a power law $M \propto l_c^{\nu}$, where $\nu = 0.97(2)$. We deduce the scaling relations $\alpha = \delta$ and $\gamma = 2$ between these exponents.

PACS numbers: 61.43.Hv, 05.40.Fb, 47.53.+n, 47.54.+r

I. INTRODUCTION

The pattern formation in nonequilibrium processes is a longstanding problem in Statistical Physics [1, 2, 3]. In special, the diffusion-limited aggregation model (DLA) [4] is a noteworthy example in which a very simple algorithm generates disorderly fractal clusters. This model was related to several physical and biological applications, such as electrodeposition [5], viscous fingering [6], bacterial colonies [7], and neurite formation [8]. In DLA model, particles released at a point distant from the cluster execute random walks until they find a neighbor site of the cluster and irreversible stick in this site. If the random walks are replaced by ballistic trajectories at random directions, we have the ballistic aggregation (BA) model [9]. In contrast to DLA, the BA model generates disordered nonfractal clusters with nontrivial scaling properties [2, 10].

Due to its importance as a fundamental model, several generalizations of DLA model were proposed [1, 2]. In particular, those models in which the particle trajectories are biased random walks were investigated [11, 12, 13, 14, 15, 16]. In these models, on the short length scales the particle trajectories are common random walks with fractal dimension 2.0, whereas on the longer length scales the bias becomes dominant and the dimensionality of the walk is 1.0. Clusters grown using this type of walk must behave like DLA model on the short length scales, while nonfractal patterns are ob-

served on the longer ones. Consequently, the mass of a cluster of size l is given by

$$M(l) = l^{d_f} f(l/l_c); \quad (1)$$

in which

$$f(x) = \begin{cases} \text{const} & \text{if } x \ll 1 \\ x^{d-d_f} & \text{if } x \gg 1 \end{cases} \quad (2)$$

Here, d is the space dimension, d_f is the DLA fractal dimension, and l_c is the crossover radius from DLA- to BA-like scaling regimes. This idea was firstly considered by Meakin [11]. In his model, the simulations start with a single seed at the center of a square lattice and the drift of all trajectories is in a fixed lattice direction. Along the walk, the particle is moved one lattice unity in the drift direction with probability P , or moves to one of its four next-neighbor sites with probability $1 - P$. The model generates patterns with a growth tendency in the opposite direction of the drift. The author argues that the crossover from the DLA-like structure on the short length scales to a compact structure on the longer ones is characterized by a length $l_c \propto P^{-1}$. However, using a renormalization group approach, Nagatani found $l_c \propto P^{1/(d-d_f)}$ [17]. Kim et. al. [12, 13] studied lattice models with a global drift to the seed, in which the particles have a higher probability to move to the nearest neighbor representing the shortest distance away from the seed. The pattern morphologies are ruled by the lattice anisotropy and their fractal dimension is 1. Nagatani [14] considered the effects of positive and negative radial drifts in DLA model. In the positive case the cluster fractal dimension is asymptotically 2, while

Electronic address: silviojr@ufv.br

eccentric patterns with dimension 1 were found for the negative case. Other models [15, 16] consist of the deposition processes on a d-dimensional substrate in which the walk drift is the substrate direction.

In the present paper, we are interested in the transition from DLA to BA models when the random or ballistic trajectories of DLA and BA models, respectively, are replaced by biased random walks with a random drift direction. The central concern of this work is the fact that all real fractals exhibit scaling only on limited ranges and, consequently, a quantitative analysis of both experiments and simulations demands a deep understanding of these crossovers. The outline of the paper is the following. In Sec. II, the model and the respective computational algorithm are described. In Sec. III, the simulation results are presented and discussed. Finally, some conclusions are drawn in Sec. IV.

II. MODELS AND METHODS

As in the original DLA and BA models, at the beginning of the simulations a unique seed localized in the center of the lattice constitutes the cluster. Then, particles are sequentially released at a circle distant from the cluster and execute biased random walks. The distance between the center of the lattice and the launching circle is denoted by R_1 . The biased walks are defined by

$$\begin{aligned} x_{n+1} &= x_n + \cos(\theta' + \theta_n) \\ y_{n+1} &= y_n + \sin(\theta' + \theta_n); \end{aligned} \quad (3)$$

where x_n and y_n are the particle coordinates at the n th step of the walk, θ' is a random angle that defines the bias direction, $\theta \in [0; 2\pi]$ is the parameter that controls the random component of the trajectories, and θ_n is a random direction. The variables θ' and θ_n are in the range $[0; 2\pi]$. Notice that θ' is defined at the beginning of the walks, whereas θ_n assumes random values for each walk step. One can see that the particular cases $\theta = 0$ and $\theta = 1$ recover the BA and DLA models, respectively. If the particle visits a site neighboring the cluster it irreversibly joins to this site. However, if the distance between the particle and the cluster is too large, i. e., larger than a killing radius R_k , the particle is excluded and a new one is released at the launching circle. In order to determine when a walker is neighboring a cluster site, its lattice position was defined as the next integer value of its real coordinates defined by Eq. (3). The majority of the results presented in this work refer to the square lattice version of the model.

The previously introduced variables (R_1 and R_k) must be as large as possible. However, computational limitations restrict the use of R_1 and R_k values. We defined $R_1 = R_{\max} + R_0$, where R_{\max} is the maximum distance from the center of the lattice of a particle belonging to the cluster. For DLA and BA models R_0 can be of the order of some lattice units [1, 2]. However, for the model




FIG. 1: Schematic representation of the model.

with biased random walks the patterns morphologies are strongly dependent on this value. Our tests suggest that the patterns become insensitive to R_0 variations when $R_0 > 300$, in agreement with the values adopted by Kim [16] for a model of deposition of biased random walks on a substrate. Thus, $R_0 = 400$ were used in all simulations. The killing radius R_k must be 10 to 100 times R_{\max} for very large DLA clusters [1] whereas a R_k only some lattice units larger than R_{\max} is necessary for the BA model. Due to the bias present in the random walks, we used the same strategy adopted by Kim [16], i. e., $R_k = 2R_{\max} + R_0$. Fig. 1 illustrates two tentatives, one successful and the other frustrated, to add a new particle to the cluster.

To analyze the transition between BA and DLA it is necessary to simulate large clusters using lattices containing up to $10^4 \times 10^4$ sites, specially when $\theta = 1$. Consequently, the computational time becomes prohibitive and an efficient algorithm is necessary. A technique commonly used to simulate large DLA and related models is to allow the particles execute long steps at random directions if they are far from the cluster [18, 19, 20, 21]. This procedure is correct because the probability of a random walker crosses the circle centered on its initial position in a given angle is uniformly distributed in the interval $[0; 2\pi]$. However, for the biased random walkers this is not true. Indeed, the probability density distributions are concentrated around the direction θ' . For θ' not very close to the unity or for large steps, the probability distributions are very well fitted by Gaussian curves centered at θ' , as illustrated in Fig. 2. Using this fact, the following procedure was adopted. If the distance between the random walker and the cluster is larger than a value $R_s + R_k$, it executes a jump of length R_s . Thus, a long




FIG. 2: Angle distributions for biased random walkers with preferential direction $\alpha = 0$ and the correspondent Gaussian fits. The curves correspond to $\alpha = 0.99$ and $R_s = 200$ (squares), $\alpha = 0.99$ and $R_s = 1000$ (triangles), and $\alpha = 0.90$ and $R_s = 200$ (circles). Other parameter sets used in the simulations provide fits better than those shown in this figure. For all couples α and R_s used in the simulations a correlation coefficient $r^2 = 0.999$ was required.

jump can not lead the walker to a distance smaller than ℓ . The jump direction is $\theta = \theta' + \phi_g$, where ϕ_g is a random number between $-\pi$ and π selected from a Gaussian distribution

$$P(\phi_g) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{\phi_g^2}{2\sigma^2}\right) \quad (4)$$

In order to obtain the Gaussian width σ for each couple α and R_s , a large number of biased random walks (10^6 – 10^7) were simulated and a histogram of the probabilities built (Fig. 2). Then, the σ value can be determined using least square Gaussian fits. The quality of fits is improved as larger R_s values and smaller α values are used. Therefore, the R_s values should be sufficiently large for reproduce good fits, in special for $\alpha \rightarrow 1$. Furthermore, several R_s values can be used in a same simulation improving the algorithm efficiency. We used two values: $R_s = 200$ and $R_s = 1000$. The σ values used in the simulations are shown in table I. Also, $\ell = 20$ were used. All tests show that the growth patterns are not sensible to the ℓ value. It is worth to note that the Gaussian distribution is not normalized in the interval $[-\pi; \pi]$ and, obviously, this is not the actual angle distribution for the present problem. However, the very good fits to the angle distributions justify the use of the Gaussian functions.

III. RESULTS AND DISCUSSIONS

The first stage of the present work was to confirm the validity of the previous defined algorithm. We simulate

| | $(R_s = 200)$ | $(R_s = 1000)$ |
|-------|---------------|----------------|
| 0.100 | 0.02585 | 0.01202 |
| 0.300 | 0.07591 | 0.03413 |
| 0.500 | 0.12535 | 0.05613 |
| 0.700 | 0.18206 | 0.08139 |
| 0.900 | 0.31933 | 0.14216 |
| 0.950 | 0.45109 | 0.20057 |
| 0.990 | 1.03575 | 0.45130 |
| 0.995 | 1.53278 | 0.63451 |

TABLE I: σ values determined with the Gaussian least square fits.

relatively small lattices containing 10^3 – 10^5 sites with and without the optimization for $\alpha = 0.99$. In Fig. 3, comparisons between clusters generated with (top) and without (bottom) the optimization are shown. Comparing the patterns, one can see that they are statistically indistinguishable. Using the mass-radius method, the fractal dimensions of the patterns generated with and without the algorithm were $d_f = 1.70 \pm 0.02$ and $d_f = 1.72 \pm 0.02$, respectively, and the exponents of the radius of gyration, defined by $R_g \propto n^\nu$ (n is the number of cluster particles) [2], were $\nu = 0.560 \pm 0.003$ and $\nu = 0.561 \pm 0.002$, respectively. These exponents reinforce the algorithm validity. Concerning the computational time, a single run to generate one of the clusters shown in Fig. 3 without the optimization takes about 1 hour in a 3 GHz Pentium IV, but the same simulation is done in 10 minutes using the optimization. Therefore, even for small lattices the simulation performance is greatly improved when our optimized algorithm is used.




FIG. 3: Two clusters generated (a) using or (b) not the optimization. In these simulations, lattices of size $L = 1000$ and $\alpha = 0.99$ were used.




FIG. 4: Morphology transition between BA and DLA growth patterns. The number of aggregated particles varies from $2.5 \cdot 10^5$ ($\lambda = 0.1$) to $5 \cdot 10^4$ ($\lambda = 0.995$).

Figure 4 shows growth patterns for distinct λ values. These patterns were generated with the optimization at lattices containing 10^3 – 10^5 sites. The simulations stopped when the aggregate reaches the lattice edge. A continuous transition from disordered and dense to ramified clusters is observed. For small λ values the patterns are essentially BA-like but, the patterns become very similar to the DLA clusters as $\lambda \rightarrow 1$. Indeed, the cluster generated with $\lambda = 0.995$ is characterized by the square lattice anisotropy, a signature of the DLA model [1, 2]. However, one expects that all patterns become asymptotically homogeneous with a finite characteristic size for the empty regions.

In order to quantify the DLA to BA morphology transition, the mean particle density in the inner regions of the cluster was evaluated. This mean density $\bar{\rho}(r)$ is defined as the ratio between the number of occupied sites and the total number of sites in a region delimited by a circle of radius r centered at the initial seed. Since one expects asymptotically nonfractal clusters, the density must reach a finite value ρ_0 as $r \rightarrow \infty$. Nevertheless, the approach to the constant density is very slow and takes a scale invariant form

$$\bar{\rho}(r) = \rho_0 + A r^{-\alpha} \quad (5)$$

Here, α is a correction to the fractal dimension and A a constant. This scaling hypothesis was also used by Liang and Kadano [10] to study the driven ballistic aggregation, in which the particles trajectories are in a single direction. They conclude that the α exponent is nonuniversal, i.e., depends on the lattice structure.

In Fig. 5, the double-logarithm plots of $\bar{\rho} - \rho_0$ as a function of r for distinct λ values are shown. The density ρ_0 was obtained by searching for the best linear fit




FIG. 5: Double-logarithm plots of $\bar{\rho} - \rho_0$ against r for $\lambda = 0.3$ (squares) and $\lambda = 0.99$ (circles). The dashed line corresponds to the slope -0.46 . The linear fits of the data provide -0.45 for $\lambda = 0.3$, and -0.47 for $\lambda = 0.99$.

in the larger linear region. To avoid the active region, we limited the fits to those data corresponding to a half of the cluster sizes. Depending on the λ value, lattices with linear size $L = 5 \cdot 10^3$ or $L = 10^4$, and 10 to 20 independent runs were used. One can observe a power law regime for $r > 10$ showing that the approach to the stationary value obeys Eq. 5. In Fig. 6, the asymptotic density ρ_0 and the α exponent are shown as a function of $1 - \lambda$, the distance from the transition point. ρ_0 acts as an order parameter, which vanishes at the critical point following a relation $\rho_0 \sim |1 - \lambda|^j$. The exponent obtained from the data of Fig. 6(a) was $j = 0.27(1)$, whereas the

exponents obtained for $L = 2 \cdot 10^3$ and $L = 10^4$ were $\gamma = 0.28(2)$ and $\gamma = 0.26(1)$, respectively. The numbers in parenthesis represent the uncertainties. In Fig. 6(b), the exponents for distinct values are shown. One can observe that γ fluctuates around the value 0.46. The smaller value found was 0.43 and the larger one 0.48. Our simulations suggest that the exponent is independent of λ and its value is $\gamma = 0.46 \pm 0.02$. The error indicated in the γ value was evaluated through an average over the data of Fig. 6(b).

In order to test the universality of the γ and ρ_0 exponents we studied two versions of the present model. In the first one, we used square lattices, but the walkers stick to the cluster if they reach a nearest or a next-nearest empty neighbor of an occupied site. In the second one, we used a hexagonal lattice. Lattices with size $L = 2000$ were used. The exponents for the first modified version were $\gamma = 0.27 \pm 0.02$ and $\gamma = 0.49 \pm 0.03$, and the exponents for hexagonal lattice were $\gamma = 0.28 \pm 0.02$ and $\gamma = 0.49 \pm 0.03$. These results lead us to conclude that these exponents are universal.




FIG. 6: (a) Stationary density ρ_0 and (b) exponent defined in Eq. (5) as a function of the distance from the transition point. The lattice size used was $L = 5 \cdot 10^3$ and simulation stopped when the cluster reaches the lattice edge.

The morphological transition between BA and DLA models was also characterized by the crossover radius defined in Eqs. (1) and (2). The number of particles $M(r)$ inside a region delimited by a circle of radius r centered at origin was evaluated. The M vs: r curves exhibit tenuous crossovers determining the transition between DLA- and BA-like scaling regimes. In Fig. 7(a), an example of this crossover is shown. Since the growth patterns scale as DLA (BA) for small (large) length scales, in order to evaluate the crossover we fitted the curves by power laws $M(r) \propto r^{d_f}$, where $d_f = 1.71$ and $d_f = 2$ were used for the initial and the final curve regions, respectively. The crossover lengths obtained through this method are drawn as a function of the distance from the transition point in Fig. 7(b). The length diverges

at $\lambda = 1$ following a power law $\xi \propto (1-\lambda)^{-\beta}$, where $\beta = 0.61(1)$. Moreover, the mass at the critical point diverges as $M_c \propto (1-\lambda)^{-\alpha}$, where $\alpha = 0.97(2)$.

As discussed in section I, these crossovers between fractal and homogeneous patterns occur due the crossover in the particle trajectories. However, the crossover length of the walker trajectories is given by (see appendix)

$$\xi_w = \frac{r}{\sin(\theta)} \propto \frac{\sin(\theta)}{\sin(\theta)}; \quad (6)$$

which diverges as $\xi_w \propto (1-\lambda)^{-\beta}$ for $\lambda \rightarrow 1$. Thus, although the transition between DLA and BA models is due to the transition in the walk dimensionality, the respective crossover lengths are not proportional.




FIG. 7: (a) Determination of the crossover between DLA- and BA-like scaling regimes for $\lambda = 0.90$. The straight lines represent the slopes 1.71 and 2, i.e., the fractal dimensions for DLA and BA models, respectively. (b) The crossover length (squares) and the correspondent mass (circles) as a function of the distance from the transition point. These results were obtained for $L = 5 \cdot 10^3$.

Notice that Eq. (5) describes the mean density behavior when $r \gg \xi$. Moreover, $\rho \propto r^{-2}$ when $r \ll \xi$ due to the cluster fractality in this length scale. Thus, using Eq. (5), we found that the mean density at the crossover can be written as

$$\rho_c = A_1 \xi^{-d_f} + A_2 \xi^{-2}; \quad (7)$$

where A_1 and A_2 are constants. But, the mean density at the crossover is given by

$$\rho_c = \frac{M_c}{\xi^2} \propto \xi^{-2}; \quad (8)$$

Comparing Eqs. (7) and (8), we have that they are consistent only if

$$d_f = 2 \quad \text{and} \quad d_f = 2; \quad (9)$$

In agreement with the scaling relation (9), the number of independent exponents are reduced from 4 to 2. Using

the exponents measured for systems with size $L = 5000$, we found $\alpha = 0.28(2)$ and $2 - \alpha = 0.25(3)$, beside $\alpha = 0.27(1)$. The difference between these values is inside of the error margins indicated in the parenthesis. The large uncertainties obtained in the exponents (5% -10%) are originated in the difficulty in the determination of the exact crossover points.

IV. CONCLUSIONS

In the present work, we studied the transition between diffusion-limited aggregation (DLA) and ballistic aggregation (BA) models. We used a model in which the random walks in the DLA model are replaced by biased random walks with a drift in a random direction. The drift is controlled by a parameter $\alpha \in [0;1]$ that leads the model from BA ($\alpha = 0$) to DLA ($\alpha = 1$) (see Eq. (3)). Also, an efficient algorithm, which allows large scaling analysis of the growth patterns, was introduced.

For any bias, the clusters are fractal (DLA-like) on the short length scales whereas nonfractal patterns are obtained on the large ones. The transition between DLA- and BA-like scaling regimes is determined by a characteristic length that diverges as $\alpha \rightarrow 1$ following a power law $\xi \sim |1 - \alpha|^{-\nu}$, where $\nu = 0.61(1)$, while the cluster mass at the crossover follows the relation $M \sim \xi^d$, where $d = 0.97(2)$. This crossover was not numerically determined in similar previous works. The density in the inner regions of the cluster reaches an asymptotic value $\rho_0 \sim \xi^{-d}$, where $d = 0.26(1)$. However, this approach is slow and follows a power law decay with a universal exponent $\alpha = 0.46(2)$ independent on the drift. These exponents obey the scaling relations $\alpha + \nu = 1$ and $d = 2 - \alpha$.

It is worth to stress two main contributions of the present work. The first one is the development of an algorithm that can be used to study other models with biased random walks as for example those related to deposition processes [15, 16], for which the determination of universality classes is a hard work. The second one is the careful quantitative characterization of the transition between DLA and BA growth models that, in our knowledge, was not previously done. The understanding of these crossovers can be an essential tool in the analysis of real fractals, which always exhibit scaling on limited ranges.

Acknowledgments

We would like to thank M. L. Martins and A. P. F. Atman for the critical reading of the manuscript. This work was partially supported by the CNPq Brazilian agency.

*

APPENDIX A: DEMONSTRATION OF EQ. (6)

For sake of simplicity, we consider Eq. (3) with a drift direction $\theta = 0$ and $x_0 = y_0 = 0$. Iterating Eq. (3) for n steps, we found

$$hx_n i = \sum_{i=1}^{X^n} h\cos(\theta_i) i; \quad (A1)$$

and

$$hx_n^2 i = \sum_{i=1}^{X^n} \sum_{j=1}^{X^n} h\cos(\theta_i) \cos(\theta_j) i; \quad (A2)$$

But,

$$h\cos(\theta_i) i = \frac{1}{2} \sum_{j=1}^Z \cos(\theta_j) d = \frac{\sin(\theta)}{2}; \quad (A3)$$

and

$$h\cos(\theta_i) \cos(\theta_j) i = \frac{\sin(\theta)^2}{2} (1 - \delta_{ij}) + \frac{1}{2} \frac{\sin(2\theta)}{2} + \delta_{ij}; \quad (A4)$$

where δ_{ij} is the Kronecker delta.

Substituting Eqs. (A3) and (A4) in Eqs. (A1) and (A2), respectively, we found

$$hx_n i = n \frac{\sin(\theta)}{2} \quad (A5)$$

and

$$\sigma_x^2(n) = n \left(\frac{1}{2} \left(1 + \frac{\sin(2\theta)}{2} \right) - \frac{\sin(\theta)^2}{2} \right); \quad (A6)$$

where $\sigma_x^2(n) = hx_n^2 i - hx_n i^2$ is the variance of the coordinate x_n . With a similar analysis, we obtained

$$hy_n i = 0 \quad (A7)$$

and

$$\sigma_y^2(n) = n \frac{1}{2} \left(1 - \frac{\sin(2\theta)}{2} \right); \quad (A8)$$

Thus, the walk mean displacement $r(n) = \sqrt{hx_n^2 i + hy_n^2 i}$ and the variance $\sigma^2(n) = \sigma_x^2 + \sigma_y^2$ are given by

$$r(n) = n \frac{\sin(\theta)}{2} \quad (A9)$$

and

$$\sigma(n) = n^{1/2} \left(1 - \frac{\sin(\theta)^2}{2} \right)^{1/2}; \quad (A10)$$

The crossover of the walk dimensionality ($d = 2$ for short times and $d = 1$ for long ones) occurs when $r(n) \sim n^{-1}$. Making equal Eqs. (A 9) and (A 10), we obtained an estimate of the characteristic number of steps N necessary for the crossover

$$N = \frac{2}{\sin(\lambda)} - 1: \quad (\text{A } 11)$$

Thus, the characteristic crossover length is

$$\xi_w = r(N) = \frac{\sin(\lambda)}{\sin(\lambda)}: \quad (\text{A } 12)$$

In Fig. (8), ξ_w is plotted as a function of $1-\lambda$. Expanding Eq. (A 12) around $\lambda = 1$, we found $\xi_w \sim |1-\lambda|^{-1}$.




FIG. 8: Double-logarithm plot of the crossover length ξ_w as a function of $1-\lambda$. The solid line represents the data obtained from Eq. (A 12) and the dashed line has a slope -1 .

-
- [1] P. Meakin, *Fractals, scaling and growth far from equilibrium* (Cambridge University Press, Cambridge, 1998).
 - [2] T. Vicsek, *Fractal Growth Phenomena* (World Scientific, Singapore, 1992).
 - [3] A.-L. Barabasi and H. E. Stanley, *Fractal Concepts on Surface growth*, (Cambridge University Press, Cambridge, 1995).
 - [4] T. A. Witten and L. M. Sander, *Phys. Rev. Lett.* 47, 1400 (1981).
 - [5] M. Matsushita, M. Sano, Y. Hayakawa, H. Honjo, and Y. Sawada, *Phys. Rev. Lett.* 53, 286 (1984).
 - [6] K. J. Malý, J. Feder, and T. J. Ssang, *Phys. Rev. Lett.* 55, 2688 (1985).
 - [7] M. Matsushita and H. Fujikawa, *Physica A* 168, 498 (1990).
 - [8] F. Caserta, H. E. Stanley, W. D. Eldred, G. Daccord, R. E. Hausman, and J. Nittmann, *Phys. Rev. Lett.* 64, 95 (1990).
 - [9] M. J. Vold, *J. Colloid Sci.* 18, 684 (1963).
 - [10] S. Liang and L. P. Kadano, *Phys. Rev. A*, 31, 2628 (1985).
 - [11] P. Meakin, *Phys. Rev. B* 28, 5221 (1983).
 - [12] Y. Kim, K. R. Choi, and H. Pak, *Phys. Rev. A* 45, 5805 (1992).
 - [13] Y. Kim and K. R. Choi, *Phys. Rev. E* 48, 1586 (1993).
 - [14] T. Nagatani, *Phys. Rev. A* 39, 438 (1989).
 - [15] M. Castro, R. Cuemo, A. Sanchez, and F. Domínguez-Adame, *Phys. Rev. E* 62, 161 (2002).
 - [16] Y. Kim, *J. Korean Phys. Soc.* 30, 511 (1997).
 - [17] T. Nagatani, *Phys. Rev. A* 37, 3514 (1988).
 - [18] R. C. Ball and R. M. Brady, *J. Phys. A* 18, L809 (1985).
 - [19] P. Meakin, *J. Phys. A* 18, L661 (1985).
 - [20] S. Tolman and P. Meakin, *Phys. Rev. A* 40, 428 (1989).
 - [21] S. C. Ferreira Jr., *Eur. Phys. J. B* 42, 263 (2004).