

Quantum fluctuations of the ultracold atom-molecule mixtures

T. Domanski

Institute of Physics, M. Curie Skłodowska University, 20-031 Lublin, Poland

We investigate evolution of the quantum coherence in the ultracold mixture of fermionic atoms and bosonic diatomic molecules. Interactions are there experimentally controlled via tuning the external magnetic field. Consequently, the fermionic atoms and their bosonic counterparts can be driven to a behavior resembling the usual BCS to BEC crossover. We analyze in some detail how this quantum coherence evolves with respect to time upon a smooth and abrupt sweep across the Feshbach resonance inducing the atom-molecule quantum fluctuations.

PACS numbers: 74.20.Mn, 03.75.Kk, 03.75.Ss

ATOM SUPERFLUIDITY

The recent experimental techniques for trapping and cooling of the atomic vapors enabled exploration of the extremely low temperature regions where quantum effects play a crucial role. An example can be the Bose-Einstein condensation (BEC) produced out of the bosonic atoms in alkali metals, polarized hydrogen, etc. Phase transition to the BEC state is there triggered purely by the quantum statistical requirements which lead to macroscopic occupancy of the lowest energy level and can occur even in absence of any interactions. Recent activities in the field of ultracold atomic systems focus on application of similar techniques to fermionic atoms like ^6Li or ^{40}K (besides even number of nucleons they consist of odd number of electrons). At ultralow temperatures such quantum effects like the Pauli principle play considerable role, but eventual quantum phase transitions would be allowed only if fermionic atoms get correlated via interactions.

Interactions between trapped atoms are routinely induced by applying the magnetic field to fermion atoms prepared in several (two or more) hyperfine configurations. From elementary considerations [1] it turns out that the involved hyperfine states experience the effective scattering described by a potential whose magnitude and sign depend on the applied field B . In particular, the various (so called) Feshbach resonances can take place. On this basis there was proposed a mechanism of the resonance superfluidity [2] with a transition occurring near the Fermi temperature $T_c = T_F$. Besides the isotropic phase there has already been observed also the exotic p-wave superfluidity [3, 4].

A unique possibility of controlling the effective interactions gives a chance for the experimental realization of the BCS to BEC crossover. The BCS limit corresponds to a case of weakly attracting fermion atoms which get coupled into the large Cooper pairs. In opposite limit, the tightly bound diatomic molecules are formed which ultimately can undergo transition to the BEC. Experimentalists are able to switch between these limits in a controllable manner and change of the interactions can

be performed either adiabatically (by slowly varying the field) [5] or in a non-adiabatic way (via the sudden sweep) [6].

In this short paper we investigate the quantum fluctuations induced by the time-dependent change of the interactions. We focus on a situation when the magnetic field is detuned from the resonant value B_0 towards the far BCS regime at higher field $B > B_0$. We consider two different processes: the smooth and the sudden switching. The fast sweep has been discussed in the literature but with some ambiguous conclusions concerning evolution of the order parameters with respect to time [7, 8, 9]. From our analysis we find that both parameters do oscillate in a damped way.

HEISENBERG EQUATIONS

In a close proximity to the Feshbach resonance (i.e. when $B = B_0$) the ultracold fermion atoms coexist and interact with the diatomic molecules. On a microscopic basis this situation can be described in terms of the two component boson-fermion Hamiltonian [2]

$$H = \sum_{\mathbf{k}} \left(\epsilon_{\mathbf{k}}^F + \frac{g}{N} \sum_{\mathbf{q}} b_{\mathbf{q}}^\dagger c_{\mathbf{k}-\mathbf{q}}^\dagger + \text{h.c.} \right) c_{\mathbf{k}} + \sum_{\mathbf{q}} \left(\epsilon_{\mathbf{q}}^B + 2(B - B_0) \right) b_{\mathbf{q}}^\dagger b_{\mathbf{q}} \quad (1)$$

which has been known and studied in the solid state physics by J. Ranninger and coworkers [10] as a phenomenological model for the high temperature superconductivity. In the present context (1) describes the atoms in two hyperfine states denoted symbolically by \uparrow and \downarrow . The second quantization operators $c_{\mathbf{k}}^{(\uparrow)}$, $b_{\mathbf{q}}^{(\uparrow)}$ correspond to fermion atoms with energy $\epsilon_{\mathbf{k}}^F = \hbar^2 k^2 / 2m$ and to diatomic molecules with energy $\epsilon_{\mathbf{q}}^B = \hbar^2 q^2 / 2(2m)$. The effect of external magnetic field is included via the detuning parameter $B - B_0$ which shifts the boson energies and hence affects efficiency of the boson-fermion coupling g [11]. As usually g is the common chemical potential and we use the grand canonical ensemble

to ensure the conservation of the total particle number $\sum_k c_k^\dagger c_k + 2 \sum_q b_q^\dagger b_q$.

We are interested here in studying the time dependent evolution of fermion and boson occupancies together with the corresponding order parameters. For this purpose we derive the Heisenberg equations of motion which for the Hamiltonian (1) are given by

$$i \frac{\partial c_{k\#} c_k}{\partial t} = (k + q - k) c_{k\#} c_k + g b_q \sum_{q^0} b_{q^0} c_{q^0}^\dagger c_{k\#} c_k + c_{k+q^0}^\dagger c_{q^0} c_k ; \quad (2)$$

$$i \sum_k \frac{\partial c_k^\dagger c_k}{\partial t} = 2g \sum_q b_q c_k^\dagger c_{k\#} c_k + b_q^\dagger c_{k\#} c_k ; \quad (3)$$

$$i \frac{\partial b_q}{\partial t} = E_q b_q + g \sum_k c_{k\#} c_k ; \quad (4)$$

$$i \frac{\partial b_q^\dagger b_q}{\partial t} = g \sum_k b_q^\dagger c_{k\#} c_k - b_q c_k^\dagger c_{k\#} ; \quad (5)$$

where $k = \frac{E_k}{\hbar}$, $E_q = \hbar^2 (B - 2) q^2$ and we set $\hbar = 1$. In general equations (2-5) are not solvable exactly. In the next section we briefly discuss an approximate method which shall be valid for the ground state and for very low temperatures.

THE SINGLE MODE APPROACH

For temperatures close to the absolute zero we can neglect the excited (finite momentum) boson states. It is sufficient to restrict attention to the $q=0$ boson level because it is macroscopically occupied. In such single mode approach [7, 8] the initial Hamiltonian (1) reduces to

$$H = \sum_k c_k^\dagger c_k + \sum_q E_0 b_q^\dagger b_q + \frac{g}{N} \sum_k b_0 c_{k\#} c_k + c_k^\dagger c_{k\#} b_0 ; \quad (6)$$

Following Anderson [12] we introduce the pseudospin notation

$$\begin{aligned} c_{k\#} c_k &= \frac{1}{2} (S_k^x + i S_k^y) \\ c_k^\dagger c_{k\#} &= \frac{1}{2} (S_k^x - i S_k^y) \\ c_k^\dagger c_k &= \frac{1}{2} (S_k^z + 1) \\ c_{k\#} c_{k\#} &= \frac{1}{2} (S_k^z - 1) \end{aligned}$$

such that $S_k = \frac{1}{2} (S_k^x, S_k^y, S_k^z)$ and $S_k = \frac{1}{2} S_k$; S_k are the usual Pauli operators. In the single mode approach we rewrite the Heisenberg equations (2-5) using the pseudospin notation

$$i \frac{\partial S_k^z}{\partial t} = 2 S_k^x + g b_0 S_k^z ; \quad (7)$$

$$i \frac{\partial S_k^z}{\partial t} = 2g b_0^\dagger S_k^x b_0 ; \quad (8)$$

$$i \frac{\partial b_0}{\partial t} = E_0 b_0 + g \sum_k S_k^x ; \quad (9)$$

$$i \frac{\partial b_0^\dagger b_0}{\partial t} = g \sum_k b_0^\dagger S_k^x b_0 ; \quad (10)$$

which are identical with expressions (5) and (6) in the Ref. [8]. One next replaces the boson operators by their time-dependent expectation values $b(t) = \langle b_0 \rangle$ and $b^\dagger(t) = \langle b_0^\dagger \rangle$.

In the stationary case when all parameters in (6) are time independent we can derive various expressions for the static expectation values [10]. Hamiltonian (6) has formally the following structure

$$H = \sum_k \tilde{\hbar}_k S_k^z + \text{const}; \quad (11)$$

so the pseudospin S_k behaves as though affected by a fictitious magnetic field $\tilde{\hbar}_k = (\hbar_0^x; \hbar_0^y; \hbar_k^z)$ where $\hbar_0^x + i \hbar_0^y = g \langle b_0 \rangle$. Following Anderson [12] we can solve this problem (11) for arbitrary temperature. In analogy to the Weiss theory of ferromagnetism we obtain that a magnitude of the pseudospin expectation value is $\langle S_k^z \rangle = \tanh \left(\frac{\hbar_k^z}{\hbar_k^2 + \hbar_0^2} \right) = \tanh \left(\frac{\hbar_k^z}{2k_B T} \right)$. Determining the angle between the z and xy components of the vector $\tilde{\hbar}_k$ we finally arrive at the stationary equations [10]

$$\begin{aligned} \langle S_k^x \rangle &= \langle \hbar_k^x S_k^z \rangle \\ &= \frac{g}{2} \frac{\hbar_0^x}{\hbar_k^2 + \hbar_0^2} \tanh \left(\frac{\hbar_k^z}{2k_B T} \right) \end{aligned} \quad (12)$$

and

$$\begin{aligned} \langle S_k^y \rangle &= \frac{1}{2} \sum_k \langle \hbar_k^y S_k^z \rangle \\ &= \frac{g}{2} \frac{\hbar_0^y}{\hbar_k^2 + \hbar_0^2} \tanh \left(\frac{\hbar_k^z}{2k_B T} \right) ; \end{aligned} \quad (13)$$

FLUCTUATIONS OF THE ORDER PARAMETERS

In the symmetry broken state (for $T < T_c$) the two component model (1) is characterized by two order parameters: $b(t)$ and another one of the fermion subsystem defined as $\langle S \rangle = \sum_k \langle S_k^z \rangle$. These quantities are complex. In the stationary case they are proportional to each other as can be seen from the equation (12). However, this relation is no longer valid when the Hamiltonian (6) depends on time. Evolution of the order parameters $b(t)$ and $\langle S \rangle(t)$ with respect to time must be determined by solving the Heisenberg equations (7-8) subject to some boundary conditions.




FIG. 1: Variation of the fermion $j(t)$ and boson $b(t)$ order parameters caused by the detuning from the Feshbach resonance. Initially, for $t < 0$, the system is in the stationary state with $\langle j(t) \rangle = \langle b(t) \rangle = 0$ where the magnitudes of both order parameters are static. Upon lifting the molecule level there appear the damped oscillations. In the left h.s. panel we illustrate the case of an abrupt detuning ($t > 0$) = 0.1g and in the right h.s. a smooth detuning (t) / t .

We analyze here such dynamics assuming that initially, for $t = 0$, the mixed atom molecule system is exactly on the Feshbach resonance $\mu = 0$. We also assume that the boson order parameter is real and $b(t = 0) = 1$ so that the fermion order parameter is real too. Value of $\mu(t = 0)$ was determined from the equation (12). For simplicity we focus on the ground state and set the boson fermion coupling g as a unit or all the energies appearing in our study.

For time $t > 0$ we change the detuning parameter in the following ways: a) via the sudden detuning as previously discussed in the Refs [7, 8, 9] and b) through gradually increasing $\mu(t) = \mu_0 + \mu_1 t$. Avoiding any constraint solutions we solved numerically the Heisenberg equations (7,8) by means of the Runge-Kutta algorithm.

By increasing the boson fermion system is pushed to the far BCS regime. For $\mu = 0.1g$ at $t = 1$ the order parameters $b(t)$ and $j(t)$ should reach very small values. Figure 1 shows that indeed temporal evolution occurs into such asymptotics and practically is achieved after several oscillations. In both cases the oscillations are clearly damped in agreement with the previous study by K. Bumet and coworkers [9]. However, this process of damping is sensitive on a particular profile of the time dependent detuning. This can be seen from the figure 1 and also in the next figure 2, where we plot the phase of the boson order parameter $b(t) = |b(t)|e^{i\theta(t)}$. For a smooth switching the oscillations seem not to look regular at all.




FIG. 2: Evolution of the phase of the boson order parameter $b(t)$ with respect to time t for: an abrupt detuning (left), and for a smoothly increasing detuning ($\mu(t) = \mu_0 + \mu_1 t$) (right). Instead of the bare angle we plot the function $\sin \theta = \text{Im } b(t) = \text{Im } j(t)$.

SUMMARY

We studied the dynamics of the ultracold fermion atoms upon the sudden and gradual detuning from the Feshbach resonance. Such situation can be experimentally realized by switching the external magnetic field from B_0 to the higher values. From the selfconsistent numerical solution of the equations of motion we find that the order parameters start oscillating with the amplitude decaying in time. Such damped oscillations depend on the specific form in which the detuning $\mu(t)$ is carried out. We also remark that the quantum oscillations of the order parameters turn out to be damped even on the level of the single mode approach without taking into account scattering to the infinite boson momenta.

ACKNOWLEDGEMENT

This work is supported by the Polish Committee of Scientific Research under the grant No. 2P 03B 06225.

-
- [1] E. T. Immanuel, P. Tommasini, M. Hussein and A. Kerman, Phys. Rep. 315, 199 (1999).
 - [2] M. Holland, S.J.J.M.F. Kokkekmans, M.L. Chiofab and R. Walser, Phys. Rev. Lett. 87, 120406 (2001).
 - [3] C. Ticknor, C.A. Regal, D.S. Jin, and J.L. Bohn, Phys. Rev. A 69, 042712 (2004).
 - [4] E. Gubankova, E.G. Mishchenko, and F. Wilczek, Phys. Rev. Lett. 94, 110404 (2005); Y. Ohashi, Phys. Rev. Lett. 94, 050403 (2005).
 - [5] M. Greiner, C.A. Regal, and D.S. Jin, Nature 426, 537 (2003).
 - [6] M. Holland and S.J.J.M.F. Kokkekmans, Phys. Rev. Lett. 89, 180401 (2002).
 - [7] A.V. Andreev, V. Gurarie, and L. Radzihovsky, Phys. Rev. Lett. 93, 130402 (2004).
 - [8] R.A. Barankov and L.S. Levitov, Phys. Rev. Lett. 93, 130403 (2004).
 - [9] M.H. Szymanska, B.D. Simons, and K. Bumet, cond-mat/0412454.

- [10] J. Ranninger and S. Robaszkiewicz, *Physica B* 135, 468 (1985); S. Robaszkiewicz, R. M. icas and J. Ranninger, *Phys. Rev. B* 36, 180 (1987); R. M. icas, J. Ranninger, and S. Robaszkiewicz, *Rev. Mod. Phys.* 62, 113 (1990); J. Ranninger, J.-M. Robin, and M. Eschrig, *Phys. Rev. Lett.* 74, 4027 (1995); T. Domanski and J. Ranninger, *Phys. Rev. B* 63, 134505 (2001); M. Cuoco and J. Ranninger, *Phys. Rev. B* 70, 104509 (2004).
- [11] T. Domanski, *Phys. Rev. A* 68, 013603 (2003).
- [12] P. W. Anderson, *Phys. Rev.* 112, 1900 (1958).