

Magnetism and the Weiss Exchange Field - A Theoretical Analysis Inspired by Recent Experiments

C. Albert*, L. Ferrari†, J. Fröhlich, B. Schlein‡
Theoretical Physics, ETH-Hönggerberg, CH-8093 Zürich

March 23, 2022

Abstract

The huge spin precession frequency observed in recent experiments with spin-polarized beams of hot electrons shot through magnetized films is interpreted as being caused by Zeeman coupling of the electron spins to the so-called *Weiss exchange field* in the film. A “Stern-Gerlach experiment” for electrons moving through an inhomogeneous exchange field is proposed. The microscopic origin of exchange interactions and of large mean exchange fields, leading to different types of magnetic order, is elucidated. A microscopic derivation of the equations of motion of the Weiss exchange field is presented. Novel proofs of the existence of phase transitions in quantum *XY*-models and antiferromagnets, based on an analysis of the statistical distribution of the exchange field, are outlined.

1 Introduction

Effects of ferromagnetism have been known since antiquity. But a mathematical understanding of the microscopic origin of ferromagnetism has remained somewhat elusive, until today! Pauli paramagnetism, ferro-, ferri- and antiferromagnetism are quantum phenomena connected to the spin of electrons and to Pauli’s exclusion principle. The theory of *paramagnetism* in (free) electron gases is quite straightforward, [1]. *Antiferromagnetism* is relatively well understood: A mechanism for the generation of antiferromagnetic exchange interactions has been proposed by Anderson [2], who discovered a close relationship between the half-filled Hubbard model and the Heisenberg antiferromagnet using perturbative methods; (see also [3] for mathematically more compelling and more general variants of Anderson’s key observation). It has been proven rigorously by Dyson, Lieb and Simon [4], using the method of infrared bounds previously discovered in [5], that the *quantum Heisenberg antiferromagnet* with nearest-neighbour exchange couplings exhibits a phase transition accompanied by spontaneous symmetry breaking and the emergence of gapless spin waves, as the temperature is lowered, in *three or more* dimensions. (The Mermin-Wagner theorem says that, in (one and) two dimensions, continuous symmetries cannot be broken spontaneously in models with short-range interactions, [6].)

Our mathematical understanding of *ferromagnetism* is far less advanced. Some kind of heuristic theory of ferromagnetism emerged, long ago, in the classic works of Heisenberg, Bloch, Stoner, Dyson, Landau and Lifshitz, and others [7]. Various insights have been gained on the basis of some form

*supported in part by the Swiss National Foundation

†present address: Collegio Papio, Ascona, Switzerland

‡present address: Department of Mathematics, Stanford University, Stanford CA 94305, USA

of mean-field theory, with small fluctuations around mean-field theory taken into account within a linear approximation. This approximation, however, is known to break down in the vicinity of the critical point of a ferromagnetic material, where nonlinear fluctuations play a crucial role, [8].

In a variety of tight-binding models of itinerant electrons, ferromagnetic order has been exhibited in the *ground state* (i.e., at zero temperature); see [9],[10],[11]. One of these models is a fairly natural two-band model in which ferromagnetism arises from a competition between electron hopping, Coulomb repulsion and an on-site Hund's rule, [11]. (Hund's rule says that the spin-tripled state of two electrons occupying the same site is energetically favoured over the spin-singlet state. It should be emphasized, however, that a mathematically rigorous derivation of Hund's rule in atomic physics from first principles has *not* been accomplished, so far.) None of the results in [9],[10],[11] comes close to providing some understanding of ferromagnetic order and of an order-disorder phase transition at *positive* temperature. It is not known how to derive, with mathematical precision, an effective Hamiltonian with explicit ferromagnetic exchange couplings from the microscopic Schrödinger equation, or a tight-binding approximation thereof, of ferromagnetic materials. But even if we resort to a phenomenological description of such materials in terms of models where ferromagnetic exchange couplings have been put in *by hand* we face the problem that we are unable to exhibit ferromagnetic order at low enough temperature and to establish an order-disorder phase transition in three or more dimensions. *No mathematically rigorous proof* of the phase transition in the *quantum Heisenberg ferromagnet* is known, to date! (Such a result has, however, been established for *classical* Heisenberg models in [5].)

Ab-initio quantum Monte Carlo simulations of models of quantum ferromagnets are plagued by the well known “sign- (or complex-phase) problem”.

Thus, until now, there are neither substantial mathematically rigorous results on, nor are there reliable ab-initio numerical simulations of, realistic models of ferromagnetic metals, such as Ni, Co or Fe! Given that ferromagnetism is among the most striking *macroscopic* manifestations - apparent, e.g., in the needle of a compass - of the *quantum-mechanical* nature of matter, this is clearly a desolate state of affairs.

In the present paper we shall not remedy this unsatisfactory situation. However, first, we attempt to draw renewed attention to it, and, second, we outline a formalism and some fairly elementary analytical observations of which we hope that they will ultimately lead to a better, mathematically rather precise understanding of ferromagnetism. Were it not known already, our analysis and the one in [11] would make clear that ferromagnetism is a non-perturbative phenomenon involving strong correlations and gapless modes. To understand it mathematically will most probably necessitate a full-fledged *multi-scale* (renormalization group) *analysis*. The formalism presented in this paper and our calculations are intended to provide a convenient starting point for such an analysis.

Analytical work on ferromagnetism may seem to be rather unfashionable. However, there are recent developments, such as spintronics, fast magnetic devices, etc. that may make work like ours appear worthwhile. Our own motivation for the work that led to this paper actually originated in studying recent experiments with beams of spin-polarized, hot electrons shot through ferromagnetically ordered films consisting of Ni, Co or Fe that were carried out in the group of H.C. Siegmann at ETH; see [12],[13]. Back in 1998, it became clear to one of us that the concept of the “*Weiss exchange field*” (see [14],[15]) would play a useful role in a theoretical interpretation of the experimental results reported in [12],[13]. More generally, the Weiss exchange field appears to offer a key to a systematic study of phase transitions in magnetic materials, magnetic order, spin precession and magnon dynamics. In this paper, we focus on elucidating the microscopic origin of the Weiss exchange field, the role it plays in the theory of magnetism, and its dynamics.

Our paper is organized as follows.

In Section 2, we describe the experiments reported in [12],[13] and sketch a phenomenological

interpretation, based on scattering theory, of the results found in these experiments, merely adding some conceptual remarks to the discussion of our experimental colleagues and describing the role played by the Weiss exchange field. We also propose some further experiments, in particular a Stern-Gerlach experiment for electrons traversing an inhomogeneous exchange field.

A mathematically precise analysis of the scattering of electrons (or neutrons, photons, ...) at dynamical targets, such as magnetic films, metallic solids, liquid droplets, ..., will appear elsewhere; (some first results appear in [16]).

In Section 3, we reformulate one-band $t - J$ and Hubbard models in terms of a dynamical Weiss exchange field. For this purpose, the standard imaginary-time functional integral formalism for the analysis of thermal equilibrium states of quantum many-body systems is recalled. The *Weiss exchange field* is seen to be a Lagrange-multiplier field in a Hubbard-Stratonovich transformation of the original functional integral that renders the action functional quadratic in the Grassmann variables describing the electronic degrees of freedom; see e.g. [15],[17]. The effective field theory of the Weiss exchange field is obtained after integrating over those Grassmann variables. The effective (imaginary-time) action functional of the exchange field and identities for Green functions of spin operators are derived.

In Section 4, we determine the leading terms of the effective action of the Weiss exchange field, \mathbf{W} , in the approximation where fluctuations of the *length* of the exchange field are neglected. For this purpose, we derive the ferro- and antiferromagnetic mean-field equations from the exact effective action of the exchange field. By solving these equations we determine the most likely length, W_0 , of the exchange field. From that point on, the length of the exchange field is frozen to be $|\mathbf{W}| \equiv W_0$.

We then consider a one-band Hubbard model with a half-filled band and find that, in this situation, the effective action of \mathbf{W} is the one of a nonlinear σ -model with a minimum that favours Néel order. This result is found on the basis of controlled perturbative calculations and goes beyond linear stability analysis of the antiferromagnetic mean-field solution, (which has been presented, e.g., in [15]). It represents a functional-integral version of Anderson's basic observations [2].

We then turn to ferromagnetically ordered mean field solutions and show that x -independent fluctuations are not a source of instability of such a solution. Then we consider a one-band Hubbard model with a weakly filled, fairly flat band. In this situation one expects that ferromagnetism prevails. Indeed, we find that, at low temperatures, the ferromagnetic mean-field equation has a non-trivial solution, and that this solution belongs to a *quadratically stable* critical point of the effective action of \mathbf{W} . This conclusion is the result of somewhat subtle calculations involving processes close to the Fermi surface, which make the dominant contribution (but would lead to small-energy denominators in a purely perturbative analysis). Details will appear in [18]. Our calculations support the idea that the one-band Hubbard model with a weakly filled, fairly flat band describes coexistence of metallic behaviour with ferromagnetic order, at sufficiently low temperatures. A similar conclusion was reached, tentatively, in [11] for some two-band Hund-Hubbard models. The methods of the present paper also apply to the model discussed in [11]; see [18].

In the next to last subsection of Section 4, we exhibit a universal Wess-Zumino term in the effective action of the Weiss exchange field \mathbf{W} and calculate its coefficient, which is purely imaginary. The Wess-Zumino term is "irrelevant" for antiferromagnets, but plays a *crucial* role in the dynamics of magnons in *ferromagnetically* ordered systems. Repeating arguments in [19], we derive the Landau-Lifshitz equations for magnons in a ferromagnet.

Finally, we draw attention to two well known arguments explaining why there is no magnetic ordering at positive temperature, in one and two dimensions; (but see [6]).

In Section 5, we sketch novel rigorous proofs, based on analyzing the effective field theory of the exchange field, \mathbf{W} , of the existence of phase transitions and magnetic order at low temperatures in a class of XY -models, Heisenberg antiferromagnets and ferromagnets of localized $SO(2n)$ - spins, for

$n = 1, 2, \dots$. See [4] for the original results. Our proof is based on establishing reflection positivity of the effective field theory of the exchange field \mathbf{W} and then using the original techniques developed in [5]; (see also [20], [21]).

It should be emphasized that the concept of the Weiss exchange field has a number of further, quite exciting applications. We hope to return to these matters in future papers.

Some of the material in this paper has a review character; but some of it is new. We hope it is fairly easy to read. If it draws renewed attention to some of the deep technical problems in the quantum theory of magnetism it has fulfilled its purpose. We gratefully dedicate this paper, belatedly, to two great colleagues and friends of the senior author (J.F.): G. Jona-Lasinio, on the occasion of his seventieth birthday, and H.-C. Siegmann, on the occasion of his retirement from ETH.

Acknowledgements. J. Fröhlich thanks S. Riesen and H.-C. Siegmann for very stimulating discussions of the experiments in [12], P. Wiegmann for some crucial advice with the calculations in Section 4, and the IHÉS for hospitality during much of the work on this paper. We all thank M. Azam for very useful and pleasant discussions on the uses of the exchange field.

2 Real and gedanken experiments involving the Weiss exchange field

We start this section with a brief description of recent experiments carried out by Oberli, Burgermeister, Riesen, Weber and Siegmann at ETH-Zürich [12], [13]. In these experiments, a beam of hot, spin-polarized electrons is shot through a thin ferromagnetic film (Ni, Co, or Fe) and the polarization of the outgoing beam is observed. Their experimental setup is as described in Fig. 1.




Figure 1: Experimental setup

The following quantities are measurable:

- i) The thickness, d , of the film; d is a few nanometers.
- ii) The average energy, E , of an incident electron; if E_F denotes the Fermi energy of the magnetic film then $E - E_F$ varies between $4eV$ and $16eV$. The group velocity of the electrons inside the film is denoted by \mathbf{v} ; it is not directly measurable, but it is comparable to $(2(E + eV)/m_{el}^*)^{1/2}$, where eV is the average potential energy of an electron and m_{el}^* its effective mass inside the film.
- iii) The degree, P_0 , and the direction, \mathbf{n}_0 , of the spin polarization of the incident electron beam; (in Fig. 1, \mathbf{n}_0 is parallel to the x -direction, \mathbf{v} to the y -direction); the same quantities, P and \mathbf{n} , for the outgoing beam.

- iv) The direction of the magnetization, \mathbf{M} , of the film (in Fig. 1 chosen to be parallel to the z -axis); the angles, θ_0 and θ , between \mathbf{n}_0 and \mathbf{M} and between \mathbf{n} and \mathbf{M} , respectively ($\theta_0 = \pi/2$, in Fig. 1). Experimentally, the angle θ is found to be considerably smaller than θ_0 , i.e., the spins of the transmitted electrons rotate into the direction of the spontaneous magnetization \mathbf{M} of the film. This is interpreted as being mainly due to an enhanced absorption of minority-spin electrons, as compared to majority-spin electrons; see (vi). (It appears that the contribution of spin flip processes accompanied by magnon emission into the film – “Stoner excitations” – to the total spin rotation is only around 5%, [12]).
- v) The spin precession angle, ϵ , between the projections of \mathbf{n}_0 and of \mathbf{n} onto the plane perpendicular to \mathbf{M} (the xy -plane of Fig. 1); ϵ is found to be “large” (tens of degrees).
- vi) Let I be the intensity of the incident beam, and let $I^+ = I^+(E)$ and $I^- = I^-(E)$ be the intensities of the outgoing beam of electrons with spin parallel or antiparallel to \mathbf{M} , respectively, assuming the incident beam has intensity I and the spins of its electrons are parallel to \mathbf{M} (\mathbf{n}_0 parallel to \mathbf{M} , $P_0 \cong 1$), or antiparallel to \mathbf{M} (\mathbf{n}_0 anti-parallel to \mathbf{M} , $P_0 \cong 1$), respectively. Then $\theta_0 = \theta = 0$, or $\theta_0 = \theta = \pi$, respectively, and $\epsilon = 0$. I^+ and I^- can be measured and yield the spin-transmission asymmetry

$$A = \frac{I^+ - I^-}{I^+ + I^-};$$

A is found to be positive and large. This is interpreted in terms of rates of transitions of electrons into unoccupied 3d states (holes): There are more unoccupied 3d states in the film with spin antiparallel to \mathbf{M} (minority spin) than with spin parallel to \mathbf{M} (majority spin). This explains qualitatively the experimental results found for A and for $\theta - \theta_0$ (see (iv)); [12].

- vii) The orbital deflection angle, α , between the directions of the incident and the transmitted beam (not indicated in Fig. 1). Experimentally α is found to be negligibly small. This tells us that the integrated Lorentz force on the electrons transmitted through the film is tiny. The precession of the spins of the electrons when they traverse the film can therefore *not* be explained by Zeeman coupling of the spins to the magnetic field inside the layer. It is mainly due to Zeeman coupling of the spins to what will be called the *Weiss exchange field*. In iron, the Weiss exchange field causing the observed spin precession would correspond to a magnetic field of roughly 8000 Tesla (which is gigantic).

A theoretical interpretation of the experimental results reported in [12] can be attempted within the formalism of *scattering theory*. If the luminosity of the incident beam is low we can consider a single incoming electron. The incoming state is described as a tensor product of a Pauli spinor, ψ_{in} , describing the incident electron and a state, ξ , of the film. Typically, ξ is the ground state (temperature $T = 0$) or a thermal equilibrium state ($T > 0$) of the film. The outgoing state, long after the interactions between an outgoing electron and the film have taken place, is more complicated and will, in general, exhibit *entanglement* between the electron and the degrees of freedom of the film. If only measurements far away from the film are performed, as in [12], the outgoing state can be described as a density matrix

$$P_{\text{out}} = (\rho_N)_{N=0}^{\infty}, \quad (2.1)$$

where ρ_N is a non-negative, trace-class operator on the Hilbert space of N *outgoing electrons* (the incident electron has knocked $N - 1$ electrons out of the film), $N = 2, 3, \dots$; ρ_0 is a non-negative number, the *absorption probability*, ρ_1 is a non-negative trace-class operator on the Hilbert space

$$\mathcal{H} = L^2(\mathbb{R}^3) \otimes \mathbb{C}^2 \quad (2.2)$$

of square-integrable Pauli spinors and describes the (generally *mixed*) state of *one* outgoing electron. “Conservation of probability” implies that

$$\rho_0 + \sum_{N=1}^{\infty} \text{Tr} \rho_N = 1. \quad (2.3)$$

The state P_{out} is obtained by taking a partial trace of the outgoing state of the *total system*, *including* the film, over the degrees of freedom of the film. This is justified, because the degrees of freedom of the film are *not* observed in the experiment. If the energy, E , of the incident electron is below (or comparable to) the threshold, $\Sigma^{(2)}$, for emission of two or more electrons from the film then

$$\rho_N = 0 \quad \text{for } N \geq 2. \quad (2.4)$$

In the interpretation of the experimental data provided in [12], this is tacitly assumed.

Experimentally, the absorption probability ρ_0 and the spin polarizations \mathbf{P}_0 and \mathbf{P} of the incoming and the outgoing electron, respectively, are measured. The vectors \mathbf{P}_0 and \mathbf{P} are given by

$$\mathbf{P}_0 \equiv P_0 \mathbf{n}_0 = \langle \psi_{\text{in}}, \boldsymbol{\sigma} \psi_{\text{in}} \rangle, \quad (2.5)$$

and

$$\mathbf{P} \equiv P \mathbf{n} = \frac{1}{\text{Tr}_{\mathcal{H}}(\rho_1)} \text{Tr}_{\mathcal{H}}(\rho_1 \boldsymbol{\sigma}), \quad (2.6)$$

where ψ_{in} is the wave function of the incident electron, $\boldsymbol{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ is the vector of Pauli matrices, and $\tilde{\rho}_1 := [\text{Tr}_{\mathcal{H}}(\rho_1)]^{-1} \rho_1$ is the *conditional* state of the outgoing electron, given that it has *not* been absorbed in the film. If (2.4) is assumed to hold then

$$\text{Tr}_{\mathcal{H}}(\rho_1) = 1 - \rho_0 \quad (2.7)$$

Since we have assumed that the incoming electron has been prepared in a pure state,

$$P_0 \equiv |\mathbf{P}_0| = 1. \quad (2.8)$$

However, there is, a priori, no reason why $\tilde{\rho}_1$ should be a pure state. If it were pure then

$$P = |\mathbf{P}| = 1. \quad (2.9)$$

It would be highly interesting to estimate, experimentally, the amount of entanglement with the film (or *decoherence*) in the state of the outgoing electron by measuring the quotient P/P_0 . If $P/P_0 < 1$ then $\tilde{\rho}_1 = [\text{Tr}_{\mathcal{H}}(\rho_1)]^{-1} \rho_1$ is *not* a pure state, anymore, meaning there is entanglement with the film. Apparently, P/P_0 has not been measured accurately, yet.

A moment’s reflection shows that spin flip processes accompanied by magnon emission in the film (“Stoner excitations”) lead to entanglement; while absorption of electrons into unoccupied 3d states need not be correlated with entanglement of the states of *those* electrons that *do* traverse the film. In fact, it is implicitly assumed in [12] that if Stoner excitations are neglected then $\tilde{\rho}_1$ is close to a pure state (at least in spin-space). The experimental techniques of [12] could be used to test this hypothesis.

Next, we express the spin transmission asymmetry A (see (vi)) in terms of outgoing states. Let ψ_{in}^+ and ψ_{in}^- be incoming states with spin polarization $\mathbf{P}_0 \equiv \mathbf{P}_0^+$ parallel to \mathbf{M} (majority spin) and $\mathbf{P}_0 \equiv \mathbf{P}_0^-$ antiparallel to \mathbf{M} (minority spin). Let $P_{\text{out}}^+ = (\rho_N^+)_{N=0}^{\infty}$ and $P_{\text{out}}^- = (\rho_N^-)_{N=0}^{\infty}$ be outgoing states corresponding to ψ_{in}^+ , ψ_{in}^- , respectively. Then

$$A = \frac{\rho_0^- - \rho_0^+}{2 - \rho_0^+ - \rho_0^-}, \quad (2.10)$$

as follows from (2.3). More interesting would be measurements of

$$\mathbf{P}^\pm = \text{Tr}(\tilde{\rho}_1^\pm \boldsymbol{\sigma}). \quad (2.11)$$

Clearly \mathbf{P}^\pm are parallel or anti-parallel to \mathbf{M} ; but their lengths $P^\pm = |\mathbf{P}^\pm|$ ought to be measured. In [12], it is tacitly assumed that $P^\pm \cong 1$, and that the states $\tilde{\rho}_1^\pm$ are close to *pure* states; but serious experimental data backing up this hypothesis appear to be lacking. It is clear that it would be invalidated if “Stoner excitations” played an important role.

In the following, we outline a phenomenological description of the experiments in [12], assuming that (2.4) and the hypothesis just discussed (purity of $\tilde{\rho}_1^\pm$) are valid. (A more detailed discussion of the scattering approach to electron transmission- and reflection experiments will be presented elsewhere.)

When an incoming electron enters the film it occupies an empty state of the film. If the film is crystalline this state belongs to a band of states; let α be the corresponding band index. The state of an electron in band α is described by a Pauli spinor

$$\phi = (\phi^+, \phi^-), \quad (2.12)$$

where ϕ^+ and ϕ^- are the components of ϕ with spin parallel or anti-parallel to the magnetization \mathbf{M} , respectively. Adopting the approximation of the Peierls substitution, the Pauli equation for ϕ in configuration space has the form

$$i\hbar \mathcal{D}_0 \phi = \mathcal{E}_\alpha(-i\hbar \mathcal{D}) \phi, \quad (2.13)$$

where $\mathcal{E}_\alpha(\mathbf{p})$ is the band function of band α ,

$$\mathcal{D}_0 = \frac{\partial}{\partial t} + i \frac{e}{\hbar} \phi^c + i \left(\mathbf{W}_0^c \cdot \frac{\boldsymbol{\sigma}}{2} \right), \quad (2.14)$$

$$\mathcal{D}_j = \frac{\partial}{\partial x^j} + i \frac{e}{\hbar} A_j + i \left(\mathbf{W}_j \cdot \frac{\boldsymbol{\sigma}}{2} \right). \quad (2.15)$$

Here ϕ^c denotes a (complex) electrostatic potential, \mathbf{W}_0^c is a (complex) *Weiss exchange field*, $(A_j) = (A_1, A_2, A_3)$ is the electromagnetic vector potential, and $(\mathbf{W}_j) = (\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_3)$ is an $SU(2)$ -vector potential responsible for *spin-orbit interactions*. As discussed in [14], (2.13) displays electromagnetic $U(1)$ -gauge invariance and $SU(2)$ -gauge invariance, i.e., covariance with respect to local $SU(2)$ -rotations in spin space. A number of important consequences of these gauge symmetries have been pointed out in [14].

According to (vii) above, effects of the electromagnetic vector potential \mathbf{A} are apparently negligible; so \mathbf{A} is set to 0. The electrostatic vector potential ϕ^c is given, approximately, by

$$\phi^c = V + i\hat{v} \quad (2.16)$$

where eV is the surface exit work, and the imaginary part, \hat{v} , of ϕ^c provides a phenomenological description of spin-independent inelastic absorption processes inside the film. Velocity-dependent spin-flip processes due to *spin-orbit interactions* appear to play a very minor role in the experiments reported in [12]; so we may set \mathbf{W}_j to 0, $j = 1, 2, 3$.

The Weiss exchange field \mathbf{W}_0^c is given by

$$\mathbf{W}_0^c = \mathbf{W} - i\mathbf{w}, \quad (2.17)$$

where the real part \mathbf{W} describes *exchange interactions* between the incoming electron and the electron density of the film, and the imaginary part, \mathbf{w} , yields a phenomenological description of spin-dependent absorption processes.

Let ϕ_{in} denote the Pauli spinor describing the state of an electron when it enters the film at some time t_0 . Eq. (2.13) can be solved for $\phi = \phi_t$, $t \geq t_0$, with $\phi_{t_0} = \phi_{\text{in}}$. The solution is explicit if $\mathbf{A} = 0$, $\mathbf{W}_j = 0$, $j = 1, 2, 3$. Let us suppose that the real part \mathbf{W} and the imaginary part \mathbf{w} of the exchange field \mathbf{W}_0^c are both anti-parallel to the magnetization \mathbf{M} of the film. Then (2.13) leads us to consider two simple, quasi-one-dimensional scattering problems in a complex potential well of depth

$$eV + ie\hat{v} \pm \frac{1}{2}(\Omega - i\omega), \quad (2.18)$$

with $\Omega = |\mathbf{W}|$ and $\omega = |\mathbf{w}|$, for electrons with spin parallel to $\mathbf{M}(+)$, or anti-parallel to $\mathbf{M}(-)$, respectively. The solution to these scattering problems can be found in every book on elementary quantum mechanics. For the purposes of interpreting the results in [12], a semi-classical treatment appears to be adequate. The group velocity, \mathbf{v}_{\pm} , of an incoming electron wave with energy peaked at E and spin up (+), or down (-), inside the film can be found by solving the equation

$$\mathcal{E}_{\alpha}(\mathbf{p}) = E - eV \mp \frac{1}{2}\Omega \quad (2.19)$$

for \mathbf{p} and then setting

$$\mathbf{v}_{\pm} = \frac{\partial \mathcal{E}_{\alpha}}{\partial \mathbf{p}}(\mathbf{p}^{\pm}), \quad (2.20)$$

where \mathbf{p}^{\pm} is a solution of (2.19) chosen such that \mathbf{v}_{\pm} points in the positive y -direction. The sojourn time τ_{\pm} of the wave inside the film is then given by

$$\tau_{\pm} = \frac{d}{v_{\pm}}, \quad \text{with } v_{\pm} = |\mathbf{v}_{\pm}|. \quad (2.21)$$

If ϕ_{in}^{\pm} is the state of the electron with spin up (+), or spin down (-), respectively, when it enters the film its state ϕ_{out}^{\pm} upon leaving the film is then given, approximately, by ¹

$$\phi_{\text{out}}^{\pm} \cong \exp\left(-i\tau_{\pm}[(E - eV) - ie\hat{v} \mp \frac{1}{2}(\Omega - i\omega)]\right) \phi_{\text{in}}^{\pm}. \quad (2.22)$$

The presence of inelastic absorption processes implies that $e\hat{v} \mp \omega/2 > 0$. Setting

$$\exp(-\tau_{\pm}[\hat{v} \mp \omega/2]) = c\sqrt{1 \pm A}, \quad (2.23)$$

we find that

$$\phi_{\text{out}}^{\pm} = c\sqrt{1 \pm A}e^{i(\theta \pm \epsilon/2)} \phi_{\text{in}}^{\pm}, \quad (2.24)$$

where A is the spin transmission asymmetry (see point (iv) and (2.10)), ϵ is the spin precession angle, and θ is an (unimportant) spin-independent phase. If $v_+ \cong v_- = v$ then $\tau_+ \cong \tau_- \cong d/v$, and (2.22) yields

$$\epsilon \cong \tau_{\pm}\Omega \cong (d/v)|\mathbf{W}|. \quad (2.25)$$

Thus, measuring ϵ and d and estimating v yields an approximate value for the size of the spin precession angular velocity Ω and hence of the size of the Weiss exchange field.

Eqs. (2.22) and (2.24) could be mistaken for equations describing *kaon-* or *neutrino oscillations* and are analogous to the equations describing the *Faraday rotation* of light traversing a magnetized medium.

¹Here and henceforth, we use units such that $\hbar = 1$.

The origin of the Weiss exchange field \mathbf{W} is hardly a mystery: The spin of an electron traversing the film in a band α apparently experiences (exchange) interactions with the spins of the occupied states of the film. Since states with spin up and with spin down are occupied asymmetrically (corresponding to the fact that $\mathbf{M} \neq 0$), the net spin density, $\mathbf{S}(x)$, at a point x in the film is different from zero. The Weiss exchange field $\mathbf{W} = \mathbf{W}(x)$ is given by

$$\mathbf{W}(x) = -J_\alpha \mathbf{S}(x), \quad (2.26)$$

where J_α is the strength of the *exchange coupling* between spins in the α^{th} band and those in the occupied band. The theoretical discussion above is based on a *mean field ansatz*: The exchange field \mathbf{W} in (2.14), (2.17) is chosen to be

$$\mathbf{W} = \langle \mathbf{W}(x) \rangle = -J_\alpha \langle \mathbf{S}(x) \rangle. \quad (2.27)$$

(Our conventions are such that $\langle \mathbf{S}(x) \rangle$ is parallel to the magnetization \mathbf{M} .) One of the surprising implications of the experimental results of [12] is that, apparently, J_α is quite large, even for rather high-lying bands (α), implying that the orbitals of states in such bands must have substantial overlap with those of states in the partially occupied, spin-polarized band.

Eq. (2.26) makes it clear that $\mathbf{W}(x)$ is a *dynamical* field. One of the main purposes of this paper is to derive the effective quantum dynamics of $\mathbf{W}(x)$ within a Lagrangian functional integral formalism and to sketch what can be accomplished with this formalism.

Before proceeding with this program, we outline some gedanken experiments in the spirit of the experiments in [12].

- i) By applying an external magnetic field to the film rotating in the xy -plane with angular velocity ω_0 , the exchange field \mathbf{W} could be made to rotate around the z -axis:

$$\mathbf{W}(t) = W \left(\epsilon \cos(\omega_0 t + \delta), \epsilon \sin(\omega_0 t + \delta), \sqrt{1 - \epsilon^2} \right). \quad (2.28)$$

A polarized beam of electrons shot through this film must exhibit *Bloch spin resonance*; but, in this experiment, it would be due to the rotation of the exchange field.

- ii) One may envisage a *Stern Gerlach experiment* for electrons. One would start by constructing a sandwich of two ferromagnetic metals, I (e.g. Fe) and II (e.g. Ni), with exchange fields \mathbf{W}_I and \mathbf{W}_{II} of different strength, joined by a transition region, a mixture of I and II, of width d_0 . The transition region would be parallel to the xy -plane (see Fig. 2). One shoots an unpolarized beam of (not very hot) electrons through the sandwich along the transition region between I and II, as shown in Fig. 2. One would expect to detect two beams emerging on the other side of the film in slightly different directions that are spin-polarized in opposite directions.

The force, f , in the z -direction on an electron with spin up/down inside the film is given, approximately, by

$$f \cong \pm \frac{|\mathbf{W}_I - \mathbf{W}_{II}|}{2d_0}. \quad (2.29)$$

It yields a change in the z -component of the momentum of the electron during its passage through the film given by

$$\Delta p_z \cong \pm \frac{|\mathbf{W}_I - \mathbf{W}_{II}|}{2d_0} \frac{d}{v}, \quad (2.30)$$

where v is the average group velocity. The deflection angle α is found from

$$\tan \alpha = \frac{\Delta p_z}{\Delta p_y}. \quad (2.31)$$




Figure 2: Stern-Gerlach Experiment

Of course, as discussed above, the intensity, I_+ of the upper beam can be expected to be much larger than the intensity, I_- , of the lower beam, due to spin-asymmetric absorption inside the film. In order to achieve adequate focussing, the incident beam could come from the tip of a scanning tunnelling microscope.

- iii) In an experimental set-up similar to the one above, one would force an electron current, spin-polarized in the z -direction, through the film. Then a Hall tension in the z -direction should be observed (Hall effect for spin currents; see [14]).

3 The Weiss exchange field in general one-band (t - J and Hubbard) tight-binding models

In this section, we study the origin and dynamics of the Weiss exchange field in simple tight-binding models. We start by considering one-band models; but it is straightforward to include higher (partially occupied or empty) bands relevant for the study of electron transmission through magnetic films.

Every site, x , of a lattice $\Gamma \stackrel{e.g.}{=} \mathbb{Z}^d$, $d = 1, 2, 3, \dots$ carries a four-dimensional state space, \mathbb{C}^4 , corresponding to an empty state, a one-electron state with spin up (+) or down (-), or a state of two electrons in a spin-singlet state. The corresponding basis vectors of \mathbb{C}^4 are denoted by $\{|0\rangle_x, |+\rangle_x, |-\rangle_x, |+-\rangle_x\}$, $x \in \Gamma$. We introduce electron creation- and annihilation operators, $c_s^\dagger(x)$ and $c_s(x)$, respectively, where $s = \pm$, $x \in \Gamma$, satisfying canonical anti-commutation relations

$$\{c_s^\#(x), c_{s'}^\#(x')\} = 0, \quad \{c_s^\dagger(x), c_{s'}(x')\} = \delta_{ss'} \delta_{xx'}, \quad (3.1)$$

with $c^\# = c$ or c^\dagger . Then

$$c_s(x)|0\rangle_x = 0, \quad \text{for } s = \pm, x \in \Gamma \quad (3.2)$$

and

$$|s\rangle_x = c_s^\dagger(x)|0\rangle_x, \quad \text{etc.} \quad (3.3)$$

Number operators are given by

$$n_s(x) = c_s^\dagger(x)c_s(x), \quad n(x) = n_+(x) + n_-(x), \quad (3.4)$$

where $n_s(x)$ measures the number of electrons at site x with spin s . By (3.1), $n_s(x)$ has eigenvalues 0 and 1, while $n(x)$ has eigenvalues 0, 1 and 2. The spin operator, $\mathbf{S}(x)$, at site x is given by

$$\mathbf{S}(x) = \frac{1}{2} \sum_{s,s'} c_s^\dagger(x) \boldsymbol{\sigma}_{ss'} c_{s'}(x) \quad (3.5)$$

where $\sigma_{ss'}$ is the (ss') -matrix element of the vector of Pauli matrices $\sigma = (\sigma_x, \sigma_y, \sigma_z)$. The operators $\mathbf{S}(x)$ vanish on $|0\rangle_x$ and $|+-\rangle_x$, while, for any linear combination $\phi(x)$ of $|+\rangle_x$ and $|-\rangle_x$,

$$\mathbf{S}(x) \cdot \mathbf{S}(x) \phi(x) = \frac{3}{4} \phi(x). \quad (3.6)$$

The dynamics of a gas of electrons moving on the lattice Γ is assumed to be generated by a $t - J$ model Hamiltonian of the form

$$H = T + U + E_{\neq}, \quad (3.7)$$

where the kinetic energy operator, T (the hopping term), is given by

$$T = \sum_{x \neq y} \hat{t}(x - y) \sum_s c_s^\dagger(x) c_s(y), \quad (3.8)$$

$$U = U_0 \sum_x n_+(x) n_-(x) \quad (3.9)$$

describes an on-site (Hubbard) repulsion, and

$$E_{\neq} = -\frac{1}{2} \sum_{x,y} J_{\neq}(x - y) \mathbf{S}(x) \cdot \mathbf{S}(y) \quad (3.10)$$

describes (effective) exchange interactions between the spins of electrons. In (3.8), $\hat{t}(x - y)$ is the amplitude for hopping of an electron from site y to site x . Selfadjointness of the Hamiltonian H implies that $\hat{t}(x - y) = \hat{t}(y - x)$. The constant U_0 is a measure of the strength of the on-site repulsion, and $J_{\neq}(x - y)$ is the exchange coupling between spins at x and y , with $J_{\neq}(0) = 0$.

The term U can be rewritten as follows. When x is empty or singly occupied $U_0 n_+(x) n_-(x)$ vanishes; when x is doubly occupied it takes the value U_0 . Thus, by (3.6),

$$U_0 n_+(x) n_-(x) = \frac{1}{2} U_0 n(x) - \frac{2}{3} U_0 \mathbf{S}(x) \cdot \mathbf{S}(x) = \frac{2}{3} U_0 : \mathbf{S}(x) \cdot \mathbf{S}(x) :, \quad (3.11)$$

where $: \cdot :$ denotes normal ordering. We now set $J(0) = \frac{4}{3} U_0$, $J(x) = J_{\neq}(x)$, for $x \neq 0$, and find

$$H = T + E, \quad (3.12)$$

with

$$E = \frac{1}{2} \sum_{x,y} J(x - y) : \mathbf{S}(x) \cdot \mathbf{S}(y) :. \quad (3.13)$$

The *Heisenberg model* corresponds to a completely flat band, i.e.,

$$T = 0, \quad \text{or} \quad \hat{t}(x) \equiv 0. \quad (3.14)$$

Our goal is to study the grand-canonical partition function

$$\Xi(\beta, \mu) := \text{Tr} \left(e^{-\beta[H - \mu \sum_x n(x)]} \right) \quad (3.15)$$

and the thermal equilibrium state given by the density matrix

$$P_{\beta, \mu} = \Xi(\beta, \mu)^{-1} e^{-\beta[H - \mu \sum_x n(x)]} \quad (3.16)$$

for the family of Hamiltonians introduced above.

Remark. Of course, these objects must first be calculated for arbitrary bounded regions, Λ , in the lattice Γ (indicated by a superscript “ Λ ”). Afterwards, one may attempt to pass to the thermodynamic limit, $\Lambda \nearrow \Gamma$, setting

$$\beta g(\beta, \mu) = - \lim_{\Lambda \nearrow \Gamma} \frac{1}{|\Lambda|} \ln \Xi^\Lambda(\beta, \mu) \quad (3.17)$$

and

$$\omega_{\beta, \mu}(\cdot) = \lim_{\Lambda \nearrow \Gamma} \text{Tr} \left(P_{\beta, \mu}^\Lambda(\cdot) \right), \quad (3.18)$$

where $|\Lambda|$ is the number of sites in Λ . These are standard matters (see e.g. [22]) and will not be discussed any further. The superscript “ Λ ” will often be suppressed in our notation.

It is convenient to study $\Xi(\beta, \mu)$ and the state $\omega_{\beta, \mu}$ by making use of *functional integration*. With every $x \in \Gamma$ and every imaginary time $t \in [0, \beta)$, we associate anticommuting (Grassmann) variables, $\bar{C}_s(x, t)$ and $C_s(x, t)$ with

$$\{C_s^{(-)}(x, t), \bar{C}_{s'}^{(-)}(x', t')\} = 0,$$

and

$$C_s^\#(x, t + \beta) = -C_s^\#(x, t). \quad (3.19)$$

The anti-periodic boundary conditions in (3.19) are a consequence of the KMS condition; see e.g. [17]. We introduce an (imaginary-time) action functional

$$S_J(\bar{C}, C) = \int_0^\beta dt \left[\left(\sum_{x,s} \bar{C}_s(x, t) \frac{\partial}{\partial t} C_s(x, t) \right) - H(\bar{C}(\cdot, t), C(\cdot, t)) - \mu \sum_{x,s} \bar{C}_s(x, t) C_s(x, t) \right], \quad (3.20)$$

where $H(\bar{C}, C)$ is obtained from the Hamiltonian H by replacing the operators $c_s(x)$ and $c_s^\dagger(x)$ by $C_s(x, t)$ and $\bar{C}_s(x, t)$, respectively. Because of the presence of products of C - and \bar{C} -fields at coinciding sites and imaginary times the quartic term needs to be regularized; (the ambiguity of the quadratic terms is an unimportant constant). We introduce an infinitesimal separation of the imaginary times, replacing $\mathbf{S}(x)$ in (3.13) by

$$\mathbf{S}^\epsilon(x, t) = \frac{1}{2} \sum_{s, s'} \bar{C}_s(x, t + \epsilon) \sigma_{ss'} C_{s'}(x, t - \epsilon) \quad (3.21)$$

in the action. Then the time ordering of the C - and \bar{C} -fields corresponds to the normal ordering of the operators c^\dagger . Because of (3.13) the regularized action is now given by

$$\begin{aligned} S_J^\epsilon(\bar{C}, C) = \int_0^\beta dt \left[\left(\sum_{x,s} \bar{C}_s(x, t) \frac{\partial}{\partial t} C_s(x, t) \right) - \sum_{x \neq y} \hat{t}(x - y) \sum_s \bar{C}_s(x, t) C_s(y, t) \right. \\ \left. + \frac{1}{2} \sum_{x,y} J(x - y) \mathbf{S}^\epsilon(x, t) \cdot \mathbf{S}^\epsilon(y, t) + \mu \sum_{x,s} \bar{C}_s(x, t) C_s(x, t) \right], \quad (3.22) \end{aligned}$$

Berezin’s integration form for anticommuting variables is formally given by

$$\mathcal{D}\bar{C}\mathcal{D}C = \prod_{t \in [0, \beta)} \prod_x \Pi_s d\bar{C}_s(x, t) dC_s(x, t). \quad (3.23)$$

Then

$$\Xi(\beta, \mu) = \lim_{\epsilon \searrow 0} \int \mathcal{D}\bar{C} \mathcal{D}C e^{S_J^\epsilon(\bar{C}, C)} \quad (3.24)$$

and the state $\omega_{\beta, \mu}$ can be reconstructed from the imaginary-time Green functions

$$\langle \Pi_j \bar{C}_{s_j}(x_j, t_j) C_{s'_j}(x'_j, t'_j) \rangle_{\beta, \mu} = \Xi(\beta, \mu)^{-1} \lim_{\epsilon \searrow 0} \int \mathcal{D}\bar{C} \mathcal{D}C e^{S_J^\epsilon(\bar{C}, C)} \Pi_j \bar{C}_{s_j}(x_j, t_j) C_{s'_j}(x'_j, t'_j); \quad (3.25)$$

see, e.g., [17], [23].

If $S_J^\epsilon(\bar{C}, C)$ were quadratic in C and \bar{C} the Berezin integrals in (3.24), (3.25) could be evaluated and expressed in terms of determinants. The only contribution to $S_J^\epsilon(\bar{C}, C)$ *not* quadratic in \bar{C} , C comes from the term

$$\int_0^\beta dt \sum_{x, y} J(x - y) \mathbf{S}^\epsilon(x, t) \cdot \mathbf{S}^\epsilon(y, t). \quad (3.26)$$

By introducing a Lagrange-multiplier field (“Hubbard-Stratonovich transformation”) this term, too, can be rendered quadratic in \bar{C} and C . The Lagrange multiplier field will turn out to be the *Weiss exchange field* \mathbf{W} .

Let K denote the matrix inverse of J , i.e., $\hat{K}(k) = \hat{J}(k)^{-1}$, where k is a point in the first Brillouin zone, B_Γ , of the lattice Γ , and \hat{J} , \hat{K} denote the Fourier transforms of J and K . If U_0 is chosen to be large enough then J and K are positive-definite, i.e., $\hat{J}(k) > 0$, $\hat{K}(k) > 0$, for $k \in B_\Gamma$. We introduce a real, vector-valued random field, $\mathbf{W}(x, t)$, $x \in \Gamma$, $t \in [0, \beta)$, with

$$\mathbf{W}(x, t + \beta) = \mathbf{W}(x, t) \quad (3.27)$$

(KMS condition for bosons; see e.g. [17]), assumed to have a Gaussian distribution

$$d\mu_K(\mathbf{W}) = e^{S_K(\mathbf{W})} \mathcal{D}\mathbf{W}, \quad (3.28)$$

where

$$S_K(\mathbf{W}) = -\frac{1}{2} \int_0^\beta dt \sum_{x, y} \mathbf{W}(x, t) K(x - y) \mathbf{W}(y, t), \quad (3.29)$$

$$\mathcal{D}\mathbf{W} = \text{const}_K \Pi_{t \in [0, \beta)} \Pi_x d^3 W(x, t), \quad (3.30)$$

and const_K is chosen such that

$$\int d\mu_K(\mathbf{W}) = 1. \quad (3.31)$$

We define²

$$S_0(\bar{C}, C) = S_{J=0}(\bar{C}, C), \quad (3.32)$$

$$S_1(\bar{C}, C; \mathbf{W}) = - \int_0^\beta dt \left(\sum_x \mathbf{W}(x, t) \cdot \mathbf{S}(x, t) \right), \quad (3.33)$$

and

$$\bar{S}(\bar{C}, C; \mathbf{W}) = S_0(\bar{C}, C) + S_1(\bar{C}, C; \mathbf{W}) + S_K(\mathbf{W}). \quad (3.34)$$

²Henceforth we suppress the regularization indicated by ϵ .

We note that $\bar{S}(\bar{C}, C; \mathbf{W})$ is quadratic in \bar{C} and C and in \mathbf{W} (separately), and that

$$\int \mathcal{D}\mathbf{W} e^{\bar{S}(\bar{C}, C; \mathbf{W})} = e^{S_J(\bar{C}, C)}, \quad (3.35)$$

as follows by quadratic completion.

It is apparent from (3.33) that $\mathbf{W}(x, t)$ plays the role of the Weiss exchange field. In order to calculate the grand canonical partition function $\Xi(\beta, \mu)$, we have to integrate the R.S. of (3.35) over \bar{C} and C . Since $\bar{S}(\bar{C}, C; \mathbf{W})$ is quadratic in \bar{C} and C , it is tempting to interchange the \mathbf{W} - and the \bar{C} , C - integrations. First doing the \bar{C} , C - integration yields

$$\int \mathcal{D}\bar{C} \mathcal{D}C e^{S_0(\bar{C}, C) + S_1(\bar{C}, C; \mathbf{W})} = \det(D\mathbf{W}), \quad (3.36)$$

where the operator $D\mathbf{W}$ acts on the space

$$\mathfrak{h} = C^\infty([0, \beta))_{ap} \otimes (l^2(\Gamma) \otimes \mathbb{C}^2) \quad (3.37)$$

(“ap” stands for “antiperiodic”) and is determined by requiring that

$$S_0(\bar{C}, C) + S_1(\bar{C}, C; \mathbf{W}) = \int_0^\beta dt \sum_{x,s} \bar{C}(x, t) (D\mathbf{W}C)(x, t).$$

Thus, $D\mathbf{W}$ is given by

$$D\mathbf{W} = \frac{\partial}{\partial t} \otimes \mathbf{1} + \mathbf{1} \otimes (\hat{t} - \mu \mathbf{1}) + \frac{1}{2} \mathbf{W}(x, t) \cdot \boldsymbol{\sigma}, \quad (3.38)$$

where $(\hat{t}h)(x) = \sum_y \hat{t}(x - y)h(y)$, $h \in l^2(\Gamma)$.

After Fourier transformation in t and x , vectors in the space \mathfrak{h} are given by two-component spinors

$$\phi(\mathbf{k}, k_0) = \begin{pmatrix} f_+(\mathbf{k}, k_0) \\ f_-(\mathbf{k}, k_0) \end{pmatrix},$$

with $\mathbf{k} \in B_\Gamma$ and $k_0 = (\pi/\beta)(2n + 1)$, $n \in \mathbb{Z}$, and

$$(D\mathbf{W}\phi)(\mathbf{k}, k_0) = (-ik_0 + t(\mathbf{k}) - \mu) \phi(\mathbf{k}, k_0) + \frac{1}{2} \int_{B_\Gamma} \frac{d\mathbf{k}'}{(2\pi)^3} \sum_{k'_0} \hat{\mathbf{W}}(\mathbf{k} - \mathbf{k}', k_0 - k'_0) \cdot (\boldsymbol{\sigma}\phi)(\mathbf{k}', k'_0), \quad (3.39)$$

where $t(\mathbf{k})$ denotes the Fourier transform of $\hat{t}(x)$. Defining the *effective action*, $S_{eff}(\mathbf{W})$, of the exchange field, \mathbf{W} , by

$$S_{eff}(\mathbf{W}) = \ln \det(D\mathbf{W}) + S_K(\mathbf{W}), \quad (3.40)$$

we find that, for example,

$$\Xi(\beta, \mu) = \int \mathcal{D}\mathbf{W} e^{S_{eff}(\mathbf{W})}. \quad (3.41)$$

Note that

$$\frac{\delta}{\delta \mathbf{W}(x, t)} e^{\bar{S}(\bar{C}, C; \mathbf{W})} = - \left(\sum_y K(x - y) \mathbf{W}(y, t) + \mathbf{S}(x, t) \right) e^{\bar{S}(\bar{C}, C; \mathbf{W})}. \quad (3.42)$$

Setting

$$\delta_{\mathbf{W}(x,t)} = \frac{\delta}{\delta \mathbf{W}(x,t)} + \sum_y K(x-y) \mathbf{W}(y,t),$$

the imaginary-time Green function corresponding to a product, $\prod_{j=1}^n S_{\alpha_j}(x_j, t_j)$, of spin operators turns out to be given by

$$\langle \prod_{j=1}^n S_{\alpha_j}(x_j, t_j) \rangle_{\beta, \mu} = (-1)^n \Xi(\beta, \mu)^{-1} \int \mathcal{D}\mathbf{W} \prod_{j=1}^n \delta_{W_{\alpha_j}(x_j, t_j)} e^{S_{eff}(\mathbf{W})}, \quad (3.43)$$

where we have used (3.40) and (3.42).

Our purpose is now to determine $S_{eff}(\mathbf{W})$ as explicitly as possible, in order to get some insight into the R.S. of (3.41) and (3.43).

4 Effective σ -models of the exchange field

If we want to apply the method of steepest descent to estimate the R.S. of (3.41) and of (3.43), we must look for the (absolute and local) maxima of $S_{eff}(\mathbf{W})$. This leads us to consider mean-field ansätze for configurations, \mathbf{W} , corresponding to local maxima of $S_{eff}(\mathbf{W})$.

Ferromagnetic mean-field theory

We set

$$\mathbf{W}(x, t) = W_0 \mathbf{n}, \quad (4.1)$$

independently of x and t , where \mathbf{n} is a unit vector, and $W_0 > 0$. Because of the symmetry of the problem under rotations of \mathbf{W} , we can choose $\mathbf{n} = \mathbf{n}_z$ to point into the z -direction. The problem of evaluating

$$g^{mf}(\beta, \mu; W_0) = \lim_{\Lambda \nearrow \Gamma} -\frac{1}{\beta|\Lambda|} S_{eff}^\Lambda(\mathbf{W} \equiv W_0 \mathbf{n}_z) \quad (4.2)$$

is elementary, because, in momentum space, all modes decouple from one another, and we can appeal to well known calculations (see e.g. [15], [17]) to obtain the answer:

$$\begin{aligned} g^{mf}(\beta, \mu; W_0) &= \frac{1}{2} \hat{K}(0) W_0^2 - \lim_{\Lambda \nearrow \Gamma} \frac{1}{\beta|\Lambda|} \ln \det(D_{\mathbf{W}=W_0 \mathbf{n}}) \\ &= \frac{1}{2} \hat{K}(0) W_0^2 - \frac{1}{\beta} \int_{B_\Gamma} \frac{d\mathbf{k}}{(2\pi)^3} \ln \left[\left(1 + e^{-\beta(\epsilon(\mathbf{k}) + W_0/2)} \right) \times \right. \\ &\quad \left. \times \left(1 + e^{-\beta(\epsilon(\mathbf{k}) - W_0/2)} \right) \right] + \text{const.}, \quad (4.3) \end{aligned}$$

where

$$\epsilon(\mathbf{k}) = t(\mathbf{k}) - \mu. \quad (4.4)$$

Formula (4.3) can also be derived directly from the definition of $\ln \det(D_{\mathbf{W}}) = \text{tr} \ln(D_{\mathbf{W}})$ by using the Poisson summation formula.

In the limit $\beta \rightarrow \infty$, expression (4.3) converges to

$$g^{mf}(\infty, \mu; W_0) = \frac{1}{2} \hat{K}(0) W_0^2 + \int_{B_\Gamma} \frac{d\mathbf{k}}{(2\pi)^3} \left[2\epsilon(\mathbf{k}) \Theta \left(-\frac{W_0}{2} - \epsilon(\mathbf{k}) \right) + \left(\epsilon(\mathbf{k}) - \frac{W_0}{2} \right) \Theta \left(\frac{W_0}{2} - |\epsilon(\mathbf{k})| \right) \right] + \text{const.}, \quad (4.5)$$

which is the ground state energy density of electrons in a constant exchange field $\mathbf{W} = W_0 \mathbf{n}_z$. The value of W_0 minimizing g^{mf} is found by solving the equation $(\partial/\partial W_0)g^{mf}(\beta, \mu; W_0) = 0$, i.e.,

$$\hat{K}(0)W_0 = \frac{1}{2} \int_{B_\Gamma} \frac{d\mathbf{k}}{(2\pi)^3} \left[(e^{\beta(\epsilon(\mathbf{k}) - \frac{W_0}{2})} + 1)^{-1} - (e^{\beta(\epsilon(\mathbf{k}) + \frac{W_0}{2})} + 1)^{-1} \right]. \quad (4.6)$$

When $\beta \rightarrow 0$ the R.S. of (4.6) tends to 0, and we conclude that, for small β , this equation only has the trivial solution, $W_0 = 0$. When $\beta \rightarrow \infty$, equation (4.6) yields the equation

$$\hat{K}(0)W_0 = \frac{1}{2} \int_{B_\Gamma} \frac{d\mathbf{k}}{(2\pi)^3} \Theta \left(\frac{W_0}{2} - |\epsilon(\mathbf{k})| \right). \quad (4.7)$$

Besides the trivial solution, this equation also has a non-trivial solution $W_0 > 0$, provided $\hat{K}(0)$ is sufficiently small (depending on \hat{t}). We recall that

$$\hat{K}(0) = \hat{J}(0)^{-1} = \left(\frac{4U_0}{3} + \hat{J}_\neq(0) \right)^{-1}. \quad (4.8)$$

Thus, for U_0/t_* , with $t_* := \max |\hat{t}(x)|$, large enough, equation (4.6) has a non-trivial solution, provided β is large enough, *even* in the Hubbard model, where $J_\neq = 0$. Note that, by (3.42) and (3.43),

$$-\sum_y K(x-y) \langle \mathbf{W}(y, t) \rangle_{\beta, \mu} = \langle \mathbf{S}(x, t) \rangle_{\beta, \mu}. \quad (4.9)$$

Thus, in mean-field theory,

$$\hat{K}(0)W_0 = M, \quad (4.10)$$

where W_0 is the non-trivial solution of equation (4.6), and M is the spontaneous magnetization in mean-field theory (i.e., $-M\mathbf{n}_z = \langle \mathbf{S}(x, t) \rangle_{\beta, \mu}^{mf}$). Thus, equation (4.7) tells us that, at zero temperature ($\beta \rightarrow \infty$),

$$M = \frac{1}{2} \int_{B_\Gamma} \frac{d\mathbf{k}}{(2\pi)^3} \Theta \left(\frac{\hat{J}(0)}{2} M - |\epsilon(\mathbf{k})| \right), \quad (4.11)$$

where $\hat{J}(0) = \sum_x J(x) = \hat{K}(0)^{-1}$. This equation has an obvious interpretation apparent from Fig. 3.

Matters simplify further for a *flat band*, $t \equiv 0$, i.e., for the Heisenberg model. Then eq. (4.7) reduces to

$$M = \begin{cases} 1/2, & -\hat{J}(0)/4 < \mu < \hat{J}(0)/4, \\ 0, & \text{otherwise,} \end{cases} \quad (4.12)$$

for $\beta \rightarrow \infty$. For, eq. (4.5) implies that

$$g^{mf}(\infty, \mu; W_0) = \frac{1}{2} \hat{K}(0) W_0^2 - \frac{W_0}{2} \Theta \left(\frac{W_0}{2} - |\mu| \right), \quad (4.13)$$




Figure 3: Spin bands

up to an unimportant constant. Thus, for $|\mu| < 1/(4\hat{K}(0)) = \hat{J}(0)/4$, $g^{mf}(\infty, \mu; W_0)$ has a quadratic minimum at $W_0 = 1/(2\hat{K}(0))$, with

$$g^{mf}\left(\infty, \mu; W_0 = \frac{1}{2\hat{K}(0)}\right) = -\frac{1}{2\hat{K}(0)},$$

and

$$\frac{\partial^2 g^{mf}}{\partial W_0^2}\left(\infty, \mu; W_0 \approx \frac{1}{2\hat{K}(0)}\right) \equiv \hat{K}(0). \quad (4.14)$$

Antiferromagnetic mean-field theory

Antiferromagnetism and Néel ordering in Hubbard models is discussed in some detail in [15], so we shall be brief. To simplify matters, we consider the lattices $\Gamma = \mathbb{Z}^d$, $d = 1, 2, 3, \dots$. Our mean-field ansatz configuration for the exchange field \mathbf{W} is given by

$$\mathbf{W}(x, t) = (-1)^{|x|} W_0 \mathbf{n}_z, \quad (4.15)$$

which describes Néel order. We skip detailed calculations and give the result for the free energy

$$g^{mf}(\beta, \mu, W_0) = \frac{1}{2} \hat{K}(\mathbf{k}_\pi) W_0^2 - \frac{1}{\beta} \int_{B_\Gamma} \frac{d\mathbf{k}}{(2\pi)^3} \ln \left[\left(1 + e^{-\beta \sqrt{\epsilon(\mathbf{k})^2 + (1/4)W_0^2}} \right) \left(1 + e^{\beta \sqrt{\epsilon(\mathbf{k})^2 + (1/4)W_0^2}} \right) \right] + \text{const}, \quad (4.16)$$

where $\mathbf{k}_\pi = (\pi, \pi, \dots, \pi)$ denotes a corner of the Brillouin zone. In order to derive (4.16), we assume that $\epsilon(\mathbf{k} + \mathbf{k}_\pi) = -\epsilon(\mathbf{k})$, i.e., we assume the band to be *half-filled*. Note that small deviations $\delta\epsilon(\mathbf{k}) = \epsilon(\mathbf{k} + \mathbf{k}_\pi) + \epsilon(\mathbf{k})$ from half filling can be taken into account with the help of perturbation theory in $\delta\epsilon(\mathbf{k})$. The idea behind the derivation of (4.16) is to consider simultaneously the contributions of \mathbf{k} and $\mathbf{k} + \mathbf{k}_\pi$, diagonalising a 4×4 matrix. The energy eigenvalues, for a fixed \mathbf{k} , are given by $\epsilon_\pm(\mathbf{k}) = \pm \sqrt{\epsilon^2(\mathbf{k}) + (1/4)W_0^2}$, each one with twofold degeneracy. From (4.16) we obtain the mean-field value of W_0 by solving the equation $\partial g^{mf}(\beta, \mu; W_0)/\partial W_0 = 0$. In the limit where $\beta \rightarrow \infty$, this equation reduces to

$$\hat{K}(\mathbf{k}_\pi) = \frac{1}{4} \int_{B_\Gamma} \frac{d\mathbf{k}}{(2\pi)^3} \frac{1}{\sqrt{\epsilon(\mathbf{k})^2 + (1/4)W_0^2}}. \quad (4.17)$$

The R.S. diverges logarithmically when $W_0 \rightarrow 0$, for a lattice Γ of arbitrary dimension. Thus (4.17) has a non-trivial solution $W_0 > 0$ for arbitrarily large values of $\hat{K}(\mathbf{k}_\pi) = (4U_0/3 + J_\neq(\mathbf{k}_\pi))^{-1}$, i.e., for arbitrarily small values of the on-site repulsion U_0 (even if $J_\neq = 0$), and hence the same is true for finite, but sufficiently large values of β . Thus

$$|\epsilon_\pm(\mathbf{k})| \geq \frac{1}{2}W_0 > 0, \quad (4.18)$$

and we conclude that a strictly positive *energy gap* $W_0 > 0$ *opens at the Fermi-surface*.

We note that the solution of (4.17) behaves like

$$W_0 \propto t_* e^{-\alpha \hat{K}(\mathbf{k}_\pi) t_*}, \quad (4.19)$$

for some constant α . For the half-filled Hubbard model ($J_\neq = 0$, $U_0 > 0$), $\hat{K}(\mathbf{k}_\pi) = 3/(4U_0)$, and we find that W_0 has an essential singularity at $U_0 = 0$.

Comparing these results with those obtained for the ferromagnetic mean-field ansatz, we conclude that, at zero temperature, in the Hubbard model with $\hat{K}(0) = \hat{K}(\mathbf{k}_\pi) = 3/(4U_0)$, a non-trivial ferromagnetic mean-field solution does *not* exist when U_0 is small, while an antiferromagnetic mean-field solution exists *for arbitrarily small values* of U_0 at *half filling*, where the (mean-field) ground state energy of the Néel state is found to be *below* the ground state energy of the ferromagnetic state.

The magnetic σ -model

As we have seen, these ansätze correspond to quadratic (local) maxima of the effective action as a function of the *length* W_0 *of the exchange field*. This suggests that the effective action $S_{eff}(\mathbf{W})$ is maximal on exchange-field configurations of constant length, $|\mathbf{W}(x, t)| \approx W_0$. It may therefore be appropriate to study the effective action (3.40) with the constraint

$$|\mathbf{W}(x, t)| = W_0, \quad \text{for all } t \in [0, \beta), \quad x \in \Gamma. \quad (4.20)$$

To study magnetic order, we consider the following *ansatz* for the effective action on configurations of exchange fields of constant length, $|\mathbf{W}(x, t)| = W_0$:

$$\begin{aligned} \tilde{S}_{eff}(\mathbf{W}) = & \int_0^\beta dt \sum_x \left\{ -\frac{W_0^2}{2} \sum_y \hat{\mathbf{W}}(x, t) K(x - y) \hat{\mathbf{W}}(y, t) + C_t |\partial_t \hat{\mathbf{W}}(x, t)|^2 + C_\nabla |\nabla \hat{\mathbf{W}}(x, t)|^2 \right\} \\ & + C_{WZ} \int_{S^{2,N}} dt ds \sum_x \hat{\mathbf{W}}(x, t, s) \cdot (\partial_t \hat{\mathbf{W}}(x, t, s) \wedge \partial_s \hat{\mathbf{W}}(x, t, s)) + \text{higher order terms}, \end{aligned} \quad (4.21)$$

where we have set $\hat{\mathbf{W}}(x, t) := W_0^{-1} \mathbf{W}(x, t)$ and where, for each $x \in \Gamma$, $\hat{\mathbf{W}}(x, t, s)$ is chosen to be a smooth extension of $\hat{\mathbf{W}}(x, t)$ from the circle $\{t | 0 \leq t \leq \beta\}$ to the northern hemisphere, $S^{2,N}$, of the 2-sphere of radius $\beta/(2\pi)$. The operator ∇ denotes the finite difference gradient on the lattice, and the corresponding term in the effective action has to be interpreted as

$$\sum_x |\nabla \hat{\mathbf{W}}(x, t)|^2 := \sum_{|x-y|=1} |\hat{\mathbf{W}}(y, t) - \hat{\mathbf{W}}(x, t)|^2.$$

The functional $\tilde{S}_{eff}(\mathbf{W})$ is the (imaginary-time) action of a *non-linear σ -model* with $SO(3)$ symmetry and with a Wess-Zumino term. This model is called *magnetic σ -model*. It is expected to

describe exchange field configurations \mathbf{W} in thermal equilibrium at inverse temperature β , provided the coefficients C_t , C_∇ and C_{WZ} are chosen in such a way that

$$\tilde{S}_{eff}(\mathbf{W}) \approx S_{eff}(\mathbf{W})|_{|W(x,t)|=W_0} + \text{const} \ , \quad (4.22)$$

on configurations $\mathbf{W}(x, t)$ satisfying the constraint $|\mathbf{W}(x, t)| = W_0$, with $S_{eff}(\mathbf{W})$ as in (3.40). More precisely, if $\mathbf{W} = \mathbf{W}_{mf} + \boldsymbol{\chi}$, where \mathbf{W}_{mf} is a mean(-exchange)-field configuration and $\boldsymbol{\chi}$ is a small perturbation of \mathbf{W}_{mf} , i.e., $\boldsymbol{\chi}(x, t) \perp \mathbf{W}_{mf}(x)$ and $|\boldsymbol{\chi}(x, t)| \ll 1$, for all $x \in \Gamma$ and all times t , then we want (4.22) to hold *exactly* to *second order* in $\boldsymbol{\chi}$. Eq. (4.21) makes it clear that, in order to calculate C_t , C_∇ and C_{WZ} such that (4.22) holds to second order in $\boldsymbol{\chi}$, it suffices to consider *time-independent* exchange fields, $\mathbf{W}(x, t) \equiv \mathbf{W}(x)$, to calculate C_∇ , while it suffices to consider *site-independent* exchange fields, $\mathbf{W}(x, t) \equiv \mathbf{W}(t)$, when calculating C_t and C_{WZ} . Obviously, the *signs* of C_t and C_∇ will determine the *stable* mean-field configuration \mathbf{W}_{mf} . For \mathbf{W}_{mf} to be *independent* of time t , C_t must be *negative* (with our sign convention). For *ferromagnetism* to prevail, C_∇ must be *negative*.

In the next two subsections, the coefficients C_t and C_∇ are determined in such a way that (4.22) holds to second order in $\boldsymbol{\chi}$, for an *appropriate choice* of \mathbf{W}_{mf} (depending on our choice of parameters in the original model). The coefficient C_{WZ} of the Wess-Zumino term will be calculated and shown to be *imaginary* in the next to last subsection of Section 4. The Wess-Zumino term is crucial in arriving at the correct *magnon dynamics* for *ferromagnetically ordered* magnets.

Stability of antiferromagnetic ordering

In this subsection we calculate the coefficients C_t and C_∇ of the magnetic σ -model (4.21), in such a way that (4.22) holds, in a regime where these coefficients can be calculated perturbatively. For the *Hubbard model* this is the regime where the band is *half-filled* and the signs of the coefficients C_t and C_∇ will indeed determine the *stable* mean exchange field to be time-independent and *antiferromagnetically* ordered.

We first derive an innocent looking, but important identity, which will also be useful in the next section. We set

$$D_{\mathbf{W}}^- = -\partial_t + \epsilon - W \ . \quad (4.23)$$

Equation (3.38) shows that

$$D_{\mathbf{W}}^T = -\partial_t + \epsilon + \frac{1}{2}(W_1\sigma_1 - W_2\sigma_2 + W_3\sigma_3) \ ,$$

where the superscript “ T ” stands for transposition, (i.e., taking the adjoint, followed by complex conjugation). Hence

$$\sigma_2 D_{\mathbf{W}}^T \sigma_2 = -\partial_t + \epsilon - W = D_{\mathbf{W}}^- \ . \quad (4.24)$$

Furthermore,

$$D_{\mathbf{W}}^* = -\partial_t + \epsilon + W = D_{-\mathbf{W}}^- \ . \quad (4.25)$$

Since $\sigma_2 = \sigma_2^* = \sigma_2^{-1}$, it follows from (4.24) that

$$\det(D_{\mathbf{W}}) = \det(D_{\mathbf{W}}^T) = \det(\sigma_2 D_{\mathbf{W}}^T \sigma_2) = \det(D_{\mathbf{W}}^-) \ , \quad (4.26)$$

and hence

$$\overline{\det(D_{\mathbf{W}})} = \det(D_{\mathbf{W}}^*) = \det(D_{-\mathbf{W}}^-) = \det(D_{-\mathbf{W}}) \ . \quad (4.27)$$

This identity shows that the real part of the logarithm of the determinant of $D_{\mathbf{W}}$ is given by:

$$\Re \ln \det(D_{\mathbf{W}}) = \frac{1}{2} \ln \det(D_{\mathbf{W}} D_{-\mathbf{W}}),$$

$$D_{\mathbf{W}} D_{-\mathbf{W}} = (\partial_t + \epsilon)^2 - \frac{W_0^2}{4} - [\epsilon, W] - (\partial_t W).$$

We are tempted to treat $[\epsilon, W] + (\partial_t W)$ as a perturbation of

$$\Lambda := (\partial_t + \epsilon)^2 - \frac{W_0^2}{4}. \quad (4.28)$$

This is justified if $|\epsilon| \ll W_0$ and $|\partial_t W| \ll W_0$. In this regime the band of electrons with spin parallel to the background field, \mathbf{W}_{mf} , is filled and the band of electrons with spin anti-parallel to \mathbf{W}_{mf} is empty, i.e., the lattice is half-filled. At half-filling, we expect the *Hubbard model* to exhibit an antiferromagnetic phase at sufficiently low temperatures. We recall that $\ln \det = \text{tr} \ln$ and expand the logarithm up to quadratic order in $[\epsilon, W]/\Lambda$ and $(\partial_t W)/\Lambda$. Terms linear in W vanish at critical points of the effective action; \mathbf{W} -independent terms are omitted. We are then left with

$$\Re \ln \det(D_{\mathbf{W}}) = -\frac{1}{4} \text{tr}(\Lambda^{-1}([\epsilon, W] + (\partial_t W))\Lambda^{-1}([\epsilon, W] + (\partial_t W))) + \text{h.o.}, \quad (4.29)$$

where h.o. stands for terms of higher order in $|\epsilon|/W_0$ or $|\partial_t W|/W_0$. Next, we pull Λ^{-1} through $[\epsilon, W] + \partial_t W$ using the identity

$$[\Lambda^{-1}, [\epsilon, W] + (\partial_t W)] = -\Lambda^{-1}[\Lambda, [\epsilon, W] + (\partial_t W)]\Lambda^{-1}, \quad (4.30)$$

which shows that the commutator only contributes higher-order terms to (4.29). Mixed terms with one spatial and one time derivative of W do not contribute to the trace. We thus arrive at

$$\Re \ln \det(D_{\mathbf{W}}) = -\frac{1}{4} \text{tr}(\Lambda^{-2}([\epsilon, W]^2 + (\partial_t W)^2)) + \text{h.o.} \quad (4.31)$$

Comparison with (4.21) then yields an expression for C_{∇} ,

$$C_{\nabla} = \frac{W_0^2}{8(2\pi)^3} \int_{B_{\Gamma}} d\mathbf{k} |\nabla \epsilon(\mathbf{k})|^2 \frac{1}{\beta} \sum_{k_0} \frac{1}{((-ik_0 + \epsilon(\mathbf{k}))^2 - W_0^2/4)^2} \xrightarrow{\beta W_0 \rightarrow \infty} \frac{1}{4(2\pi)^3 W_0} \int_{B_{\Gamma}} d\mathbf{k} |\nabla \epsilon(\mathbf{k})|^2,$$

which is positive. Thus the real part of the σ -model action (4.21) for the Hubbard-model exhibits its maximum (w.r.t. time-independent exchange fields) on configurations for which

$$\int_0^{\beta} dt \sum_x |\nabla \hat{\mathbf{W}}(x, t)|^2$$

is *maximal*, and this is the case for a Néel ordered (staggered) exchange field.

From (4.31) we can also read off the C_t -coefficient in (4.21) to be given by

$$C_t = -\frac{W_0^2}{8} \frac{1}{\beta} \sum_{k_0} \frac{1}{((-ik_0 + \epsilon(\mathbf{k}))^2 - W_0^2/4)^2} \xrightarrow{\beta W_0 \rightarrow \infty} -\frac{1}{4W_0},$$

which is manifestly negative. From these calculations and the mean field equation (4.17) we conclude that the *Hubbard model at half filling* exhibits a *stable* antiferromagnetic phase if the on-site repulsion U_0 is large enough as compared to the hopping term, and if the temperature is sufficiently low.

Linear stability of the ferromagnetic mean exchange field

In this subsection we analyze conditions that imply stability of *ferromagnetically* ordered exchange fields. In the *Hubbard model*, we expect a stable ferromagnetic phase to prevail when the band is weakly filled and fairly flat. In such a regime, since there is no energy gap at the Fermi surface, the operator Λ defined in (4.28) has zeros and the perturbative expansion of the last subsection breaks down. We therefore have to resort to a linear stability analysis around a ferromagnetically ordered mean exchange field. We will see that the stability of the ferromagnetically ordered exchange field is due to contributions close to the Fermi surface.

The stability of a ferromagnetically ordered exchange field w.r.t. x -independent fluctuations can be shown quite easily, since the operator $D_{\mathbf{W}}$ is diagonal in momentum space when $\mathbf{W}(x, t) = \mathbf{W}(t)$ is x -independent. We can then write

$$\lim_{\Lambda \nearrow \Gamma} \frac{1}{|\Lambda|} \ln \det(D_{\mathbf{W}}) = \int_{B_{\Gamma}} \frac{d\mathbf{p}}{(2\pi)^3} \ln \det \left(\partial_t + \hat{\epsilon}(\mathbf{p}) + \frac{\mathbf{W}(t)}{2} \cdot \boldsymbol{\sigma} \right). \quad (4.32)$$

In the operator formalism outlined at the beginning of Section 3,

$$\det \left(\partial_t + \hat{\epsilon}(\mathbf{p}) + \frac{\mathbf{W}(t)}{2} \cdot \boldsymbol{\sigma} \right) = \text{const} \times \text{Tr}_{\mathbb{C}^4} U_{\epsilon, \mathbf{W}}(\beta, 0),$$

where

$$\partial_t U_{\epsilon, \mathbf{W}}(t, s) = h_{\epsilon, \mathbf{W}}(t) U_{\epsilon, \mathbf{W}}(t, s), \quad (4.33)$$

$$U_{\epsilon, \mathbf{W}}(t, t) = \mathbf{1}_4, \quad (4.34)$$

and

$$h_{\epsilon, \mathbf{W}}(t) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \hat{\epsilon}(\mathbf{p}) \mathbf{1}_2 + \frac{\mathbf{W}(t)}{2} \cdot \boldsymbol{\sigma} & 0 \\ 0 & 0 & 0 & 2\hat{\epsilon}(\mathbf{p}) \end{pmatrix}.$$

The trace of the propagator $U_{\epsilon, \mathbf{W}}(\beta, 0)$ is

$$\text{Tr}_{\mathbb{C}^4} U_{\epsilon, \mathbf{W}}(\beta, 0) = 1 + e^{-\beta \hat{\epsilon}(\mathbf{p})} \text{Tr}_{\mathbb{C}^2} U_{\epsilon=0, \mathbf{W}}(\beta, 0) + e^{-2\beta \hat{\epsilon}(\mathbf{p})}. \quad (4.35)$$

Using the Hölder-inequality for traces (see e.g. [21]), we find that

$$\left| \text{Tr}_{\mathbb{C}^2} (U_{0, \mathbf{W}}(\beta, 0)) \right| \leq \text{Tr}_{\mathbb{C}^2} (U_{0, \mathbf{W}=W_0 \cdot \mathbf{n}}(\beta, 0)), \quad (4.36)$$

for an arbitrary *constant* unit vector \mathbf{n} . From this inequality we can conclude, that x -independent fluctuations of an exchange field of constant length around a ferromagnetic mean-field configuration do *not* decrease the free energy and hence are not a source of instability.

Since the calculation is rather easy, we give an explicit formula for the coefficient C_t of the magnetic σ -model when the exchange field is ferromagnetically ordered. Because of the stability of the system with respect to x -independent fluctuations we must find a negative sign for this coefficient. We use the Hölder inequality (4.36), in order to get an upper bound for the large- β behaviour of the trace in (4.35) and find, for large β , and up to a \mathbf{W} -independent constant

$$S_{eff}(\mathbf{W}) \approx (V_- - V_+) \ln \det(\partial_t + W) + S_K(\mathbf{W}), \quad (4.37)$$

where $V_{\pm} = \text{Vol}\{\mathbf{p}|\hat{\epsilon}(\mathbf{p}) \pm W_0/2 < 0\}$. In order to calculate C_t we take a look at the real part of $\ln \det(\partial_t + W)$,

$$\Re \ln \det(\partial_t + W) = \frac{1}{2} \ln \det[(\partial_t + W)(-\partial_t + W)] = \frac{1}{2} \ln \det(\Lambda + (\partial_t W)),$$

where $\Lambda := -\partial_t^2 + (1/4)W_0^2$. We treat $(\partial_t W)$ as a perturbation of Λ and proceed in the same way as in the last subsection to arrive at

$$\Re \ln \det(\partial_t + W) = -\frac{1}{4} \text{tr}(\Lambda^{-2}(\partial_t W)^2) + \text{h.o.},$$

where we omitted \mathbf{W} -independent terms. Finally, we find for the coefficient C_t

$$C_t = -\frac{W_0^2(V_- - V_+)}{8(2\pi)^3\beta} \sum_{k_0} \frac{1}{(k_0^2 + W_0^2/4)^2} \xrightarrow{\beta W_0 \rightarrow \infty} -\frac{V_- - V_+}{4(2\pi)^3 W_0}, \quad (4.38)$$

which is indeed manifestly negative.

Next, we investigate the stability of the ferromagnetically ordered mean-field configurations w.r.t. time-independent, but x -dependent perturbations, which amounts to the calculation of C_{∇} in the magnetic σ -model. This calculation is somewhat harder, because the p -dependent modes in the operator $D_{\mathbf{W}}$ do not decouple. We perturb the ferromagnetic mean-field configuration $\mathbf{W} = W_0 \mathbf{n}_z$ by time-independent fluctuations, $\boldsymbol{\chi}$, and calculate their contribution to the real part of the effective action to second order in $|\boldsymbol{\chi}|/W_0$. Since the effective action has a quadratic minimum w.r.t. the length of the exchange field, we can assume $\boldsymbol{\chi}$ to be orthogonal to \mathbf{n}_z . Then (omitting $\boldsymbol{\chi}$ -independent terms)

$$\Re S_{eff}(W_0 \mathbf{n}_z + \boldsymbol{\chi}) = -\frac{1}{2} \sum_{x,y} \boldsymbol{\chi}^T(x) K(x-y) \boldsymbol{\chi}(y) - \frac{1}{4} \text{tr}(\Lambda^{-1}\{\epsilon, \boldsymbol{\chi}\} \Lambda^{-1}\{\epsilon, \boldsymbol{\chi}\}) + \text{h.o.}, \quad (4.39)$$

where

$$\Lambda = -\partial_t^2 + \epsilon^2 + \frac{W_0^2}{4} + W_0 \epsilon \sigma_3, \\ \chi = \frac{1}{2} \boldsymbol{\chi} \cdot \boldsymbol{\sigma}.$$

In the trace we only keep contributions quadratic in $|\mathbf{p}|/W_0$, where \mathbf{p} is the momentum labelling the modes of $\boldsymbol{\chi}$ (derivative expansion). Spin degrees of freedom and Matsubara frequencies can be summed over explicitly, and - after a straightforward but tedious calculation - we obtain a formula for the coefficient, C_{∇} , of the term

$$\frac{\beta}{W_0^2} \sum_x |\nabla \boldsymbol{\chi}(x)|^2$$

in the expansion of the second term on the R.S. of (4.39), which, for large values of β , is given by

$$C_{\nabla} \approx (2\pi)^{-3} \int_{B_F} d\mathbf{k} \frac{|\nabla \epsilon(\mathbf{k})|^2}{4W_0} \left\{ -\Theta\left(\epsilon(\mathbf{k}) - \frac{W_0}{2}\right) \frac{\beta W_0}{4} \frac{1}{\cosh \beta \left(\epsilon(\mathbf{k}) - \frac{W_0}{2}\right) + 1} \right. \\ \left. + \Theta\left(\frac{W_0}{2} - |\epsilon(\mathbf{k})|\right) \left[1 - \frac{\beta W_0}{4} \left(\frac{1}{\cosh \beta \left(\epsilon(\mathbf{k}) + \frac{W_0}{2}\right) + 1} + \frac{1}{\cosh \beta \left(\epsilon(\mathbf{k}) - \frac{W_0}{2}\right) + 1} \right) \right] \right. \\ \left. - \Theta\left(-\epsilon(\mathbf{k}) - \frac{W_0}{2}\right) \frac{\beta W_0}{4} \frac{1}{\cosh \beta \left(\epsilon(\mathbf{k}) + \frac{W_0}{2}\right) + 1} \right\}. \quad (4.40)$$

By plugging $\mathbf{W} = W_0 \mathbf{n}_z + \boldsymbol{\chi}$, with $\boldsymbol{\chi}$ orthogonal to \mathbf{n}_z , into the σ -model action (4.21), one easily sees that the coefficient C_∇ just derived coincides with the corresponding coefficient in the σ -model.

For the *Hubbard model* we can deduce sufficient conditions for the linear stability of the ferromagnetic mean-field configuration (i.e., for C_∇ to be *negative*) from (4.40), in the regime where β is large. We denote by v_F^\pm the Fermi velocities of the upper/lower spin band, namely

$$|\epsilon(\mathbf{k}) \pm \frac{W_0}{2}| \approx v_F^\pm |\mathbf{k} - \mathbf{k}_F^\pm|, \quad \text{for } \mathbf{k} \text{ close to the Fermi momenta } \mathbf{k}_F^\pm,$$

by A_F^\pm the area of the Fermi surface of the upper/lower spin band, and by V_F^\pm the respective volumes. Then, for large β , a sufficient condition for linear stability is that

$$\frac{W_0}{2} (A_F^+ v_F^+ + A_F^- v_F^-) > v_{\max}^2 (V_F^- - V_F^+), \quad (4.41)$$

where $v_{\max} = \max\{|\nabla \epsilon(\mathbf{k})| \mid |\epsilon(\mathbf{k}) - W_0/2| < 0 < \epsilon(\mathbf{k}) + W_0/2\}$.

We now combine this stability condition with the mean-field equation (4.7) for the Hubbard-model, i.e.,

$$W_0 = \frac{2U_0}{3} \frac{V_F^- - V_F^+}{(2\pi)^3},$$

to arrive at the stability condition

$$\frac{U_0}{3} \frac{A_F^+ v_F^+ + A_F^- v_F^-}{(2\pi)^3} > v_{\max}^2.$$

Details of these and other related calculations will appear in [18].

Magnon dynamics and the WZ -term

In this subsection we consider the Wess-Zumino-term in (4.21). We first consider an x -independent exchange field $\mathbf{W}(t)$ of constant length $|\mathbf{W}(t)| = W_0$ and calculate the variation of the first term on the R.S. of (3.40) for $\epsilon = 0$, given a variation, $\delta \mathbf{W}$, of the exchange field:

$$\delta \ln \det(D_{0,\mathbf{W}}) = \text{tr}(\delta D_{0,\mathbf{W}} (D_{0,\mathbf{W}}^* D_{0,\mathbf{W}})^{-1} D_{0,\mathbf{W}}^*), \quad (4.42)$$

where

$$\delta D_{0,\mathbf{W}} = \frac{1}{2} \delta \mathbf{W}(t) \cdot \boldsymbol{\sigma} \equiv \delta W(t).$$

In order to evaluate the R.S. of (4.42), we expand $(D_{0,\mathbf{W}}^* D_{0,\mathbf{W}})^{-1}$ in a Neumann series in powers of $(\partial_t W)$,

$$(D_{0,\mathbf{W}}^* D_{0,\mathbf{W}})^{-1} = \Lambda^{-1} + \Lambda^{-1} (\partial_t W) \Lambda^{-1} + \dots,$$

where $\Lambda = -\partial_t^2 + (1/4)W_0^2$. Plugging this series into (4.42), we find that

$$\delta \ln \det(D_{0,\mathbf{W}}) = \delta S^{(1)}(\mathbf{W}) + \delta S^{(2)}(\mathbf{W}) + \dots \quad (4.43)$$

where

$$\delta S^{(1)}(\mathbf{W}) = \text{tr}(\delta W \Lambda^{-1} D_{0,\mathbf{W}}^*), \quad (4.44)$$

$$\delta S^{(2)}(\mathbf{W}) = \text{tr}(\delta W \Lambda^{-1} (\partial_t W) \Lambda^{-1} D_{0,\mathbf{W}}^*), \quad (4.45)$$

etc.. Using the cyclicity of the trace and the fact that $\text{Tr}(\boldsymbol{\sigma}) = 0$ (where “Tr” is the trace on 2×2 matrices), we see that

$$\delta S^{(1)}(\mathbf{W}) = \alpha(W_0) \int_0^\beta dt \text{Tr}(W(t) \delta W(t)),$$

where

$$\alpha(W_0) = \beta^{-1} \sum_{k_0 \in Z_\beta} \frac{1}{k_0^2 + (1/4)W_0^2},$$

and $Z_\beta = (\pi/\beta)(2\mathbb{Z} + 1)$. Since

$$\text{Tr}(W(t) \delta W(t)) = \frac{1}{2} \delta \text{Tr}(W(t)^2),$$

$\delta S^{(1)}$ only depends on variations of the length of $\mathbf{W}(t)$ and *vanishes* if the length is held fixed.

By (4.45),

$$\delta S^{(2)}(\mathbf{W}) = \delta S_I^{(2)}(\mathbf{W}) + \delta S_{II}^{(2)}(\mathbf{W}),$$

where

$$\delta S_I^{(2)}(\mathbf{W}) = -\text{tr}(\delta W \Lambda^{-1} (\partial_t W) \Lambda^{-1} \partial_t) = -\frac{1}{2} \sum_{j=1}^3 \text{tr}(\delta W_j \Lambda^{-1} (\partial_t W)_j \Lambda^{-1} \partial_t)$$

and

$$\delta S_{II}^{(2)}(\mathbf{W}) = \text{tr}(\delta W \Lambda^{-1} (\partial_t W) \Lambda^{-1} W) = -\frac{i}{4} \sum_{j,l,m} \epsilon_{jlm} \text{tr}(\delta W_j \Lambda^{-1} (\partial_t W)_l \Lambda^{-1} W_m).$$

Here we have used that $\text{Tr}(\sigma_j \sigma_l) = 2\delta_{jl}$ and $\text{Tr}(\sigma_j \sigma_l \sigma_m) = -2i\epsilon_{jlm}$. Since ∂_t , Λ and $W_j(t)$ are real operators, $\delta S_I^{(2)}$ is real and $\delta S_{II}^{(2)}$ purely imaginary. The real part has been calculated in the last subsection. Turning to $\delta S_{II}^{(2)}(\mathbf{W})$, we observe that

$$\begin{aligned} \delta S_{II}^{(2)}(\mathbf{W}) &= \text{tr}(W \delta W (\partial_t W) \Lambda^{-2}) + \text{higher derivative terms} \\ &= -\frac{i}{4} \gamma(W_0) \int_0^\beta dt \sum_{j,l,m} \epsilon_{jlm} W_j(t) \delta W_l(t) \partial_t W_m(t) + \text{h.d.t.} \\ &= -\frac{i}{2} \int_0^\beta dt \hat{\mathbf{W}}(t) \cdot (\delta \hat{\mathbf{W}}(t) \wedge \partial_t \hat{\mathbf{W}}(t)) + \mathcal{O}((\beta W_0)^{-1}) + \text{h.d.t.}, \end{aligned} \quad (4.46)$$

where, in the last equation, we have used that

$$\gamma(W_0) := \beta^{-1} \sum_{k_0 \in Z_\beta} \frac{1}{(k_0^2 + (1/4)W_0^2)^2} \xrightarrow{\beta W_0 \rightarrow \infty} \frac{2}{W_0^3}. \quad (4.47)$$

We note that $\text{h.d.t.} \sim \mathcal{O}((\beta W_0)^{-1})$.

Recall that $\mathbf{W}(t)$ has periodic boundary conditions at $t = 0, \beta$. Viewing the imaginary-time circle $[0, \beta)$ as the equator of a sphere S_β^2 of radius $\beta/(2\pi)$, we may extend the unit-vector field $\hat{\mathbf{W}}(t)$ from the equator of S_β^2 to a continuous unit-vector field $\hat{\mathbf{W}}(t, s)$, $0 \leq s \leq \beta/2$, on the entire sphere in an

arbitrary way, but with $\hat{\mathbf{W}}(t, \beta/4) = \hat{\mathbf{W}}(t)$. The first term on the R.S. of (4.46) then turns out to be the variation of the functional

$$\begin{aligned} S_{WZ}(\mathbf{W}) &= -\frac{i}{2} \int_0^\beta dt \int_0^{\beta/4} ds \hat{\mathbf{W}}(t, s) \cdot (\partial_s \hat{\mathbf{W}}(t, s) \wedge \partial_t \hat{\mathbf{W}}(t, s)) \\ &= -\frac{i}{2} \int_{S^{2,N}} \hat{\mathbf{W}} \cdot (d\hat{\mathbf{W}} \wedge d\hat{\mathbf{W}}), \end{aligned} \quad (4.48)$$

where “ d ” denotes the exterior derivative, and $S^{2,N}$ is the northern hemisphere of S^2 . We recognize (4.48) to be the Wess-Zumino term in (4.21) for an x -independent (i.e., ferromagnetically ordered) exchange field. The corresponding coefficient C_{WZ} depends on ϵ and will be determined later on. The R.S. of (4.48) is expressed in standard differential-form notation, which makes it manifest that $S_{WZ}(\mathbf{W})$ is *independent* of the radius, $\beta/2\pi$, of the sphere.

The leading term on the R.S. of (4.46) is also the variation of

$$S'_{WZ}(\mathbf{W}) = \frac{i}{2} \int_{S^{2,S}} \hat{\mathbf{W}} \cdot (d\hat{\mathbf{W}} \wedge d\hat{\mathbf{W}}),$$

where $S^{2,S}$ is the southern hemisphere of S^2 . One observes that

$$S_{WZ}(\mathbf{W}) - S'_{WZ}(\mathbf{W}) = -\frac{i}{2} \int_{S^2} \hat{\mathbf{W}} \cdot (d\hat{\mathbf{W}} \wedge d\hat{\mathbf{W}}) = 2\pi i n, \quad (4.49)$$

where $n = n(\hat{\mathbf{W}}) \in \mathbb{Z}$ is the degree of the map $\hat{\mathbf{W}} : S^2 \rightarrow S^2$. It follows that, apparently, the Wess-Zumino action S_{WZ} is only determined *modulo an integer multiple of $2\pi i$* . Due to (4.49), $\exp(S_{WZ}(\mathbf{W}))$ is a *single-valued* functional of \mathbf{W} .

If the exchange field $\mathbf{W}(x, t)$ is *antiferromagnetically* ordered, i.e., the Néel field

$$\mathbf{N}(x, t) \equiv (-1)^{|x|} \mathbf{W}(x, t) = W_0 \mathbf{n}$$

is a constant field pointing in the direction of some unit vector \mathbf{n} , the Wess-Zumino term in (4.21) obviously tends to 0 in the formal continuum limit. It follows that, in antiferromagnetically ordered states, at sufficiently low temperatures, there are gapless Goldstone bosons with dispersion

$$\omega(\mathbf{k}) \approx v|\mathbf{k}|, \quad (|\mathbf{k}| \approx 0), \quad (4.50)$$

for some velocity v .

In *ferromagnetically ordered systems*, however, the Wess-Zumino term does *not* disappear. Using (4.37) and (4.48), the coefficient C_{WZ} in (4.21) is found to be

$$C_{WZ} = -\frac{i}{2} \frac{(V_- - V_+)}{(2\pi)^3}, \quad (4.51)$$

in the limit where $\beta W_0 \rightarrow \infty$. Had we *normalized* the Fourier modes and were we in a finite volume, we would find that $\exp(\tilde{S}_{eff}(\mathbf{W}))$ is single-valued. From (4.21) we derive the equations of motion

$$C_{WZ} \hat{\mathbf{W}}_x \wedge \dot{\hat{\mathbf{W}}}_x = \sum_y \tilde{K}(x - y) \hat{\mathbf{W}}_y + \alpha \hat{\mathbf{W}}_x, \quad (4.52)$$

with $\hat{\mathbf{W}}_x(t) = \hat{\mathbf{W}}(x, t)$, $\dot{\hat{\mathbf{W}}}_x = \partial_t \hat{\mathbf{W}}_x$, where $\tilde{K}(x - y) = W_0^2 K(x - y) + C_\nabla \delta_{|x-y|,1}$, and where α is a Lagrange multiplier arising from the constraint $|\hat{\mathbf{W}}_x(t)|^2 = 1$, for all x and all t . Taking the vector product of (4.52) with $\hat{\mathbf{W}}_x$, and doing the Wick rotation from imaginary time back to real time [23], we obtain the equations of motion for *magnons* in a ferromagnet

$$iC_{WZ} \dot{\hat{\mathbf{W}}}_x = \hat{\mathbf{W}}_x \wedge \sum_y \tilde{K}(x - y) \hat{\mathbf{W}}_y, \quad (4.53)$$

which is the well known *Landau-Lifshitz equation*; see also [19].

If J , and hence K , are of short range and if the equilibrium state of the system were ferromagnetically ordered, i.e.,

$$\mathbf{W}(x, t) = W_0 \mathbf{n} + \boldsymbol{\chi}(x, t), \quad W_0 > 0,$$

where \mathbf{n} is a unit vector and $\boldsymbol{\chi}(x, t)$ is assumed to be small, then the dispersion of magnons is found to be

$$\omega(\mathbf{k}) = \frac{|\mathbf{k}|^2}{2M}, \quad \text{for some constant } M > 0,$$

as expected. If \mathbf{n} points in the z -direction then $\boldsymbol{\chi}$ lies essentially in the $x - y$ plane. We define

$$\phi(x, t) = \chi_1(x, t) + i\chi_2(x, t).$$

Passing to the formal continuum limit, the action functional for the complex magnon field ϕ of a ferromagnet is seen to be given by

$$S_{\text{magnon}}(\bar{\phi}, \phi) = \int_0^\beta dt \int dx [\bar{\phi}(x, t) \partial_t \phi(x, t) + \frac{1}{2M} \bar{\phi}(x, t) (\Delta \phi)(x, t)], \quad (4.54)$$

which is the action for a system of *conserved non-relativistic bosons* with vanishing chemical potential. Since $W_1 = \chi_1 = (\phi + \bar{\phi})/2$ is an observable field, the chemical potential actually necessarily vanishes, and there is no “magnon condensation”; (just like photons do not form any Bose condensates).

Absence of symmetry breaking in one and two dimensions at positive temperatures

For $\beta < \infty$, we can study the fluctuations of the modes $\mathbf{W}(k_0, \mathbf{k})$ of the exchange field with $k_0 = 0$,

i.e., of $\int_0^\beta \mathbf{W}(x, t) dt$.

We set

$$\int_0^\beta \mathbf{W}(x, t) dt = \beta(\mathbf{W}^{mf} + \chi_l(x) \mathbf{n}_z + \boldsymbol{\chi}_t(x)),$$

where \mathbf{W}^{mf} is a solution of the ferromagnetic or the antiferromagnetic mean-field equation. Then, in d spatial dimensions, we conclude from our stability analysis

$$\langle \chi_t(x)^2 \rangle_{\beta, \mu} \propto \int_{B_\Gamma} \frac{d^d k}{k^2}$$

which, for $d = 1$ and 2 , is infrared-divergent. Thus, transversal fluctuations around the mean field are gigantic and hence destroy long-range order; (this is the Mermin-Wagner theorem, see, e.g., [6]).

In two dimensions, there is an alternative way to understand the absence of ordering when $\beta < \infty$: Setting $\mathbf{W}(x, t) = \tilde{\mathbf{W}}(x)$ (ferromagnetic short-range order), or $\mathbf{W}(x, t) = (-1)^{|x|} \tilde{\mathbf{W}}(x)$ (antiferromagnetic short-range order), where $\tilde{\mathbf{W}}(x)$ is slowly varying, and passing to the formal continuum limit (lattice spacing $\rightarrow 0$), one finds that the action $S_{eff}(\mathbf{W})$ has (approximate) critical points, indeed local maxima, on time-independent configurations $\tilde{\mathbf{W}}(x)$ with the properties:

$$|\tilde{\mathbf{W}}(x)| \approx W_0, \quad \text{for all } x,$$

where W_0 is a solution of the (ferromagnetic or antiferromagnetic) mean-field equation,

$$\tilde{\mathbf{W}}(x) \rightarrow W_0 \mathbf{n}, \quad \text{as } |x| \rightarrow \infty,$$

where \mathbf{n} is an arbitrary unit vector, and the degree of the map $\tilde{\mathbf{W}}(x)$, as measured by the integer (“winding number”)

$$\frac{1}{4\pi W_0^3} \int d^2x \tilde{\mathbf{W}}(x) \cdot \left(\frac{\partial \tilde{\mathbf{W}}(x)}{\partial x_1} \wedge \frac{\partial \tilde{\mathbf{W}}(x)}{\partial x_2} \right),$$

is non-zero. Such configurations are called “instantons”. The contributions of instanton configurations to a functional integral, such as (3.43), destroy long-range order.

Remark. Generally speaking, it is hard to justify the use of steepest descent in approximately evaluating functional integrals, such as (3.41) and (3.43), because there isn’t any large constant $\mathcal{N} = \hbar^{-1}$ multiplying $S_{eff}(\mathbf{W})$. If, however, we place \mathcal{N} identical species of spin- $\frac{1}{2}$ -fermions on each site $x \in \Gamma$, coupled to each other only through exchange interactions between the total spin operators, and if we set $K_0 = \mathcal{N}K_0$ then $S_{eff}(\mathbf{W}) \equiv S_{eff}^{(K_0)}(\mathbf{W})$ is replaced by $\mathcal{N}S_{eff}^{(K_0)}(\mathbf{W})$, and steepest descent becomes reliable.

5 Reflection positivity and phase transitions in Heisenberg-models

In order to establish the existence of phase transitions in some of the models introduced in Section 3, we recall the method of *reflection positivity* ([5],[4],[20],[21]). For simplicity, we choose the lattice Γ to be given by \mathbb{Z}^d . We *decompose* Γ *into two equal halves*

$$\Gamma_+ = \{x \in \Gamma | x_1 > 0\}, \quad \Gamma_- = \{x \in \Gamma | x_1 \leq 0\}, \quad (5.1)$$

where x_j is the j^{th} component of $x \in \Gamma$. Let Π be the plane in \mathbb{R}^d orthogonal to the 1-direction at height $x_1 = \frac{1}{2}$, and let ϑ denote reflection at Π . Clearly, ϑ maps Γ_- onto Γ_+ , and conversely.

An exchange coupling matrix, $J(x, y)$, is said to be *reflection positive* iff

$$\sum_{x, y} \overline{f(x)} J(x, \vartheta y) f(y) \geq 0, \quad (5.2)$$

for all functions f on Γ with $\text{supp } f \subseteq \Gamma_+$. Note that this condition is *independent* of the diagonal elements, $J(x, x)$, of J ; so, by choosing them appropriately, we can always ensure that J is either positive-definite or negative-definite. By computing some integrals with a Gaussian measure of mean 0 and covariance $-J^{-1}$ (J negative-definite) we see that if J is reflection positive then so is $-J^{-1}$; see e.g. [20],[21]. Furthermore, if J is reflection-positive then so is $-\tilde{J}$, where \tilde{J} is given by

$$\tilde{J}(x, y) = (-1)^{|x+y|} J(x, y) = (-1)^{|x|+|y|} J(x, y) \quad (5.3)$$

with $|x| = \sum_{j=1}^d x_j$. (Note that if J is positive-definite then \tilde{J} is positive-definite, too.) The following simple calculation proves our claim. Let

$$\tilde{f}(x) = (-1)^{|x|} f(x).$$

Since J is assumed to be reflection-positive,

$$\begin{aligned} 0 &\leq \sum_{x,y} \overline{\tilde{f}(x)} J(x, \vartheta y) \tilde{f}(y) \\ &= \sum_{x,y} \overline{f(x)} (-1)^{|x|+|y|} J(x, \vartheta y) f(y) \\ &= - \sum_{x,y} \overline{f(x)} (-1)^{|x+\vartheta y|} J(x, \vartheta y) f(y) \\ &= - \sum_{x,y} \overline{f(x)} \tilde{J}(x, \vartheta y) f(y), \end{aligned}$$

for an arbitrary function f with $\text{supp } f \subseteq \Gamma_+$ (hence $\text{supp } \tilde{f} \subseteq \Gamma_+$). These considerations are summarized as follows:

$$\text{If } J \text{ is reflection-positive then } -K := -J^{-1} \text{ is reflection-positive, too;} \quad (5.4)$$

and

$$\text{if } -J \text{ is reflection-positive (i.e., } J \text{ is reflection-negative)}$$

$$\text{then } -\tilde{K} := -\widetilde{J^{-1}} \text{ is reflection-positive, too.} \quad (5.5)$$

In both situations, we can choose the diagonal elements of J such that J and K are positive-, or negative-definite.

A reflection-positive matrix J is called *ferromagnetic*, while a reflection-negative matrix is called *antiferromagnetic*. It is easy to see that, for nearest-neighbour exchange couplings, these notions of “ferro-” and “antiferromagnetic” coincide with the familiar ones.

Let J be ferromagnetic and negative-definite. Then the Gaussian measure

$$d\mu_{-J}(\phi) = \text{const} \cdot e^{\frac{1}{2} \sum_{x,y} \phi(x) J(x,y) \phi(y)} \prod_x d^n \phi(x),$$

where $\phi(x) \in \mathbb{R}^n$, $n = 1, 2, 3, \dots$, and the constant is chosen such that $d\mu_{-J}$ is normalized, is *reflection-positive*, in the sense that

$$\int d\mu_{-J}(\phi) \overline{(\Theta F)(\phi)} F(\phi) \geq 0, \quad (5.6)$$

for an arbitrary bounded function F that depends only on the variables $\{\phi(x) | x \in \Gamma_+\}$; the space of all such functions is denoted by \mathcal{F}_+ . The function ΘF in (5.6) is defined by

$$(\Theta F)(\phi) = F(\phi_\vartheta), \quad (5.7)$$

where $\phi_\vartheta(x) = \phi(\vartheta x)$. If $\phi^{(1)}, \dots, \phi^{(l)}$ are independent Gaussian random fields with distributions given by $d\mu_{-J^{(t)}}(\phi^{(t)})$, where $J^{(t)}$ is ferromagnetic and $J^{(t)} < 0$, $t = 1, \dots, l$, then the distribution

$$d\mu(\phi) = \prod_{t=1}^l d\mu_{-\tau^{(t)} J^{(t)}}(\phi^{(t)}) \quad (5.8)$$

of the random field ϕ given by

$$\phi(x, t) = \phi^{(t)}(x), \quad t = 1, \dots, l,$$

is reflection-positive, for arbitrary $\tau^{(t)} > 0$, $t = 1, \dots, l$; as is easily seen.

These considerations have the following consequence that will play a crucial role in our analysis of phase transitions: Let $\mathbf{W}(x, t)$, $x \in \Gamma$, $t \in [0, \beta)$, with $\mathbf{W}(x, t + \beta) = \mathbf{W}(x, t)$, be the *imaginary time exchange field*, and let $d\mu_K(\mathbf{W}) = \exp(S_K(\mathbf{W}))\mathcal{D}\mathbf{W}$ be its Gaussian distribution, as introduced in Section 3, (3.27) through (3.31). If the exchange couplings J are *ferromagnetic* (in the sense specified above) and positive-definite then $-K = -J^{-1}$ is ferromagnetic and negative-definite; hence the Gaussian measure

$$d\mu_K(\mathbf{W}) \text{ is reflection-positive,} \quad (5.9)$$

as follows from (3.29), (5.6) and (5.8).

Next, we suppose that the exchange couplings J are *antiferromagnetic* (in the sense specified above). Then, by (5.5), $-\tilde{K} = -\widetilde{J^{-1}}$ is *reflection-positive* and negative-definite, for J positive-definite. We introduce a random field, \mathbf{N} , (“N” for “Néel”), by setting

$$\mathbf{N}(x, t) = (-1)^{|x|}\mathbf{W}(x, t). \quad (5.10)$$

If \mathbf{W} has distribution $d\mu_K(\mathbf{W})$, with $K = J^{-1}$, then \mathbf{N} has distribution $d\mu_{\tilde{K}}(\mathbf{N})$, and hence

$$d\mu_{\tilde{K}}(\mathbf{N}) \text{ is reflection-positive.} \quad (5.11)$$

We set

$$\mathbf{N}_\vartheta(x, t) = \mathbf{N}(\vartheta x, t), \quad (5.12)$$

and (see (5.7))

$$(\Theta F)(\mathbf{N}) = F(\mathbf{N}_\vartheta), \quad (5.13)$$

for an arbitrary function F of $\{\mathbf{N}(x, t) | x \in \Gamma, t \in [0, \beta)\}$.

We consider the models introduced in Section 3 without hopping term but with \mathcal{N} Fermion species. They have an action functional given by

$$\begin{aligned} \bar{S}(\bar{C}, C; \mathbf{W}) = & \sum_{a=1}^{\mathcal{N}} \int_0^\beta dt \sum_x \left[\bar{C}_s^{(a)}(x, t) \left(\frac{\partial}{\partial t} - \mu \right) C_s^{(a)}(x, t) - \nu_a \mathbf{W}(x, t) \cdot \mathbf{S}^{(a)}(x, t) \right] \\ & - \frac{1}{2} \int_0^\beta dt \left(\sum_{x,y} \mathbf{W}(x, t) K(x-y) \mathbf{W}(y, t) \right), \end{aligned} \quad (5.14)$$

where $\nu_a = \pm 1$, for all $a = 1, \dots, \mathcal{N}$, and $\hat{K}(k) = \left(\frac{4}{3}U_0 + \hat{J}_\neq(k) \right)^{-1}$; see (3.20), (3.29), and (3.32) through (3.34). Since the first term on the R.S. of (5.14) does not couple different sites in the lattice, the effective action of the exchange field \mathbf{W} , see (3.36) and (3.40), is given by

$$S_{eff}(\mathbf{W}) = -\frac{1}{2} \int_0^\beta dt \left(\sum_{x,y} \mathbf{W}(x, t) K(x-y) \mathbf{W}(y, t) \right) + \sum_a \sum_x \ln \det(D_{\nu_a} \mathbf{W}_{(x, \cdot)}), \quad (5.15)$$

with $D_{\mathbf{W}}$ given in (3.38), for $\hat{t}(x) \equiv 0$. We define

$$\begin{aligned} f_x(\mathbf{N}) &= \prod_{a=1}^{\mathcal{N}} \det(D_{\nu_a}(-1)^{|x|} \mathbf{N}(x, \cdot)) \\ &= \prod_{a=1}^{\mathcal{N}} \det(D_{\nu_a} \mathbf{W}(x, \cdot)), \end{aligned} \quad (5.16)$$

Then it follows that

$$\begin{aligned} \overline{\Theta f_x(\mathbf{N})} &= \overline{f_x(\mathbf{N}_{\vartheta}(x, \cdot))} \\ &= \prod_{a=1}^{\mathcal{N}} \overline{\det(D_{\nu_a}(-1)^{|x|} \mathbf{N}_{\vartheta}(x, \cdot))} \\ &= \prod_{a=1}^{\mathcal{N}} \overline{\det(D_{-\nu_a}(-1)^{|\vartheta x|} \mathbf{N}(\vartheta x, \cdot))} \\ &= \prod_{a=1}^{\mathcal{N}} \det(D_{\nu_a}(-1)^{|\vartheta x|} \mathbf{N}(\vartheta x, \cdot)) \\ &= f_{\vartheta x}(\mathbf{N}), \end{aligned} \quad (5.17)$$

where, in the third equation, we have used that $(-1)^{|x|} = -(-1)^{|\vartheta x|}$, and, in the fourth equation, we have inserted identity (4.27). Equation (5.17) implies that

$$\overline{\Theta \left(\prod_{x \in \Gamma_+} f_x(\mathbf{N}) \right)} = \prod_{x \in \Gamma_-} f_x(\mathbf{N}). \quad (5.18)$$

From this equation and equations (5.11) and (5.6) we conclude the following simple, but quite fundamental

Result A Suppose the exchange couplings J are *antiferromagnetic* (in the sense introduced above) and positive-definite. We set

$$d\mu_{eff}(\mathbf{N}) = \text{const} \cdot \prod_x f_x(\mathbf{N}) d\mu_{\tilde{K}}(\mathbf{N}) \quad (5.19)$$

$$= \text{const}' \cdot \exp \left(S_{eff}(\mathbf{N}) \right) \mathcal{D}\mathbf{N}, \quad (5.20)$$

where the constants are chosen such that $\int d\mu_{eff}(\mathbf{N}) = 1$, and

$$\begin{aligned} S_{eff}(\mathbf{N}) &:= -\frac{1}{2} \int_0^b dt \left(\sum_{x,y} \mathbf{N}(x, t) \tilde{K}(x-y) \mathbf{N}(y, t) \right) \\ &\quad + \sum_a \sum_x \ln \det \left(D_{\nu_a}(-1)^{|x|} \mathbf{N}(x, \cdot) \right), \end{aligned} \quad (5.21)$$

with $\tilde{K} = \widetilde{J^{-1}}$; see (5.15), (5.10), (5.11).

Then the measure $d\mu_{eff}(\mathbf{N})$ is *reflection-positive*, in the sense that

$$\int d\mu_{eff}(\mathbf{N}) \overline{(\Theta F)(\mathbf{N})} F(\mathbf{N}) \geq 0, \quad (5.22)$$

for arbitrary functions $F \in \mathcal{F}_+$ (where \mathcal{F}_+ is defined right above (5.7)).

Next, we suppose that the number of fermion species, $\mathcal{N} = 2\mathcal{M}$ is *even*, with

$$\nu_1 = \dots = \nu_{\mathcal{M}} = 1, \nu_{\mathcal{M}+1} = \dots = \nu_{2\mathcal{M}} = -1. \quad (5.23)$$

We define

$$g_x(\mathbf{W}) = \det(D_{\mathbf{W}(x,\cdot)})^{\mathcal{M}} \det(D_{-\mathbf{W}(x,\cdot)})^{\mathcal{M}}.$$

Identity (4.27) then implies that $g_x(\mathbf{W}) \geq 0$, for all $x \in \Gamma$, and, setting

$$\mathbf{W}_{\vartheta}(x, t) = \mathbf{W}(\vartheta x, t),$$

we find that

$$\overline{\Theta g_x(\mathbf{W})} = \Theta g_x(\mathbf{W}) = g_{\vartheta x}(\mathbf{W}). \quad (5.24)$$

Defining

$$d\mu_{eff}(\mathbf{W}) = \text{const} \cdot \prod_x g_x(\mathbf{W}) d\mu_K(\mathbf{W}) \quad (5.25)$$

$$= \text{const}' \cdot \exp\left(S_{eff}(\mathbf{W})\right) \mathcal{D}\mathbf{W}, \quad (5.26)$$

where $K = J^{-1}$, $S_{eff}(\mathbf{W})$ is as in (5.15), and the constants are chosen such that $\int d\mu_{eff}(\mathbf{W}) = 1$, we arrive at

Result F Suppose the exchange couplings J are *ferromagnetic* (in the sense introduced above). Then the measure $d\mu_{eff}(\mathbf{W})$ is *reflection-positive*, i.e.

$$\int d\mu_{eff}(\mathbf{W}) \overline{(\Theta F)(\mathbf{W})} F(\mathbf{W}) \geq 0, \quad (5.27)$$

for arbitrary functions $F \in \mathcal{F}_+$.

From Results A and F one obtains the following *infrared (spin-wave) bounds*, originally discovered in [5] and generalized in [20],[21].

(IRA) Under the hypotheses of Result A (in particular, J *antiferromagnetic*), one has the inequalities

$$\begin{aligned} 0 &\leq \langle \widehat{\mathbf{N}}(\mathbf{k}, k_0) \cdot \widehat{\mathbf{N}}(-\mathbf{k}, -k_0) \rangle_{\beta, \mu} \\ &\leq \beta W_0^2 \delta_0(\mathbf{k}) \delta_{k_0, 0} + \left(\widehat{K}(\mathbf{k}) - \widehat{K}(0) \right)^{-1} \end{aligned} \quad (5.28)$$

for some constant $W_0^2 \geq 0$. Note that $\widehat{\mathbf{N}}(-\mathbf{k}, -k_0) = \overline{\widehat{\mathbf{N}}(\mathbf{k}, k_0)}$, because $\mathbf{N}(x, t)$ is real.

If W_0^2 is strictly positive then the state $\langle (\cdot) \rangle_{\beta, \mu}$ exhibits *long-range order*, and if we couple \mathbf{N} to an arbitrarily small external field, $\varepsilon \mathbf{n}$, $|\mathbf{n}| = 1$, then

$$\lim_{\varepsilon \rightarrow 0} \langle \mathbf{N}(x, t) \rangle_{\beta, \mu, \varepsilon \mathbf{n}} = W_0 \mathbf{n}.^3 \quad (5.29)$$

³This claim is heuristic, but can be replaced by an equivalent, mathematically rigorous one, [20].

Inequality (5.28) implies that, for an arbitrary bounded function $f(x, t)$ of rapid decay in $x \in \Gamma$ and periodic in $t \in [0, \beta)$,

$$\begin{aligned} 0 &\leq \langle |\mathbf{N}(f)|^2 \rangle_{\beta, \mu} \\ &\leq \beta W_0^2 |\hat{f}(0, 0)|^2 + \sum_{k_0 \in \frac{2\pi}{\beta} \mathbb{Z}} \int_{B_\Gamma} d\mathbf{k} \frac{|\hat{f}(\mathbf{k}, k_0)|^2}{\widehat{\tilde{K}}(\mathbf{k}) - \widehat{\tilde{K}}(0)}, \end{aligned} \quad (5.30)$$

where $\mathbf{N}(f) = \int_0^\beta dt \sum_x \mathbf{N}(x, t) f(x, t)$.

Returning to identities (3.42) and (3.43), with

$$\mathbf{S}(x, t) = \sum_{a=1}^{\mathcal{N}} \nu_a \mathbf{S}^{(a)}(x, t), \quad (5.31)$$

and recalling that $\mathbf{W}(x, t) = (-1)^{|x|} \mathbf{N}(x, t)$, we conclude from (5.30) that

$$0 \leq \langle (\int_0^\beta \mathbf{S}(0, t) dt)^2 \rangle_{\beta, \mu} \quad (5.32)$$

$$\leq \beta \tilde{K}(0) + \beta W_0^2 \widehat{\tilde{K}}(0)^2 + \int_{B_\Gamma} d\mathbf{k} \frac{\widehat{\tilde{K}}(\mathbf{k})^2}{\widehat{\tilde{K}}(\mathbf{k}) - \widehat{\tilde{K}}(0)}. \quad (5.33)$$

We recall that $-\tilde{K}$ is reflection-positive, and the diagonal elements are chosen such that \tilde{K} (or, equivalently, K and J) is positive-definite. Therefore $\widehat{\tilde{K}}(\mathbf{k}) > 0$, for all $\mathbf{k} \in B_\Gamma$. Furthermore, if J is of short range then \tilde{K} is of short range, too, i.e., $\tilde{K}(x)$ decreases rapidly in $|x|$, and, by reflection-positivity of $-\tilde{K}$, $\widehat{\tilde{K}}(0) \leq \widehat{\tilde{K}}(\mathbf{k})$. It follows that

$$0 \leq \widehat{\tilde{K}}(\mathbf{k}) - \widehat{\tilde{K}}(0) \leq \kappa |\mathbf{k}|^2, \quad (5.34)$$

for some constant κ . Thus, under the assumptions just specified, the third term on the R.H.S. of (5.32) is bounded above by a *finite* constant, $I(K, d)$, in dimension $d \geq 3$, but is (*infrared-*) *divergent* in one and two dimensions; ($I(K, d = 1, 2)$ is finite *only* if J is of *long-range*; see [20],[21]).

A fairly elementary inequality due to Bruch and Falk, see [4], yields the following lower bound

$$\langle (\int_0^\beta \mathbf{S}(0, t) dt)^2 \rangle_{\beta, \mu} \geq C(\beta, \mu, \mathcal{N}) \beta \quad (5.35)$$

for some constant $C(\beta, \mu, \mathcal{N}) \geq 0$. If the on-site repulsion U_0 , the chemical potential μ and the value of $\tilde{K}^{-1}(0) = \int_{B_\Gamma} d\mathbf{k} \widehat{\tilde{K}}^{-1}(\mathbf{k})$ are tuned appropriately such that, at zero temperature, the band of each species of fermions is half-filled then $C(\beta, \mu, \mathcal{N}) > 0$ and $\lim_{\mathcal{N} \rightarrow \infty} C(\beta, \mu, \mathcal{N}) = \infty$.

Thus, we conclude that, for appropriate choices of U_0 , μ and $\tilde{K}^{-1}(0)$, and for a *sufficiently large* number, \mathcal{N} , of fermion species,

$$W_0 \text{ is strictly positive, for } \beta \text{ large enough} \quad (5.36)$$

Hence, by (3.42) and (5.29),

$$\begin{aligned}
\lim_{\varepsilon \rightarrow 0} \langle \mathbf{S}(x, 0) \rangle_{\beta, \mu, \varepsilon \mathbf{n}} &= - \sum_y K(x-y) \lim_{\varepsilon \rightarrow 0} \langle \mathbf{W}(y, 0) \rangle_{\beta, \mu, \varepsilon \mathbf{n}} \\
&= -(-1)^{|x|} \sum_y \tilde{K}(x-y) \lim_{\varepsilon \rightarrow 0} \langle \mathbf{N}(y, 0) \rangle_{\beta, \mu, \varepsilon \mathbf{n}} \\
&= -(-1)^{|x|} \widehat{\tilde{K}}(0) W_0 \mathbf{n} \neq 0,
\end{aligned} \tag{5.37}$$

with $M = \widehat{\tilde{K}}(0) |W_0|$ the local magnetization, i.e., the system exhibits *Néel order*, for sufficiently large values of β and \mathcal{N} . A similar result has first been proven in [4].

Arguments related to those used in the proof of the Mermin-Wagner theorem (see [6]) show that if $M > 0$ then there are gapless spin waves (see e.g. [24]), in accordance with the mean-field picture of Section 4.

Next, we exploit Result F. It implies that, under condition (5.23), the following infrared bounds hold.

(IRF) Under the hypotheses of Result F (in particular, J *ferromagnetic* and (5.23) holds), the inequalities

$$\begin{aligned}
0 &\leq \langle \widehat{\mathbf{W}}(\mathbf{k}, k_0) \cdot \widehat{\mathbf{W}}(-\mathbf{k}, -k_0) \rangle_{\beta, \mu} \\
&\leq \beta W_0^2 \delta_0(\mathbf{k}) \delta_{k_0, 0} + \left(\widehat{K}(\mathbf{k}) - \widehat{K}(0) \right)^{-1}
\end{aligned} \tag{5.38}$$

for a constant $W_0^2 \geq 0$, are valid. (See [5],[20],[21]). The consequences of these inequalities are perfectly analogous to those discussed for antiferromagnets above. We set

$$\mathbf{S}(x, t) = \sum_{a=1}^{\mathcal{M}} \left(\mathbf{S}^{(a)}(x, t) - \mathbf{S}^{(a+\mathcal{M})}(x, t) \right), \tag{5.39}$$

see (5.23). Then, under appropriate conditions on U_0 , μ and $K^{-1}(0) = J(0)$, and for $d \geq 3$,

$$M = \lim_{\varepsilon \rightarrow 0} \langle \mathbf{S}(x, t) \rangle_{\beta, \mu, \varepsilon \mathbf{n}} = -\widehat{K}(0) W_0 \mathbf{n} \neq 0 \tag{5.40}$$

for β and \mathcal{M} large enough, which proves long-range order at low temperatures, for \mathcal{M} large enough; (presumably $\mathcal{N} = 2\mathcal{M} = 2$ suffices if J is nearest-neighbour and $d \geq 3$).

Unfortunately, there is, as yet, no such result on the existence of a phase transition in a quantum Heisenberg *ferromagnet* with only one species of fermions (electrons), i.e., $\mathcal{N} = 1$. The reason is that this model is *not* described by a reflection-positive distribution, $d\mu_{eff}(\mathbf{W})$, for the exchange field \mathbf{W} . Setting

$$f_x(\mathbf{W}) = \det(D_{\mathbf{W}(x, \cdot)}), \tag{5.41}$$

$d\mu_{eff}$ is given by

$$d\mu_{eff}(\mathbf{W}) = \text{const} \cdot \prod_x f_x(\mathbf{W}) d\mu_K(\mathbf{W}). \tag{5.42}$$

Now, if J is ferromagnetic $d\mu_K(\mathbf{W})$ is reflection-positive if we set $\mathbf{W}_\vartheta(x, t) = \mathbf{W}(\vartheta x, t)$. However, using (4.27),

$$\begin{aligned}
\overline{\Theta f_x(\mathbf{W})} &= \overline{\det(D_{\mathbf{W}(\vartheta x, \cdot)})} \\
&= \det(D_{-\mathbf{W}(\vartheta x, \cdot)}) \\
&= f_{\vartheta x}(-\mathbf{W}) \neq f_{\vartheta x}(\mathbf{W}).
\end{aligned}$$

Thus, $d\mu_{eff}(\mathbf{W})$, as given in (5.42), is *not* reflection-positive, and there is therefore *no* reason why the Infrared Bounds (IRF) should hold. A very similar problem is encountered in the study of *Bose-Einstein condensation* in lattice models of non-relativistic, interacting bosons.

In order to get some preliminary insights into phase transitions for the Heisenberg *ferromagnet*, one may consider the following “*static approximation*”: one replaces $f_x(\mathbf{W})$ by

$$f_x^{(0)}(\mathbf{W}) = \det(D_{\mathbf{W}^{(0)}(x)}), \quad (5.43)$$

where

$$\mathbf{W}^{(0)}(x) = \frac{1}{\beta} \int_0^\beta dt \mathbf{W}(x, t) \quad \left(= \frac{1}{\beta} \mathbf{W}(x, k_0 = 0) \right),$$

which is time-independent. Then

$$\overline{\Theta f_x^{(0)}(\mathbf{W})} = f_{\vartheta x}^{(0)}(-\mathbf{W}) = f_{\vartheta x}^{(0)}(\mathbf{W}), \quad (5.44)$$

as is easily checked, and $f_x^{(0)}(\mathbf{W}) > 0$. Replacing $d\mu_{eff}(\mathbf{W})$ by

$$d\mu_{eff}^{(0)}(\mathbf{W}) = \prod_x f_x^{(0)}(\mathbf{W}) d\mu_K(\mathbf{W}),$$

we conclude that $d\mu_{eff}^{(0)}(\mathbf{W})$ is reflection-positive if J is ferromagnetic, hence the infrared bounds (5.38) hold, and we conclude that, in the *static approximation*, for appropriate choices of U_0 , μ and $J(0)$, a phase transition accompanied by continuous symmetry breaking occurs at sufficiently low temperatures.

If one replaces electrons with spin $\frac{1}{2}$ by fermions with spin s , or by $2s$ identical species of spin- $\frac{1}{2}$ fermions (with $\nu_a = 1$, for all $a = 1, \dots, 2s$), and if one then takes $s \rightarrow \infty$ (rescaling the spin operators, $\mathbf{S}(x, t) \mapsto \frac{1}{s} \mathbf{S}(x, t)$) then the static approximation becomes *exact*. This is not surprising, because the limit $s \rightarrow \infty$ corresponds to the *classical limit*, as shown in [25], and the spin- s Heisenberg model approaches the classical Heisenberg model for which the existence of a phase transition has been established in [5]. Our formalism, involving the exchange field \mathbf{W} , offers a neat and simple way of recovering some of the results in [25]; (we leave this as an exercise to the reader).

Remarks.

1. We have outlined a proof of existence of a phase transition for a class of Heisenberg *antiferromagnets*; (the proof is mathematically rigorous; some missing details can be inferred from [5], [4]). Of course, this is not a surprise. In [4], Dyson, Lieb and Simon have already proven a similar result *also* using reflection positivity (albeit in a different manner). Their method of proof, too, breaks down for quantum *ferromagnets*.
2. It appears that to understand phase transitions in quantum *ferromagnets* and the related phenomenon of *Bose-Einstein condensation* for lattice gases of non-relativistic bosons in a *mathematically rigorous way*, we would have to resort to a full-fledged renormalization group analysis - a technically rather demanding task.
3. The fact that the function $f_x(\mathbf{W})$ defined in (5.41) (see also (5.19)) is neither positive, nor even real is the origin of the “*sign (complex phase) problem*” in numerical simulations of quantum ferro- and antiferromagnets involving, e.g., the Monte Carlo method applied to (5.42). The great quality of methods based on reflection positivity is that they are applicable to the analysis of certain *complex* distributions $d\mu_{eff}$.

4. If one considers models in d dimensions with $SO(2n)$ spins, $n = 1, 2, 3, \dots$, then for $d \geq 3$ one can prove a ferromagnetic phase transition, using arguments very similar to those described above.
5. After completion of this paper, work of Bach et al. [27] appeared to which we draw the reader's attention.

References

- [1] W. Pauli, *Über Gasentartung und Paramagnetismus*, Z. Physik **41**, 81 (1927).
- [2] P.W. Anderson, *New approach to the theory of superexchange interactions*, Phys. Rev. **115**, 2-13 (1959).
- [3] N. Datta, R. Fernández, J. Fröhlich, *Effective Hamiltonians and phase diagrams for tight-binding models*, J. Stat. Phys. **96**, 545-611 (1999).
- [4] F. Dyson, E. H. Lieb and B. Simon, *Phase transitions in quantum spin systems with isotropic and non-isotropic interactions*, J. Stat. Phys. **18**, 335 (1978).
- [5] J. Fröhlich, B. Simon and T. Spencer, *Infrared Bounds, Phase Transitions And Continuous Symmetry Breaking*, Commun. Math. Phys. **50**, 79 (1976).
- [6] N.D. Mermin, H. Wagner, *Absence of ferromagnetism or antiferromagnetism in one- or two-dimensional isotropic Heisenberg models*, Phys. Rev. Lett. **17**, 1133-36 (1966).
- [7] W. Heisenberg, Z. Physik **49**, 619 (1928).
F. Bloch, Z. Physik **61**, 206 (1930).
E.C. Stoner, Proc. R. Soc. **A165**, 372 (1938).
F.J. Dyson, *General Theory of Spin-Wave Interactions*, Phys. Rev. **102**, 1217-1230 (1956).
F.J. Dyson, *Thermodynamic Behavior of an Ideal Ferromagnet*, Phys. Rev. **102**, 1230-1244 (1956).
L.D. Landau, E.M. Lifshitz, *Course of Theoretical Physics Volume 9, Statistical Physics*, Part 2, by E.M. Lifshitz and L.P. Pitajewski, London: Pergamon 1981.
- [8] E. Brezin, J. C. Le Guillou and J. Zinn-Justin, *Field Theoretical Approach To Critical Phenomena*, in *Phase Transitions and Critical Phenomena*, C. Domb and M.S. Green (eds.) Volume 6, 125-247, London: Academic Press 1976.
D.J. Amit, *Field Theory, the Renormalization Group and Critical Phenomena*, New York a.o.: McGraw-Hill 1978.
- [9] A. Mielke, H. Tasaki, *Ferromagnetism in the Hubbard model*, Commun. Math. Phys. **158**, 341-71 (1993).
- [10] H. Tasaki, *Ferromagnetism in the Hubbard model: A constructive approach*, Commun. Math. Phys. **242**, 445-72 (2003).
- [11] J. Fröhlich, D. Ueltschi, *Hund's Rule and Metallic Ferromagnetism*, J. Stat. Phys. **118**, 973-996 (2005).

- [12] W. Weber, S. Riesen and H. C. Siegmann, *Magnetization Precession by Hot Spin Injection*, Science **291**, 1015, (2001).
- [13] D. Oberli, *The ferromagnetic spin filter*, PhD thesis ETH Zürich, Nr. 12933, (1998).
- [14] J. Fröhlich, U. M. Studer and E. Thiran, *Quantum theory of large systems of nonrelativistic matter*, in *Fluctuating geometries in statistical mechanics and field theory* (Les Houches 1994), F. David, P. Ginsparg, J. Zinn-Justin (eds.); Elsevier 1996, [arXiv:cond-mat/9508062].
- [15] E. Fradkin, *Field theories of condensed matter systems*, Redwood City, California a.o.: Addison-Wesley 1994.
- [16] J. Reuteler, *QM particles scattering off a dynamical target, and decoherence*, ETH-diploma thesis, spring 2005.
- [17] T. Chen, J. Fröhlich and M. Seifert, *Renormalization group methods: Landau-Fermi liquid and BCS superconductor*, in *Fluctuating geometries in statistical mechanics and field theory*, F. David, P. Ginsparg, J. Zinn-Justin (eds.); Elsevier 1996, [arXiv:cond-mat/9508063].
- [18] M. Azam, L. Ferrari, J. Fröhlich, to appear.
- [19] H. Leutwyler, *Nonrelativistic effective Lagrangians*, Phys. Rev. D **49** 3033 (1994), [arXiv:hep-ph/9311264].
- [20] J. Fröhlich, R. Israel, E.H. Lieb, B. Simon, *Phase Transitions and Reflection Positivity I. General Theory and Long Range Lattice Models*, Commun. Math. Phys. **62**, 1-34, (1978).
- [21] J. Fröhlich, *Mathematical Physics of Phase Transitions and Symmetry Breaking*, Bulletin of the American Mathematical Society, **84**, 165-193 (1978).
- [22] D. Ruelle, *Statistical mechanics: rigorous results*, Singapore: World Scientific, 1999.
- [23] L. Birke, J. Fröhlich, *KMS, ETC*. Rev. Math. Phys. **14**, 829-872, (2002), [arXiv:math-ph/0204023].
- [24] J. Fröhlich and T. Spencer, *Phase transitions in statistical mechanics and quantum field theory*, in *New Developments in Quantum Field Theory and Statistical Mechanics*; M. Lévy, P. Mitter (eds.), New York, London: Plenum 1977.
- [25] E.H. Lieb, *The Classical Limit of Quantum Spin Systems*, Commun. Math. Phys. **31**, 327-340, (1973).
- [26] J. Dereziński, V. Jaksic and C. Pillet, *Perturbation theory of W^* -dynamics, Liouvilleans and KMS-states*, Rev. Math. Phys. **15** 447, (2003).
V. Bach, J. Fröhlich and I. M. Sigal, *Return to equilibrium*, J. Math. Phys. **41** 3985 (2000) .
J. Fröhlich, M. Merkli, *Another Return of 'Return to Equilibrium'*, Commun. Math. Phys. **251**, 235-262, (2004).
- [27] V. Bach, E. H. Lieb, J.P. Solovej, *Generalized Hartree-Fock Theory and the Hubbard Model*, [arXiv:cond-mat/9312044].