

New asymptotic behaviour of the surface-atom force out of thermal equilibrium

Mauro Antezza^a, Lev P. Pitaevskii^{a,b} and Sandro Stringari^{a,c}^a Dipartimento di Fisica, Università di Trento and BEC Center,
INFN-CNR, Via Sommarive 14, I-38050 Povo, Trento, Italy^b Kapitza Institute for Physical Problems, ul. Kosygina 2, 119334 Moscow, Russia^c Ecole Normale Supérieure and Collège de France,
Laboratoire Kastler Brossel, 24 Rue Lhomond 75231 Paris, France

(Dated: April 14, 2024)

The Casimir-Polder-Lifshitz force felt by an atom near the surface of a substrate is calculated out of thermal equilibrium in terms of the dielectric function of the material and of the atomic polarizability. The new force decays like $1/z^3$ at large distances (i.e. slower than at equilibrium), exhibits a sizable temperature dependence and is attractive or repulsive depending on whether the temperature of the substrate is higher or smaller than the one of the environment. Our predictions can be relevant for experiments with ultracold atomic gases. Both dielectric and metal substrates are considered.

PACS numbers: 34.50.Dy, 12.20.-m, 42.50.Vk, 42.50.Nn

An intense effort has been devoted in the recent years to study the force felt by an atom near the surface of a substrate (see, for example, [1, 2] and references therein). These studies are motivated both by the possibility of technological applications [3] as well as by the fundamental search for stronger constraints on hypothetical non Newtonian forces [4]. Experimental and theoretical research has also recently focused on the forces acting on ultracold atomic gases, including atomic beams [5, 6, 7], Bose-Einstein condensates [2, 8, 9, 10] and degenerate Fermi gases [11].

The force generated by the surface contains in general two independent components. The first one is related to zero-point fluctuations of the electromagnetic field. At short distances z (typically less than fractions of microns) this force behaves like $1/z^4$ and is the analog of the van der Waals-London interatomic force. At larger distances the inclusion of relativistic retardation effects gives rise to a different $1/z^5$ dependence characterizing the so-called Casimir-Polder regime [12, 13, 14]. Notice that both the van der Waals-London and Casimir-Polder forces depend on the temperature only through the dielectric properties of the substrate. The resulting temperature dependence is usually negligible.

The second component of the force is due to the thermal fluctuations of the electromagnetic field. This effect was first considered by Lifshitz [15] who employed the theory of electromagnetic fluctuations developed by Rytov [16]. We will refer to it as to the Lifshitz force. At distances larger than the thermal photon wavelength $\lambda_T = \hbar c/k_B T$ (corresponding to $7.6 \mu\text{m}$ at room temperature) this force decays, at thermal equilibrium, like $1/z^4$, is attractive, is proportional to the temperature and provides the leading contribution to the total force. Conversely, the zero-point fluctuation component prevails at smaller distances. In the present paper we are interested in the thermal component of the force. The Lifshitz force was originally evaluated at full thermodynamic equilibrium. A non trivial issue is the study of the force out

of thermal equilibrium [17, 18], characterizing configurations where the temperature of the substrate and the one of the surrounding walls located at large distances (hereafter called environment temperature) do not coincide. The problem is relevant both for practical reasons, due to the possibility of tuning the two temperatures independently, and for a better understanding of the interplay between zero-point and thermal fluctuation effects. A first important investigation of the surface-atom force out of thermal equilibrium was carried out by Henkel et al. [17] who calculated the force generated by a dielectric substrate at finite temperature by assuming that the environment temperature is zero. The principal motivation of that paper was the study of the force at short distances. In this Letter we show that, out of thermal equilibrium, the force acting on the atom exhibits a new asymptotic behaviour, characterized by a $1/z^3$ decay at large distances.

Let us consider an atom placed in vacuum at distance z from the flat surface of a substrate made of a material with dielectric function $\epsilon(\omega) = \epsilon^0(\omega) + i\epsilon''(\omega)$. We choose a coordinate system with the xy plane coinciding with the interface and the z axis such that the substrate occupies the region with $z < 0$ and the vacuum the region with $z > 0$. In this Letter we assume that the substrate is locally at thermal equilibrium at a temperature T_S which can differ from the environment temperature T_E , the global system being out of thermal equilibrium, but in a stationary regime. The total electromagnetic field will be in general the sum of the radiation produced by the substrate and the one of the environment. In particular this latter radiation will be partially absorbed and re-emitted from the substrate. The forces produced by the T_E and T_S components of the radiation add incoherently. In typical experiments with ultracold atomic gases the environment temperature is determined by the chamber containing the substrate and the trapped atoms. We further treat the atoms as being at zero temperature in the sense that the surrounding radiation is not able to popu-




FIG. 1: Surface-atom potential energy $V^{neq}(z) = \int_z^{R+1} dz' F^{neq}(z')$ calculated from eq.(9), for different thermal configurations.




FIG. 2: Surface-atom force $F^{neq}(z)$ calculated from eq.(9), for different thermal configurations.

the radiation of the substrate and of the environment. Their additivity can be checked by evaluating separately the two contributions and verifying that their sum, at thermal equilibrium, reproduces the Lifshitz force F_{th}^{eq} [22]. The full surface-atom force out of equilibrium can be finally written in the convenient form

$$F^{neq}(T_S; T_E; z) = F^{eq}(T_E; z) + F_{th}^{neq}(T_S; 0; z) - F_{th}^{neq}(T_E; 0; z) \quad (9)$$

where the equilibrium force $F^{eq}(T; z)$ is given by (3) while $F_{th}^{neq}(T; 0; z)$ is defined by eq.(8).

In figs. 1, 2 and 3 we show the explicit results for the potential energy, the force and the gradient of the force obtained starting from eq.(9) as a function of the




FIG. 3: Surface-atom gradient of the force $-\partial F^{neq}(z)/\partial z$ calculated from eq.(9), for different thermal configurations.

distance from the surface for different choices of T_S and T_E . Calculations have been done for a sapphire substrate and for rubidium atoms. For $F^{eq}(T; z)$ we have used the predictions of [2]. The figure clearly shows that the thermal effects out of equilibrium are sizable (solid lines), thereby providing promising perspectives for future measurements of the surface atom force at large distances. In particular in order to increase the attractive nature of the force it is much more convenient to heat the substrate by keeping the environment at room temperature (lower solid line) rather than heating the whole system (dashed line). When $T_S < T_E$ (upper solid line) the force exhibits a characteristic change of sign reflecting a repulsive nature at large distances (see also discussion below). At short distances the thermal correction to the force becomes smaller and smaller and is determined by the temperature of the substrate. We have reported the results for the potential, for the force and for the gradient of the force because the corresponding predictions can be of interest for different types of experiments with ultra cold gases. Experiments based on the study of the center of mass oscillation of a trapped gas are sensitive to the gradient of the force [2]. The corresponding frequency shifts produced by the surface-atom interaction have been recently measured [10] in conditions of thermal equilibrium in agreement with the predictions of theory [2]. Conversely experiments based on Bloch oscillations are sensitive to the force itself [11, 23]. Finally one can also think at interference experiments with Bose-Einstein condensates in a double well potential. For large separations between the wells the position of the corresponding interference fringes are sensitive to the potential [24].

In the last part of the letter we discuss the large z behaviour of the out of equilibrium force. In this limit only values $q \ll 1$ are important in the evaluation of the integral (8). By making the substitution $q^2 = 1 - \ell^2$; and

the $z \rightarrow 1$ expansion we find that the force (8) exhibits the non trivial asymptotic behaviour

$$F_{th}^{neq}(T;0;z)_{z \rightarrow 1} = \frac{\sim 0}{z^3} \frac{1}{c} \int_0^{\infty} d! \frac{!}{e^{-!} = k_B T} f(!) : \quad (10)$$

Notice that the force exhibits a slower $1=z^3$ decay with respect to the one holding at thermal equilibrium where it decays like $1=z^4$ (see eq. (5)). In the above equation we have introduced the function

$$f(!) = (j'(!) - 1)j + (j''(!) - 1)j^2 \frac{2 + j'(!) - 1}{2j'(!) - 1} \quad (11)$$

which depends on the optical properties of the substrate. For temperatures much smaller than the energy $\sim !_c = k_B$, where $!_c$ is the lowest characteristic frequency of the dielectric substrate, we can replace $f(!)$ with its low frequency limit $(j''_0 + 1) = \frac{j''_0}{j'_0} + 1$. The force (9) felt by the atom then approaches the asymptotic behaviour

$$F^{neq}(T_S;T_E;z)_{z \rightarrow 1} = \frac{0}{6} \frac{k_B^2 (T_S^2 - T_E^2)}{z^3 c} \frac{j''_0 + 1}{j'_0} \quad (12)$$

holding at low temperature and at distances larger than $\lambda_T = \frac{c}{T}$ where λ_T is the thermal photon wave length calculated at the relevant temperatures T_S and T_E [25]. Eq.(12) shows that, at large distances, the new force is attractive or repulsive depending on whether the substrate temperature is higher or smaller than the environment one. Furthermore it exhibits a stronger temperature dependence with respect to equilibrium and contains explicitly the Planck constant. The new dependence of

$F^{neq}(T;0;z)$ on temperature and distance can be physically understood by noticing that the main contribution to the z -th dependent part of the electric energy U_E arises from the black-body radiation impinging on the surface in a small interval of angles, of order of $(\lambda_T = z)^2$, near the angle of total reflection. This radiation creates slowly damping evanescent waves in vacuum. As a result $F^{neq}(T;0;z)$ turns out to be, in accordance with eq.(12), of order of $(\frac{2}{T} = z^3) U_{BB}$, where U_{BB} / T^4 is the energy density of the black-body radiation.

Equation (12) holds for a dielectric substrate where j''_0 is finite. For a metal, if one uses the Drude model, one has $j''_0(!) = 4 - !^2$ with the real part $j'_0(!)$ remaining finite as $! \rightarrow 0$ so that one finds $f(!) \rightarrow \frac{j''_0(!) - 1}{2j'_0(!) - 1} = \frac{1}{2} = 0.5$. At low temperatures eq.(10) then gives rise to a different temperature dependence

$$F^{neq}(T_S;T_E;z)_{z \rightarrow 1} = \frac{0}{z^3 c} \frac{(3=2)^{3=2} - k_B^{3=2} (T_S^{3=2} - T_E^{3=2})}{2} \quad (13)$$

where $(3=2) = 2.61$ is the usual Riemann function.

In conclusion in this Letter we have calculated the surface-atom force out of thermal equilibrium and pointed out the occurrence of a new asymptotic behaviour at large distances. Our predictions could be tested in experiments with ultracold atomic gases trapped close to the surface of a substrate.

We are grateful to E. Cornell, J. Obrecht, J. McGuirk and D. M. Harber for many useful comments. It is also a pleasure to thank I. Carusotto, C. Henkel and S. Reynaud for insightful discussions. Financial support from MURST is acknowledged.

-
- [1] J.F. Babb et al., Phys. Rev. A 70, 042901 (2004).
 [2] M. Antezza, L.P. Pitaevskii, and S. Stringari, Phys. Rev. A 70, 053619 (2004).
 [3] H.B. Chan et al., Phys. Rev. Lett. 87, 211801 (2001).
 [4] S.D. Inopoulis and A.A. Geraci, Phys. Rev. D 68, 124021 (2003). R.S. Decca et al., Annals Phys. 318, 37-80 (2005).
 [5] C.I. Sukenik et al., Phys. Rev. Lett. 70, 560 (1993).
 [6] A. Landragin et al., Phys. Rev. Lett. 77, 1464 (1996).
 [7] F. Shimizu, Phys. Rev. Lett. 86, 987 (2001); H. Oberst et al., Phys. Rev. A 71, 052901 (2005).
 [8] Y.J. Lin et al., Phys. Rev. Lett. 92, 050404 (2004).
 [9] T.A. Pasquini et al., Phys. Rev. Lett. 93, 223201 (2004).
 [10] D.M. Harber, M. Obrecht, J. McGuirk, and E.A. Cornell, cond-mat/0506208.
 [11] I. Carusotto et al., Phys. Rev. Lett., in press.
 [12] H.B.G. Casimir and D. Polder, Phys. Rev. 73, 360 (1948).
 [13] I.E. Dzyaloshinskii, E.M. Lifshitz and L.P. Pitaevskii, Advances in Physics 38, 165 (1961).
 [14] P. Milonni, The Quantum Vacuum: An Introduction to Quantum Electrodynamics (Academic Press, Boston, 1994); K.A. Milton, J. Phys. A 37, R209 (2004).
 [15] E.M. Lifshitz, Doklady Akademii Nauk SSSR 100, 879 (1955).
 [16] S.M. Rytov, Y.A. Kravtsov and V.I. Tatarskii, Elements of random fields, Principles of Statistical Radiophysics (Springer, Berlin, 1989), enlarged edition of original Rytov's book published in Russian in 1953.
 [17] C. Henkel, K. Joulain, J.-P. Mulet, and J.-J. Greffet, J. Opt. A: Pure Appl. Opt. 4, S109 (2002).
 [18] A.E. Cohen and S. Mukamel, Phys. Rev. Lett. 91, 233202 (2003).
 [19] J.P. Gordon, and A. Ashkin, Phys. Rev. A 21, 1606 (1980).
 [20] J.M. Wyllie and J.E. Sipe, Phys. Rev. A 30, 1185 (1984); J.E. Sipe, J. Opt. Soc. Am. B 4, 481 (1987).
 [21] C. Henkel et al., Opt. Commun. 186, 57 (2000).
 [22] M. Antezza, to be published.
 [23] G. Roati et al., Phys. Rev. Lett. 92, 230402 (2004).
 [24] Y. Shin et al., Phys. Rev. Lett. 92, 050405 (2004).
 [25] The static limit, which is obtained by replacing $j''(!)$ with j''_0 , can be also used to simplify the evaluation of eq. (8). In the case of a sapphire substrate at ordinary temperatures ($j''_0 = 9.4$) this approximation is very accurate at

all distances.