

Supersolid Order from Disorder: Hard-Core Bosons on the Triangular Lattice

R. G. Melko,¹ A. Paramakanti,² A. A. Burkov,¹ A. Vishwanath,² D. N. Sheng,³ and L. Balents¹

¹Department of Physics, University of California, Santa Barbara, California 93106

²Department of Physics, University of California, Berkeley, California 94720

³Department of Physics and Astronomy, California State University, Northridge, California 91330

(Dated: April 14, 2024)

We study the interplay of Mott localization, geometric frustration, and superfluidity for hard-core bosons with nearest-neighbor repulsion on the triangular lattice. For this model at half-filling, we demonstrate that superfluidity survives for arbitrarily large repulsion, and that diagonal solid order emerges in the strongly correlated regime from an order-by-disorder mechanism. This is thus an unusual example of a stable supersolid phase of hard-core lattice bosons at a commensurate filling.

Introduction: Recent experiments on ^4He under pressure suggest [1] that a supersolid phase, in which long-range diagonal (crystalline) order and long-range off-diagonal (superfluid) order coexist, may arise at temperatures $T < 100 \text{ mK}$. Such a phase had been theoretically envisioned long ago [2], based on the proposition that a nonvanishing density of zero point defects in the solid (interstitials or vacancies) could Bose condense and form a superfluid "on top" of the existing crystalline background. However, early estimates [3] of the superfluid fraction in clean ^4He solid were very small ($\sim 10^{-4}$), and recent numerics with realistic interatomic potentials find no evidence for zero point defects or appreciable interparticle exchanges [4] in the crystal.

Since the microscopic conditions under which clean ^4He might exist in a supersolid phase are still unclear, it is useful to focus on simple lattice models of bosons in order to understand the different mechanisms by which a supersolid might emerge. Such "lattice supersolids" are superfluids which also break the lattice translation symmetries [5]. Numerical studies of interacting bosons on the square lattice find that stable supersolid phases form upon doping away from a half-filled checkerboard or striped solid [6, 7, 8]. These appear to be examples of the "defect condensation" mechanism, where doped bosons (holes) act as interstitial (vacancy) sites in the crystal.

In this letter we explore a different route to supersolidity based on the competition between Mott localization physics and geometric frustration. To illustrate this, we focus on a simple model of hard-core bosons at half-filling on the triangular lattice, interacting via a nearest-neighbor repulsive term,

$$H = \sum_{\langle ij \rangle} t \left(b_i^\dagger b_j + b_j^\dagger b_i \right) + \sum_{\langle ij \rangle} V \left(n_i - \frac{1}{2} \right) \left(n_j - \frac{1}{2} \right); \quad (1)$$

For this model, we show that a supersolid phase emerges for $V/t > 10$. The supersolid phase arises from an order-by-disorder effect in a strongly correlated superfluid. This route to supersolidity is thus complementary to the "defect condensation" mechanism which starts from a perfect crystal and considers a small defect-density induced supersolid. Our results are based on a combination of analytical arguments, projected boson wavefunctions,




FIG. 1: (color online). (a) Ground state superfluid density, calculated with exact diagonalization, quantum Monte Carlo, and variational wavefunction, with $J_z = 1$. The inset is ρ_s in the limit $J_z \rightarrow 1$, calculated by ED (for several system sizes N), and VWF (in the thermodynamic limit). (b) The diagonal order parameter squared. Examples of the QMC extrapolations to $N \rightarrow 1$ are illustrated in Fig. 2.

quantum Monte Carlo and exact diagonalization studies. We thus go well beyond the earlier spin-wave work [9] on this model. It would be interesting to look for signatures of this supersolid in experiments, e.g. on interacting cold bosonic atoms in an optical lattice [10], where both a zero momentum condensate peak and Bragg peaks at the ordering wavevectors would be expected in the usual time-of-flight momentum spectroscopy.

Alternatively, the Hamiltonian we study can be viewed, via the standard mapping from hardcore bosons to $S = 1/2$ spins, as an XXZ model:

$$H = \sum_{\langle ij \rangle} J_\parallel S_i^x S_j^x + S_i^y S_j^y + J_z S_i^z S_j^z; \quad (2)$$

with $J_z = V$ and $J_\parallel = 2t$. A spin Hamiltonian such as Eq. (2) may be realized in a Mott insulating phase of cold bosons on an optical lattice, if the bosons possess two internal states which can play the role of the spin

Strong coupling analysis for $J_z = J_z - 1$: We now turn to directly consider the strong-coupling limit. A simple qualitative picture (which we will reinforce with a rigorous analysis a little later) can be obtained by mapping the XXZ model onto a transverse field Ising model:

$$H = \sum_{\langle ij \rangle} J_z S_i^z S_j^z + \sum_i h_e S_i^x; \quad (7)$$

with $h_e = J_z \hbar S_i^x$. We have assumed here that, as suggested by the variational wavefunction analysis, superfluidity survives in the limit of small $J_z = J_z$. This transverse field Ising model has been studied earlier [11, 13] in various contexts, using Ginzburg-Landau theory and Monte Carlo numerics. It is well established that this model has diagonal long-range order, in particular of antiferromagnetic type, for small $h_e = J_z$. This analogy suggests that the supersolid behavior of the XXZ model should be viewed, similarly to Eq. (7), as arising microscopically from an order-by-disorder mechanism [14].

One might worry that the above analysis was based on a mean-field treatment of the superfluid component of the order, which may be unreliable at small $J_z = J_z$. We now turn to a more exact argument, which strongly argues for a ground state with broken spatial symmetries. In particular, we prove that for $J_z = J_z \neq 0$, the ground states are three-fold degenerate and transform non-trivially under lattice rotations.

In this limit ($J_z = 1$), the XXZ model reduces to

$$H_1 = \sum_{\langle ij \rangle} J_z \hat{P}_C (S_i^x S_j^x + S_i^y S_j^y) \hat{P}_C; \quad (8)$$

where $\hat{P}_C$ is a projection operator onto the classical Ising antiferromagnetic ground state manifold. Consider an $L \times L$ system with periodic boundary conditions, i.e. on the torus. We define the quantities $E_a = 0$, $a = 1; 2; 3$, as the number of frustrated bonds along a straight closed path (loop) through L sites along the a^{th} principal axis of the triangular lattice. These quantities are independent of translations of the loop. Moreover, it is straightforward to show that they commute with H_1 in Eq. (8) (and indeed any local Hamiltonian in this manifold). Because each triangle has one frustrated bond, one also finds $E_1 + E_2 + E_3 = L$. Under a six-fold rotation of the system, the E_a cyclically permute. Hence a unique ground state with $E_1 = E_2 = E_3$ that is also rotationally invariant can be found only for L a multiple of three. For $L \notin 0 \pmod{3}$, the ground state thus forms at least a three-fold degenerate rotational multiplet, as promised (it can be spanned by three states with angular momentum $\ell = 0; 1; 2 \pmod{3}$).

More refined arguments indicate that this degeneracy is split only by terms of order $O(e^{-cL})$ (with c independent of L) for $0 < J_z = J_z - 1$, and $L \neq 1$. Hence a non-trivial rotational multiplet remains the ground state with exponential accuracy in the thermodynamic limit. This behavior is inconsistent with a simple uniform superfluid,




FIG. 2: (color online). Finite-size scaling of QMC data for ρ_s (a) and $\ln \rho_s^2 = S(k_0) = L^2$ (b). Here, $J_z = 1$ and the inverse temperature is $\beta = 10$. Error bars (illustrated) are much smaller than the symbol sizes. Dotted lines show linear extrapolations for the very largest system sizes. The thick solid (blue) line is a fit to $\ln \rho_s^2 / L^2$, expected asymptotically when there is no diagonal LRO.

which is expected to have only states with excitation energies above the ground state scaling as a power-law of L . The most natural possibility consistent with this behavior is broken spatial symmetry of the ground state. Other arguments can be presented against more "exotic" possibilities. Having concluded that the system must break lattice symmetries, and already argued from our variational wavefunction that superfluidity persists for $J_z \neq 1$, we thus come to the conclusion that the ground state at large J_z is a supersolid, as our numerical simulations below prove.

Exact numerical results: To address the properties of the spin-1/2 XXZ model in Eq. (2) we have carried out exact diagonalization (ED) studies, as well as Stochastic Series Expansion (SSE) quantum Monte Carlo (QMC) simulations [16] in the grand canonical framework. Using straightforward modifications [15] to the original SSE directed-loop algorithm [17], the QMC is able to explore all regions of the finite-temperature phase diagram for small $J_z = J_z$. However for $J_z = J_z > 10$, the QMC is observed to experience a dynamical freezing due to the development of large energy barriers for local updates. In Fig. 1, we summarize our results for ED over the entire range of coupling strengths, and QMC in the weak to intermediate coupling regime (where we have been able to reliably equilibrate data).

The superfluid density is estimated by the stiffness, Eq. (6), easily measured using winding numbers in the QMC simulations [17, 18]. For a given $J_z = J_z$, ρ_s is extrapolated to $L \rightarrow \infty$ using finite-size scaling on systems of size $N = L \times L$ (Fig. 2a). The results show that for small $J_z = J_z$ the system is clearly in a robust superfluid

ground state with a large s . The stiffness value decreases with increasing $J_z=J_\perp$, however it remains finite to the largest values of $J_z=J_\perp$ studied with the QMC. The survival of s at large $J_z=J_\perp$ is in stark contrast to the behavior at $J_z=J_\perp = 1$, where the phase transition to a fully-polarized ferromagnetic phase is evident by the vanishing stiffness. Further, ED calculations (Fig. 1a, inset) strongly suggest that s survives to $J_z=J_\perp \rightarrow 1$, although system sizes are too small to allow a reliable extrapolation to $N \rightarrow 1$ for this data. The small value of s for $J_z \rightarrow 1$ indicates that the supersolid is very close in parameter space to an insulating state.

In the above analytical arguments, the survival of s at large $J_z=J_\perp$ is accompanied by the development of simultaneous Ising order (the supersolid phase). To address this prediction, we compute the static structure factor

$$S(k) = \frac{1}{N} \sum_{ij} e^{ik \cdot (r_i - r_j)} \langle S_i^z S_j^z \rangle \quad (9)$$

This structure factor reveals the development of sharp Bragg peaks for moderate values of $J_z=J_\perp$. The peaks occur at the corners of the hexagonal Brillouin zone ($k_0 = (4\pi/3, 0)$), in complete agreement with the above analytical discussion. The presence of diagonal long range order (LRO) can be demonstrated rigorously through a finite-size scaling analysis of $S(k_0)$ (Fig. 2b). Specifically, in a LRO state, the diagonal order parameter squared, $\langle S_i^z \rangle^2 = S(k_0) = L^2$, should scale to a constant as $L \rightarrow 1$ (with a leading order-correction varying as $1/L$ [20]). In contrast, in the absence of diagonal LRO, spin-spin correlations are short ranged, and $\langle S_i^z \rangle^2$ should tend to zero as $1/L^2$ in the asymptotic limit. Employing this analysis on the QMC data for all $J_z=J_\perp$ studied reveals the position of the zero-temperature phase transition from the superfluid to supersolid state to be between $J_z=J_\perp = 4$ and $J_z=J_\perp = 5$ (Fig. 1).

Finally, we note that the grand canonical QMC simulations indicate no significant deviations from half-filling in the supersolid phase for system sizes up to $L = 36$. In the ED, the minimum-energy ground state is also found to lie within the $\langle S_i^z \rangle = 0$ subspace. These observations suggest the occurrence of the antiferromagnetic rather than the ferromagnetic supersolid, however QMC simulations on much larger system sizes may be necessary in order to distinguish conclusively between the two.

Conclusions: We have presented a comprehensive study of the phase diagram of interacting hard-core bosons at half-filling (zero field XXZ model) on the triangular lattice. We have found that a subtle interplay between repulsion and quantum fluctuations produces a supersolid ground state in this system in the strongly correlated regime. This model provides an unusual example of a supersolid at a commensurate density arising from an order-by-disorder mechanism rather than defect condensation. This distinct mechanism completely avoids the instability of the defect-mediated square lattice supersolid to phase separation [8]. Here, the supersolid has

boson density equal to or extremely close to $1/2$, which is strongly favored by the maximal frustration-induced entropy of the corresponding classical Ising antiferromagnetic states.

The authors would like to thank A.W. Sandvik, D.J. Scalapino and D. Stamper-Kum for insightful discussions. Supercomputer time was provided by NCSA under grant number DMR020029N. Financial support was provided by National Science Foundation Grants DMR02-11166 (RGM), DMR-9985255 (LB, AAB), DMR-0307170 (DNS), the Packard Foundation (LB, AAB), ACS-PRF 41752-AC10 (DNS) and DOE LDRD DE-AC03-76SF00098 (AP and AV). During completion of this work we became aware of parallel numerical works [21], which reach similar conclusions.

-
- [1] E. Kim and M.H.W. Chan, *Nature* 427, 225 (2004); *Science* 305, 1941 (2004).
 - [2] A.F. Andreev and I.M. Lifshitz, *Sov. Phys. JETP* 29, 1107 (1969); G. Chester, *Phys. Rev. A* 2, 256 (1970).
 - [3] A.J. Leggett, *Phys. Rev. Lett.* 25, 1543 (1970).
 - [4] D.M. Ceperley and B. Bernu, *Phys. Rev. Lett.* 93, 155303 (2004).
 - [5] K.-S. Liu and M.E. Fisher, *J. Low Temp. Phys.* 10, 655 (1973).
 - [6] A. van Otterlo and K.H.W. Agenblast, *Phys. Rev. Lett.* 72, 3598 (1994); A. van Otterlo et al, *Phys. Rev. B* 52, 16176 (1995).
 - [7] G.G. Batrouni and R.T. Scalettar, *Phys. Rev. Lett.* 84, 1599 (2000); F. Hebert et al, *Phys. Rev. B* 65 014513 (2001).
 - [8] P. Sengupta, L.P. Pryadko, F. Alet, M. Troyer, and G. Schmit, *cond-mat/0412338* (unpublished).
 - [9] G. Murthy, D. Arovas and A. Auerbach, *Phys. Rev. B* 55, 3104 (1997).
 - [10] L.M. Duan, E. Demler, and M.D. Lukin, *Phys. Rev. Lett.* 91, 090402 (2003).
 - [11] D. Blankschtein, M. Ma, A.N. Berker, G.S. Grest, and C.M. Soukoulis, *Phys. Rev. B* 29, R5250 (1984).
 - [12] G.H. Wannier, *Phys. Rev.* 79, 357 (1950); R.M.F. Houtappel, *Physica* 16, 425 (1950).
 - [13] R.M. Oosner and S.L. Sondhi, *Phys. Rev. B* 63, 224401 (2001); R.M. Oosner, S.L. Sondhi, and P. Chandra, *Phys. Rev. B* 64, 144416 (2001).
 - [14] In contrast, generalizations of our analytical arguments predict a simple uniform superfluid on the kagome lattice at all $J_z=J_\perp$. This prediction has recently been confirmed by QMC calculations [15].
 - [15] S.V. Isakov and R.G. Melko (unpublished).
 - [16] A.W. Sandvik and J. Kurkijarvi, *Phys. Rev. B* 43, 5950 (1991); A.W. Sandvik, *J. Phys. A* 25, 3667 (1992).
 - [17] O.F. Syljäsén and A.W. Sandvik, *Phys. Rev. E* 66, 046701 (2002).
 - [18] E.L. Pollock and D.M. Ceperley, *Phys. Rev. B* 36, 8343 (1987).
 - [19] M. Boninsegni, *J. Low Temp. Phys.* 132, 39 (2003).
 - [20] D.A. Huse, *Phys. Rev. B* 37, R2380 (1988).
 - [21] D. Heidarian and K. Damle, *cond-mat/0505257*; S. Wessel and M. Troyer, *cond-mat/0505298*;