## A ccuracy of Sem iclassics: C om parative A nalysis of W K B and Instanton A pproaches.

V A Benderskii

Institute of Problem s of Chemical Physics, RAS 142432 Moscow Region, Chemogolovka, Russia and Laue-Langevin Institute, F-38042, Grenoble, France

E V Vetoshkin Institute of Problem s of Chemical Physics, RAS 142432 Moscow Region, Chemogolovka, Russia

E. I. Kats

Laue-Langevin Institute, F-38042, Grenoble, France and L.D.Landau Institute for Theoretical Physics, RAS, Moscow, Russia (D ated: A pril 14, 2024)

We analyze quantitatively the accuracy of eigenfunction and eigenvalue calculations in the fram e work of W KB and instanton sem iclassical methods. We show that to estimate the accuracy it is enough to compare two linearly independent (with the same quantum number) solutions to the Schrodinger equations with the potential under study, and with the approximating piecew ise sm ooth potential. The main advantage of the approach is related to the appropriate choice of the approximating potential, providing absolutely convergent majorant series for the solutions. We test our method for a strongly anharm onic one dimensional potential, but the basic ideas inspiring our work and its results can be applied to a large variety of interesting chemical and physical problem swhich are of relevance to various molecular system s.

PACS num bers: 05.45.-a, 72.10.-d

### I. IN TRODUCTION

It is a textbook wisdom that if the de Broglie wavelengths of particles are small in comparison with the characteristic space scales of a given problem, then the problem can be treated sem iclassically. The commonly used W K B m ethod (phase integral approach) [1] - [3] is intended for the conditions of "geom etrical optics", in which the gradient of the action is large, but slow ly variable (this is suggested also by it containing the factor h, since we are dealing with a sem iclassical approximation, in which h is taken as small). The corresponding condition can be form ulated m ore quantitatively [1] as follows: must vary only slightly over distances of the order of itself

$$\frac{d(x)=2}{dx} \quad 1; \tag{1}$$

where (x) = 2 h=p(x), and p(x) is a classical momentum. However, this simple criterion is not a practical tool to estimate how accurate could be found sem iclassical solutions of particular problems, since nothing is specied regarding the convergence of the sem iclassical series. The criterion (1) does not work to estimate the magnitude of the error involved in the approximate calculation of physical quantities (e.g., matrix elements), neither to not the domain of validity in the complex plane in which the sem iclassical solutions are dened. Indeed from (1) one can conclude only that higher order corrections to sem iclassical wave functions are small in the asymptotic regions, but this mathematical criterion has almost nothing to do with say the physical accuracy of sem iclassical matrix elements which depends on the wave function accuracy in space regions providing main contributions into the matrix elements under consideration. For example, the energy eigenvalues are determined by the asymptotical region of the linear turning points (i.e., the region distant from these points), and as well by the proximity region to the second order turning points, since in the both regions the wave functions possess the largest values.

W ithin the W KB m ethod such kind of a physical accuracy estimation has been performed long ago by N. and P.O. From an [4]. They analyzed higher order corrections to the sem iclassical wave functions and found that although those are really small over 1 = 2 ( 1 is sem iclassical parameter), the corrections are proportional to the factor  $[(E = ) U]^2$ , (where E is energy and U is potential), and thus the function has non-integrable singularity at the linear turning points where  $[(E = ) U]^2 = 0$  (or, within the alternative to W KB sem iclassical form alism so-called extrem e tunneling trajectory or instanton instanton approach [5] - [11], the corrections are singular in the second order turning points). To sum ount this problem in [4] (see also [13], and [14]) the analytical continuation of the correction function into the complex plane has been proposed, and it gives in practically bulky expressions even for simplem odel

potentials. Since this problem has relevance far beyond W KB treatment of a particular model potential, this is an issue of general interest to develop a simple and convenient in practice quantitative method to study the accuracy of the sem iclassical approach, and it is the immediate motivation of the present paper to develop a system atic procedure how to do it.

The idea of our approach is to construct two linearly independent continuous (with continuous rst derivatives) approximate solutions to the Schrödinger equation, which in the asymptotic region coincide with sem iclassic solutions, and in the vicinity of the turning points – with the exact solutions of the so-called comparison equation (i.e. the exact solution of the Schrödinger equation for the chosen appropriately approximate near the turning points potentials  $V_c$  (X), henceforth will be referred to as the comparison potential). A linearly independent currect criterion is form ulated as the majorant inequalities for a certain matrix (which we nd in the explicit analytical form and calculated numerically) connecting our approximate and exact solutions in the nite space interval (not only in the vicinity of the turning points). Since this method has largely gone unnoticed in the study of sem iclassics, we found it worthwhile to present its derivation in a short and explicit form, and also to point out its practical usability.

The remainder of this paper is organized as follows. In section II we present the basic expressions necessary for our investigation. In this section we also present the main steps and qualitative idea of our method. Section III contains our results. We derive the inequalities which enable us to nd the nite space interval (not at the isolated points) where the solutions have to be matched, and calculate the 2 2 coordinate dependent matrix connecting the approximate and exact solutions. Since the sem iclassical solutions of the harm onic potential coincide with the exact solutions, the accuracy of any sem iclassically treated problem depends crucially on its potential energy anharm onicity. That is why as the touchstone to test our method the results presented in the section III are applied to an anharm onic oscillator in section IV. We end with some brief conclusions in the same section.

### II. SEM ICLASSICAL EQUATIONS IN THE W KB AND INSTANTON FORMS

Technically the basic idea how to overcom e the di culty of the sem iclassical solutions in the vicinity of the turning points is reduced to an appropriate (adm itting exact analytic solutions) approximation of the potential near the turning points. A fler that step one has to match the asymptotics of this exact solution to the Schrodinger equation for an approximate potential with the sem iclassical solutions to the Schrodinger equation for the potential under consideration (i.e. approximate solutions of the exact potential) far from the turning points. To illustrate main ideas of any sem iclassic method (and to retain compactness and transparency of expressions) we discuss here a one dimensional case. As it is wellknown [1] in the W K B method solutions to the Schrodinger equation are sought in the form

$$= A \exp \frac{i}{h} ; \qquad (2)$$

where for the function called action the one particle Schrodinger equation (traditionally term ed as H am ilton Jacoby equation) reads as

$$\frac{1}{2m} \quad \frac{\theta}{\theta x}^2 = E \quad U; \qquad (3)$$

where m is a particle mass, E is its energy, and U is external eld potential. Since the system is supposed quasiclassical in its properties, we seek in the form of a series expanded in powers of h. Depending on normalization prefactor A (x) entering (2) can be also found but the corresponding equation (referred traditionally as transport equation) plays a pure passive role since it is fully determined by the action found as the solution of the H am ilton - Jacoby equation

$$\frac{\mathrm{i}h}{\mathrm{m}} = \frac{1}{2} \frac{\mathrm{e}^2}{\mathrm{e}x^2} \mathbf{A} + \frac{\mathrm{e}\mathbf{A}}{\mathrm{e}\mathbf{x}} \frac{\mathrm{e}}{\mathrm{e}\mathbf{x}} + \frac{\mathrm{h}^2}{2\mathrm{m}} \frac{\mathrm{e}^2\mathbf{A}}{\mathrm{e}\mathbf{x}^2} = 0; \qquad (4)$$

where in the spirit of the sem iclassical approximation the last term  $(/ h^2)$  is neglected. Technically of course more convenient to use instead of han expansion over equivalent but dimensionless parameter 1 1 we w ill call in what follows as sem iclassical parameter and de ne as

$$\frac{m_{0}a_{0}^{2}}{h}$$
 1; (5)

where  $a_0$  is a characteristic length of the problem, e.g. the tunneling distance,  $_0$  is a characteristic frequency, e.g. the oscillation frequency around the potential minimum. Evidently the sem iclassical parameter 1, and by its physical meaning it is determined by the ratio of the characteristic potential scale over the zero oscillation energy. We put h = 1, and use  $_0$  and  $a_0$  to set corresponding dimensionless scales, i.e. we introduce dimensionless energy  $E = _0$ , dimensionless coordinate X  $x=a_0$ , dimensionless potential V  $U = _0$  (except where explicitly stated

to the contrary and dimensions are necessary for understanding or numerical estimations). The analogous to (2), (3) procedure for the Schrödinger equation in the imaginary time (instanton formalism, corresponding to the W ick rotation in the phase space, when coordinates remain real valued x ! x but conjugated m omenta become imaginary  $p_x$  !  $ip_x$ ) can be formulated as the following substitution for the wave function (cf. to (2))

$$= A_{E} (X) \exp ( E_{E});$$
(6)

where the action  $_{\rm E}$  and we use the subscript E to denote so-called Euclidean action obtained from the WKB action

after the W ick rotation. Perform ed above rotation is not a harm less change of variables. The deep m eaning of this transform ation within the instanton approach is related to redistribution of di erent terms between the H am ilton – Jacoby and the transport equations. Indeed, like that is in the W K B m ethod, eigenvalues for the ground and for the low –lying states are of the order of <sup>0</sup>, while all other terms in (3) are of the order of <sup>1</sup>. Therefore to perform a regular expansion over <sup>1</sup> for the substitution (6) one has to remove the energy term from the H am ilton – Jacoby equation, and to include this term into the transport equation. Besides in the rst order over <sup>1</sup> one can neglect the term with the second derivative of the prefactor. As a result of this redistributions the both equations are presented as

$$\frac{1}{2} \frac{\Theta_{E}}{\Theta X}^{2} = V(X);$$
(7)

instead of the W KB Ham ilton - Jacoby equation (3), and the transport equation is

$$\frac{\partial A_E}{\partial X} \frac{\partial}{\partial X} + \frac{1}{2} \frac{\partial^2}{\partial X^2} A_E = A_E : \qquad (8)$$

O ne can easily note by a simple inspection of the W KB (3), (4) and of the instanton (7), (8) equations that although the both sem iclassicalm ethods can be form ulated neglecting terms of the order of <sup>1</sup>, therefore possessing the sam e accuracy over <sup>1</sup>, the solutions evidently coincide in the asymptotic classically forbidden region V (X) = , but their behavior, number and type of turning points (where any sem iclassic approximation does not work) are quite di erent. For example in the W KB form alism there are two turning points where V (X) (=) = 0 around each minimum of the potential, while in the instanton approach since the energy does not enter the H am ilton - Jacoby equation (thus one can say that no classically accessible regions at all) the turning points are extrem alpoints (minim a for the case) of the potential. Furtherm ore as a consequence of this di erence, in the W KB m ethod all turning points are linear, whereas in the instanton approach they are second order (quadratic over X).

### III. ACCURACY OF SEM ICLASSICAL APPROX IM AT ION

A rm ed with this know ledge we are in the position now to construct our approximants. Let us introduce besides the comparison potential  $V_c$  (X), one more specially chosen potential  $V_{sc}$  (henceforth will be referred to as the sem iclassical potential). This potential is chosen by the requirement that the exact solutions to the Schrodinger equation with  $V_{sc}$  coincide asymptotically with the sem iclassical solutions to the Schrodinger equation with the potential V (X) the problem under study. Thus according to the construction, the sem iclassical wave function sc satis es the equation

$$\int_{SC}^{1} \frac{d^{2}}{dX^{2}} = 2^{2} V_{SC} (X) - ;$$
(9)

and from here we can relate the sem iclassical potential ( $V_{sc}$  (X )) with the bare one (V (X ))

$$V_{sc}^{(1;2)} = V(X) - \frac{1}{2^2} A^{-1} - \frac{d^2 A}{dX^2}$$
; (10)

in the vicinity of the storder or of the second order (superscripts 1 or 2) turning points.

Since near the turning point X<sub>0</sub> the prefactors A<sup>(1)</sup> / K X<sub>0</sub> j<sup>1=4</sup>, and A<sup>(2)</sup> / (X X<sub>0</sub>)<sup>n</sup> (where n is an integer number which occurs from the transport equation (8) solution at the energy = n + (1=2)) the potential V<sub>sc</sub><sup>(1)</sup> at X ! X<sub>0</sub> is singular and negative, and V<sub>sc</sub><sup>(2)</sup> has the same singularity (/ (X X<sub>0</sub>)<sup>2</sup>) but positive. The di erence is due to the fact that near the W KB linear turning points we have deal with the V<sub>sc</sub><sup>(1)</sup> well, whereas near the second order instanton turning points one has to treat the potential barrier V<sub>sc</sub><sup>(2)</sup>. It m ight be useful to illustrate the essential features of the introduced above potentials V<sub>c</sub> and V<sub>sc</sub> applying the de nition (10) to a simple (but the generic touchstone) example of the follow ing anharm onic oscillator

$$V(X) = \frac{1}{2} [X^{2} + X^{3} + X^{4}]:$$
(11)

We show in Fig. 1 the sem iclassical and the comparison potentials associated with (11) for the W KB (Fig. 1a) and instanton (Fig. 1b) methods (= 1.25, = 0.5, and the energy window corresponds to n = 3 excited state of the potential (11)).

The key elements to construct our approximants are the following combinations related to probability ows to and from the turning points

$$J(X_{1}) = \int_{SC}^{1} \frac{d_{SC}}{dX} = \int_{C}^{1} \frac{d_{C}}{dX} = 2 \int_{X_{1}}^{2} \frac{d_{C}}{dX} = 2 \int_{X_{1}}^{2} \frac{d_{C}}{dX} = 2 \int_{SC}^{2} \frac{$$

where  $X_1 < X_0$  and analogously for  $X_2 > X_0$  the ow function J( $X_2$ ) is given by (12) where the integration limits are from  $X_2$  to +1. Since the exact wave functions are continuous with continuous rst derivatives (providing due to these features the continuity of the density probability currents), the idea of our procedure is to require the sam e from the approximate wave functions.

The integrals entering J (X<sub>1</sub>) and J (X<sub>2</sub>) can be calculated easily for any form of the potential, and the maximum accuracy of the any sem iclassical approach can be achieved upon the matching of the approximate solutions at the characteristic points X<sup>#</sup><sub>1,2</sub> where J (X<sup>#</sup><sub>1,2</sub>) = 0. The points X<sup>#</sup><sub>1,2</sub> do exist in the case when the potentials V<sub>c</sub> and V<sub>sc</sub> intersect in the region where the approximate wave functions sc and c are monotone ones. It is easy to realize (see e.g., Fig. 1) that the both points occur in the vicinity of the linear turning points for the potentials with  $d^2V=dX^2 > 0$ . One such a point disappears when the potential turning point becomes the in ection point, and there are no X<sup>#</sup><sub>1,2</sub> points at all for  $d^2V=dX^2 < 0$ . In the vicinity of the second order turning point the comparison potential V<sub>c</sub> is a parabolic one. The curvature of the latter potential corresponds to a certain renorm alization (/ <sup>2</sup>) of the characteristic oscillation frequency ( $d^2V=dX^2$ )<sub>X = X<sub>0</sub></sub>. Note in passing that the approach we are advocating here conceptually close (although not identical) to the scale transform ation proposed by M iller and G ood [15] and further developed in [16].

Thus the conditions  $J(X_{1/2}^{*}) = 0$  allow us to construct well controlled approximate solutions to the Schrödinger equation. The accuracy of the approximation depends on the deviation of the approximate wave functions from the exact ones in the vicinity of the characteristic points  $X_{1/2}^{*}$ . In own turn, the deviation is determined by the higher over  $(X_{0})$  terms of the potential V (X) which are not included in the harm onic comparison potential  $V_{c}$ . Include explicitly the corresponding higher order terms to distinguish  $V_{c}$  and  $V_{sc}$  potentials, we ind in the vicinity of the linear turning points

$$V_{sc}^{(1)} = V_c ' - \frac{c_1}{2} (X - X_0)^2 + \frac{!^2}{2} (X - X_0)^2;$$
 (13)

where the universal num erical constant  $c_1 = 5=32$ , and the second term in the rhs. is related to deviation of the bare potential from the linear one. The same manner near the second order turning points

 $V_{sc}^{(2)} = V_c ' - \frac{C_2}{2} (X - X_0)^2 + (X - X_0)^3;$  (14)

where the universal constant  $c_2 = n (n - 1)=4$  is zero for the lowest vibrational states n = 0;1, and the last term describes non-parabolicity of the potential. Note that unlike the sem iclassical action which within the instanton m ethod is independent of quantum numbers n, the position of the characteristic points does depend on n, and the  $X_{1,2}^{\#}$  points are placed near the boundaries of the classically accessible region.

Now we are in the position to construct the approximant wave functions

which are the solutions to the Schrodinger equation with the following piecewise smooth approximating potential

$$\nabla (X) = \frac{V_{c}(X); X_{1}^{\#} < X < X_{2}^{\#}}{V_{sc}; X < X_{1}^{\#}; X > X_{2}^{\#}} :$$
(16)

The wave functions calculated according to (15) in the fram ework of the instanton approach close to the W eber functions in the classically accessible regions, but their exponentially decaying tails in the classically forbidden regions correspond to the exact (bare) potential, not to its harm onic approximant. Analogously in the W KB m ethod these functions (15) coincide with the sem iclassical ones out of the interval ( $X_1^{\#}$ ;  $X_2^{\#}$ ), and with the A iry functions in this interval.

To proceed further on we have to relate our approximant wave functions (15) and two linearly independent solutions to the bare Schrödinger equation  $_1$  and  $_2$ . It can be written down formally as

$$(X) = {}^{\sim}(X) + {}^{Z_{X}}_{X_{0}} dX {}^{0}_{V} (X {}^{0}) G (X ; X {}^{0}) (X {}^{0}); \qquad (17)$$

where v = V(X)  $\forall (X), G(X; X^0)$  is the G reen function for the Schrödinger equation with the potential (16),

 $G(X;X_{1}) \quad const[_{1}(X)_{2}(X_{1}) \quad _{1}(X_{1})_{2}(X)];X_{1} \quad X; \qquad (18)$ 

where the constant in (18) is the W ronskian equal to  $(2)^{1}$  in the instanton m ethod, and  $i(2)^{1}$  within the W K B approach. In (17) (1; 2) (the same de nition for  $\gamma$ , and we take the turning point X<sub>0</sub>, where the functions and  $\sim$  are close to each other at the lower integration lim it.

The solution to the integral equation (17) is expressed as the Neumann series expansion,

$$(X) = {}^{\sim}(X) + {}^{Z_{X}}_{X_{0}} dX_{1} v (X_{1}) G (X; X_{1}) {}^{\sim}(X_{1}) + ;$$
(19)

and the m -th order term can be factorized and estim ated as

$$\frac{1}{m!} \int_{X_0}^{Z_X} dX_1 v(X_1) G(X;X_1)^{\sim}(X_1) :$$
(20)

The integrals entering this estim ation

$$Z_{X} = dX_{1}v(X_{1})G(X;X_{1})^{*}(X_{1}) = L_{12=22}(X)^{*}_{1}(X) + L_{11}=21(X)^{*}_{2}(X)$$
(21)

contain the 2 2 m atrix with the following matrix elements

$$L_{ij} = \int_{X_0}^{X_{X}} dX^{0} (X^{0}) v(X^{0}) (X^{0}) (X^$$

It is convenient to introduce the matrix  $\hat{C}^{(n)}$  relating the n-th order wave function correction (n) with the wave function  $\tilde{C}^{(n)}$ 

$$\begin{pmatrix} n \\ 1 \\ 2 \end{pmatrix}^{(n)} = \hat{C}^{(n)} \quad \tilde{1}^{1}_{2} \quad ;$$
 (23)

and the full connection matrix between the exact and approximate wave functions

$${}^{1}_{2} = \hat{C} \qquad {}^{-1}_{2} ;$$
 (24)

is

$$\hat{C} = {\stackrel{X}{}} \hat{C}^{(n)}$$
: (25)

A coording to the inequality (20)

$$\hat{C}^{(n)} = \frac{1}{n!} \hat{C}_{0}^{n};$$
 (26)

where

$$\hat{C}_{0} = \begin{array}{c} L_{12} & L_{11} \\ L_{22} & L_{12} \end{array} ; \qquad (27)$$

C on bining nally the expressions (25) – (27) we end up with the upper and lower bounds for the correction matrix  $\hat{C}$  estimation

$$1 + \hat{C}_0 \quad \hat{C} \quad \exp(\hat{C}_0):$$
 (28)

We conclude from (20), (22) that the Neum ann series posses an absolute convergence if all the matrix elements  $L_{ij}$  are nite. Besides, unlike the correction functions introduced within the From an approach [4], the integrals in (22) have no singularities on the realaxis. Evidently the integrals (22) are nite with the oscillating W K B functions, since the perturbation potential v (X) is not zero only in the close proximity to the characteristic points  $X_{1,2}^{\#}$ . However in the instanton method due to mixing of increasing and decreasing exponents, the matrix elements  $L_{22}$  is divergent. Despite of this divergency the product  $L_{22} \sim 1$ , we are only interested in, is nite, and it is convenient to perform one more transform ation to exclude explicitly this divergency.

Technically one can easily eliminate the both o -diagonal elements of the matrix  $\hat{C}_0$  and thus to get rid of the divergency of the (exponentially decreasing solution  $\tilde{c}_1$ ) amplitude due to the contribution to the  $\tilde{c}_1$  the exponentially increasing solution  $\tilde{c}_2$ . These linear transform ations renorm alize the correction matrix elements  $L_{22}$  and  $L_{11}$  as follows

$$L_{22}(X)^{-1}(X) = L_{22}(X)^{-2}(X); L_{11}(X)^{-2}(X) = L_{11}(X)^{-1}(X);$$
(29)

where the renorm alized m atrix elements  $\mathrm{L}_{22}$  and  $\mathrm{L}_{11}$  read as

$$L_{22}(X) = \frac{Z_{X}}{X_{0}} dX \frac{(-1)(X)(-2)(X)}{(-1)(X)(-2)(X)} - (X^{0}) V(X^{0}) - (X^{0}) U(X^{0}) - ($$

and

$$L_{11}(X) = \frac{Z_{X}}{X_{0}} dX \frac{(-1 (X^{0})^{2} (X^{0})^{2} (X^{0})^{2})}{(-1 (X^{0})^{2} (X^{0})^{2})} (X^{0}) V(X^{0}) (X^{0})^{2} (X^{0}) L_{12}(X) :$$
(31)

Now all the integrals entering  $L_{11}$ ,  $L_{22}$ , and  $L_{12}$  are convergent and nite at any X, the correction matrix  $\hat{C_0}$  is transformed into the diagonal and positively de ned matrix  $\hat{C_0}$ 

$$\hat{C}_{0} = \begin{array}{c} L_{11} & L_{12} & 0 \\ 0 & L_{12} & L_{22} \end{array}$$
(32)

The n-th order corrections in the instanton approach satisfy the inequality

$$0 \qquad {}^{(n)}_{1} \qquad (n!)^{1} (L_{11} \quad L_{12})^{\sim}_{1}; 0 \qquad {}^{(n)}_{2} \qquad (n!)^{1} (L_{12} \quad L_{22})^{\sim}_{2}: \qquad (33)$$

Explicit sum mation of rhs in (33) gives us the upper and the lower bound limits for the solutions of the initial Schrodinger equation, i.e. the stripe where increasing and decreasing solutions are con ned

$$j_{1}^{2}(X) j j_{1}(X) j j_{1}(X) j exp(L_{11} L_{12}); j_{2}^{2}(X) j j_{2}(X) j j_{2}(X) j exp(L_{12} L_{22}):$$
 (34)

It is ourm ain result in this paper, and the stripe (34) gives the accuracy of the sem iclassical instanton m ethod. B esides we are in the position now to estim ate the contribution of increasing sem iclassical solutions into decreasing ones (w hat is relevant to solve eigenvalue problem s). The sum m ation convergent m a prant series enables us to estim ate the upper bound for this contribution

$$\frac{L_{11}}{L_{22}} (1 \exp(L_{12})):$$
(35)

Therefore at  $L_{12} = 1$  the summation of all order perturbation terms enhances the 1-st order correction by the factor  $L_{12}=L_{22}$ . Analogously the majorant estimates described above can be used to construct the connection matrices linking the sem iclassical solutions through the turning points. The comparison of the bare connection matrices (see e.g., [3], [9], [11]) with the matrices calculated accordingly to (22) - (30) provides the estimates for the eigenvalue accuracy.

As it was mentioned already, for the W KB method the procedure is even more simple, since no any divergency and therefore no need to perform the transformation (29). The similar to (20) - (21) factorization gives the  $\hat{C}_0$  matrix (cf. with (27) for the instanton approach)

$$\hat{C}_{0} = i \begin{array}{ccc} L_{12} & L_{11} \\ L_{22} & L_{12} \end{array} ;$$
(36)

and the estimations for the n-th order contribution (cf. with (33) can be form ulated now as

$$f^{(2n)} j = \frac{1}{(2n)!} = 0^{n};$$
 (37)

and

$$\mathbf{j}\hat{\mathbf{C}}^{(2n+1)}\mathbf{j} = \frac{1}{(2n+1)!} \, {}^{n}\hat{\mathbf{C}}_{0} \, ; \tag{38}$$

where we denote  $L_{12}^2$   $L_{11}L_{22}$ . Correspondingly to (37), (38) the diagonal and o -diagonal correction m atrix elements are bounded from above

$$p_{11=22} j \cos^{-1} + i L_{12} \frac{p_{--}}{p_{--}};$$
 (39)

and

$$p_{12=21}$$
 j  $L_{11} \frac{p_{--}}{p_{--}}$  : (40)

The whole procedure we employed is rationalized in the Fig. 2, where we compare the solutions to the comparison equation with the anharm onic oscillator sem iclassical wave functions computed within the instanton and W KB approaches and indicate the optim alm atching points  $X^{\#}$  found accordingly to the condition J ( $X^{\#}$ ) = 0.

## IV. ANHARMONIC OSCILLATOR

In closing let us illustrate how our estimations (34), (38) work for a strongly anham onic potential (11). A lthough it is not great trium photo re-derive the known results, our derivation illustrates several characteristic features of the correction matrix techniques derived in the section III: better accuracy, rapid convergence, simple disposal of divergences, and ease of computation in particular. The main message of our consideration in the precedent section III is that the quantitative accuracy of the sem iclassics depends crucially on the proximity of the sem iclassical wave functions to the solutions of the comparison equation in the region of the asymptotically smooth matching. Therefore, it is tempting to improve the accuracy by taking into account the anham onic corrections to the comparison potential  $V_c(X)$ . However, since the eigenvalues and the norm alization of the wave functions are alm ost independent of the detailed behavior in the vicinity of the linear turning points (because near these points, situated at the boundaries of the classically accessible region, the probability density (i.e. j j<sup>2</sup>) is exponentially sm all) this idea is useless for the W K B approach. In contrast with this, for the instanton method, the accuracy can be improved considerably upon including the anham onic corrections into the comparison potential. Indeed, within the instanton method the accuracy is determined by the vicinity of the second order turning points where the wave functions acquire the largest values (and just in this region the smooth matching described above has to be performed).

Let us rem ind rst the traditional (but form ulated within the sem iclassical fram ework) perturbation theory. K eeping in the transport equation (4) the second derivative of the prefactor A, the system of the equations of H am ilton -Jacoby (3) and the exact transport equation

$$\frac{d^2 A}{dX^2} = 2 \quad \frac{dA}{dX} \frac{d}{dX} + \frac{d^2}{dX^2} \qquad A = 0$$
(41)

are exactly equivalent to the Schrodinger equation under consideration. For the second order turning points the anharm onic corrections

$$V_{\rm p} = \frac{1}{2} X^{3} + X^{4}$$
(42)

can be considered as a perturbation and it is convenient to include this perturbation  $V_p$  into the transport equation (41). Then the comparison equation is reduced to the inhomogeneous W eber equation, and its solutions can be expanded over the W eber functions D (X) [20] (see also [13], [14])

$$(X) = N \quad D \quad (X) + \begin{array}{c} X \\ b_k \quad D \\ k \end{array}$$
 (43)

ı.

where N<sup>2</sup> = 1+  $\stackrel{P}{=} \stackrel{b^2}{\stackrel{h}{=}} _{,and} \stackrel{b^2}{=} _{,and$ 

$$k = 1; 3; and k = 0; 2; 4$$
 (44)

for the cubic and fourth order anham onic corrections respectively. Explicitly these non-zero expansion coe cients can be found by straitforward calculations, and they are

$$b_{0} = \frac{3}{2} + \frac{1}{2}; b_{1} = 3^{2}; b_{1} = (+1); b_{2} - \frac{1}{2}(-1) + \frac{1}{2}; b_{2} - \frac{1}{2} + \frac{3}{2}; (45)$$

$$b_{3} = \frac{1}{3}(-1)(-2)(-3); b_{2} - \frac{1}{3}; b_{4} = \frac{1}{4}(-1)(-2)(-3)(-4); b_{3} - \frac{1}{4}:$$

On equal footing we can different nd the perturbative corrections to the Bohr-Sommerfeld quantization rules, and therefore the eigenvalues. The calculation is straightforward, though deserves some precaution and rather tedious. Skipping a large am ount of tedious algebra we end up with the fractional part of the quantum number

and up to the second order over the anharm onic perturbation  $V_{\rm p}$  we  $\$ nd

$${n \choose n} = \frac{15^2}{2} n^2 + n + \frac{11}{30} + \frac{3}{2} n^2 + n + \frac{1}{2}$$
 (47)

However the described standard perturbative approach leads to qualitatively wrong features of the solutions. For example, the wave functions (43), (45) are represented as a product of independent exponential factors and dependent of polynom ials. As it is well known [1] in one dimension the n-th excited state wave function must have n zeros (and the number of zeros may not be changed by any perturbation). However in the m-th order perturbation theory approximation, the wave function (43) corresponding to a certain excited state n contains H emmittian polynom ials up to the order n + 3m or n + 4m for the cubic or quartic anharm onic perturbations, respectively. Therefore some false zeros of the wave function appears in the standard perturbation theory, and the region where the function oscillates become s m ore and m ore wide in the higher order over perturbations approximation. The contributions of these qualitatively and quantitatively incorrect higher order terms become dominating in the asymptotic region at j j 1,

1. It conforms with the classical results due to Bender and W u [17], [18] who have shown that for the quartic anham onic potential (= 0, > 0 in (11)) the convergency radius is zero. Moreover, for = 0 and for an arbitrary sm all (11) is the cubic anham onic potential, i.e., the decay one. Thus it should have only complex eigenvalues, what is not the case for the eigenvalues calculated within the perturbation theory. The method we developed in section III enables us not only to estimate more accurate the anham onic corrections to the eigenvalues, and to bring the whole schema of the calculations in a more elegant form. Our noting of the correction matrices is not merely to surpass a technical di culty of the standard perturbative method, it is more one of principle, and we will show that the method has no draw backs of the perturbation theory.

The proof proceeds as follows. Let us consider instanton method for the anhamonic potential (41) possessing one second order turning point X = 0. As it was shown in the section III one has to indialso two other characteristic points which are the roots of the equation (12). We denote the points as  $X_{L}^{\#}$ , and  $X_{R}^{\#}$  (to refer by the self-explanatory subscripts L and R to the left and to the right from the turning point X = 0). At the next step using the correction matrices introduced in the section III, we can de ne form ally the transform ation of our approximate wave functions (15) into the unknown exact wave functions as

$${}^{\mathrm{L}}_{\mathrm{R}} = \widehat{\mathrm{C}} \quad {}^{\sim}_{\mathrm{R}} ; \qquad (48)$$

where as above the subscripts L and R refer to the wave functions in the regions to the left and to the right from the turning point X = 0. We do not know the correction matrix  $\hat{C}$  but we do know (see (23) - (31)) the boundary estimations for the matrix.

The Eq. (48) can be used also to correct the known at the second order turning point the connection matrix  $\hat{M}$  [3], [11]. Indeed the connection matrices link the sem iclassical solutions in the X-regions to the left and to the right from the turning points. For the isolated second order turning point (we are dealing within the instanton method), the connection matrix  $\hat{M}$  links the exponentially increasing and decreasing solutions in the space regions separated by the turning point. The condition ensuring the correct asymptotic behavior is the quantization rule for this case which can be form ulated as  $M_{11} = 0$  ( $M_{1j}$  are the matrix elements of the connection matrix  $\hat{M}$ ). Since in the regions to the left and to the right from the turning point our approximate solutions  $\tilde{C}$  coincide by their de nition (15) with the sem iclassical ones, the correction matrix method enables us to correct the quantization rule too. Namely, the quantization rules are form ulated within the connection matrix technique read now as

$$C_{22}^{R}T_{2}C_{22}^{L} = C_{21}^{R}C_{21}^{L}\frac{\sin^{2}()}{T_{2}} + (C_{22}^{L}C_{21}^{R} + C_{21}^{L}C_{22}^{R})\cos() = 0; \qquad (49)$$

where the Stokes constant for the second order turning point [3] is

$$T_2 = -\frac{P_2}{(1-1)};$$
 (50)

and  $C_{ij}^{R,L}$  are the correction matrices at the  $X_R^{\#}$  or  $X_L^{\#}$  characteristic points respectively. Expanding the G am ma function entering (49) around the integer numbers, i.e., as above (46), = n + n one can nd from the equation (49) the fractional part of the quantum number. If we were known the correction matrix  $\hat{C}$  the solution of (49) would provide the exact eigenvalues. But we do know only the estimations from below and from above for the  $\hat{C}$  matrix. In the same spirit we can calculate the estimations for the correction matrices (and therefore for the eigenvalues) within the W K B approach.

Luckily it turns out that the mathematical nature of the sem iclassical problem is on our side here, and, in fact, even the rst order estimation from below  $\hat{C}^{(0)}$  (23) gives already the accuracy comparable with the standard perturbation procedure, and the estimation from above (25) gives the eigenvalues almost indistinguishable from the "exact" ones obtained by the numerical diagonalization of the Ham iltonian. The same true for the wave functions found by the correction matrix technique. We show in Fig. 3 j  $_3$  for the same anham onic potential (11) with = 1.25, = 0.5. C learly the exact numerical results and those obtained by our correction matrix techniques are correct qualitatively and in the very good quantitative agreement (indistinguishable starting from the second order approximation) unlike the situation with the standard perturbation theory. Besides we present in the table the eigenvalues of the anham onic potential. We take the anham onic coe cients and in (11) so large, that corresponding perturbations of the eigen values are of the order of the bare harm onic frequency (one in our dimensionless units = 1.2; = 0.5). In the table the eigenvalues found by the numerical diagonalization are presented in the colum n I. The colum n II contains the harm onic approximation results, the colum n III is the second order perturbation theory (47), and the colum ns IV and V results are obtained by applying our correction matrix technique: estimation from below with the rst order correction matrix (23) in the colum n IV, and the estimation from above with the matrix (25) in the colum n V.

To conclude, as we have shown how to estimate the corrections to the main technical tool for the sem iclassical approach, the connection matrices linking the solutions to the left and to the right from the turning points. Everything (e.g., the upper and the lower bounds for  $_n$ ) is determined by the matrices  $L_{12}$  and  $L_{22}$ . Approximating to (11) potential is found from (13) and after that straightforward computing according to (22) the matrices  $L_{1j}$  and their renorm alization (30) leads to the corrections we are looking for, presented in the Fig. 4, which allow us to estimate the accuracy of the sem iclassical eigenstates and eigenfunctions. We conclude that even for a strongly anharm onic potential the both m ethods (W K B and instanton) are fairly accurate ones (about 5%) up to the energy close to the potential

barrier top (in the region of negative curvature, we already discussed above). It is worth noting that in the fram e work of the conventional perturbation theory (due to zero convergency radius with respect to coe cient in (11)) pure com putational problems to get the sam e accuracy become nearly unsum ountable, see e.g., [17], [19]. Our notings show that to estim ate quantitatively the sem iclassical accuracy it is enough to com pare two linearly independent (with the sam e quantum number) solutions of the initial potential under study, and of the approximating piecew is an ooth potential. The main advantage of the approach is related to the appropriate (13), (16) choice of the approximating potential, providing absolutely convergent majorant series (17) for the solutions. A ctually our correction matrix technique is a fairly universal one and enables to estimate (and im prove!) the sem iclassical accuracy for arbitrary one dimensional potentials with any combination of the turning and of the crossing points.

#### A cknow ledgm ents

The research described in this publication was made possible in part by RFFR Grants. One of us (E K.) is indebted to INTAS Grant (under No. 01-0105) for partial support, and V B. and E.V. are thankful to CRDF Grant RU-C1-2575-M O-04.

- [1] L D Landau, E M Lifshits, Quantum M echanics (non-relativistic theory), Pergam on Press, New York, 1965.
- [2] R P Feynm an, A R Hibbs, Quantum Mechanics and Path Integrals, M cG raw Hill Book Company, New York, 1965.
- [3] J.H. eading, An Introduction to Phase-Integral M ethods, W iley Interscience, London, 1962.
- [4] N From an, P.O. From an, JW KB Approximation, North-Holland Publishing Company, Amsterdam, 1965.
- [5] A M Polyakov, NuclPhys. B, 129, 429 (1977).
- [6] S.C. olem an, A spects of Sym m etry, C am bridge U niversity P ress, C am bridge, 1985.
- [7] A Schm id, Ann. Phys., 170, 333 (1986).
- [8] V A Benderskii, D E M akarov, P G G rinevich, Chem . Phys., 170, 275 (1993).
- [9] V A Benderskii, D E M akarov, C A W ight, Chem ical Dynam ics at Low Temperatures, W illey-Interscience, New York, 1994.
- [10] V A Benderskii, E.V. Vetoshkin, Chem. Phys., 257, 203 (2000).
- [11] V A Benderskii, E.V. Vetoshkin, E.IKats, JETP, 95, 645 (2002).
- [12] R E Langer, Phys. Rev., 51, 669 (1937).
- [13] F W JO lver, J.Res.Nat.Bur.Stand., 63 B, 131 (1959).
- [14] F W J.O lver, A symptotics and Special Functions, A cad. Press., New York, 1974.
- [15] S.C.M iller, R.H.Good, Phys. Rev., 91, 174 (1954).
- [16] P.Pechukas, J.Chem. Phys., 54, 3864 (1971).
- [17] C M Bender, T S.W u, Phys. Rev., 184, 1231 (1969).
- [18] C M Bender, T S.W u, Phys. Rev. D, 1, 1620 (1973).
- [19] A.V. Turbiner, Sov. Phys. U sp., 27, 668 (1984).
- [20] A Erdelyi, W Magnus, F O berhettinger, F G. Tricom i, Higher Transcendental Functions, voll vol.3, M cG raw Hill, N ew York (1953).
- [21] V A Benderskii, E. V. Vetoshkin, E. IK ats, Phys. Rev. A, 69, 062508 (2004).

# Figure Caption

Fig.1

The characteristic sem iclassical potentials  $V_{sc}$  (dot-dashed lines) and  $V_c$  (dashed lines) for the bare anharm onic potential (11): = 1.25, = 0.5 ( = 33, and the energy window corresponds to n = 3):

(a) instanton approach;

(b)WKBmethod.

Fig.2

Sem iclassical wave function  $_3$  (n = 3) for the anharm onic potential (11) with = 1.25, = 0.5 ( = 33). Stars indicate the matching points, dashed lines show the solutions to the comparison equations, and:

(a) - solid line traces the instanton solution;

(b) – solid line shows the W KB wave function.

Fig.3

C om parison of the exact j  $_3 f$  (solid line) for the anham onic potential (11) with = 1.25, = 0.5 ( = 33) with two low est (zero and rst order) approximations of the correction matrix method (dashed and dot-dashed lines). Note that the second order approximation with the relative accuracy 10<sup>2</sup> is indistinguishable from the exact numerical results.

Fig.4

Corrections to the decreasing solutions in for the anham onic potential (11);

= 0:5; n = 3; = 33;

(1, 3) -instanton m ethod, (2, 4) -W KB, (1, 2) -the rst order corrections L<sub>11</sub>, (3, 4) -the upper bound estimation sum m ing up all terms. The vertical dashed lines indicate the values of the cubic anharmonic term () where the in ection point and new extrem a of the potential are appeared.

Table

E igenvalues of the anham onic potential (11) (= 12; = 0.5).

I the eigenvalues found by the num erical diagonalization;

II the harm on ic oscillator eigenvalues;

III the eigenvalues in the second order perturbation theory (47);

IV the eigenvalues estim ated from below by the correction m atrix (23);

V the estimation from above with the matrix (25).




Fig. 1




Fig. 2




Fig. 3




Fig. 4

n =	Ι	Π	III	IV	V
0	0.0149	0.0152	0.0149	0.0149	0.0149
1	0.0433	0.0455	0.0434	0.0433	0.0433
2	0.0697	0.0758	0.0701	0.0699	0.0695
3	0.0942	0.1061	0.0950	0.0946	0.0939
4	0.1168	0.1364	0.1182	0.1178	0.1162
5	0.1382	0.1667	0.1395	0.1396	0.1373