

Multipole ordering in f-electron systems

Katsunori Kubo, Takashi Hotta

Advanced Science Research Center, Japan Atomic Energy Research Institute, Tokai, Ibaraki 319-1195, Japan

Abstract

In order to investigate multipole ordering in f-electron systems from a microscopic viewpoint, we study the so-called δ models on three kinds of lattices, simple cubic (sc), bcc, and fcc, based on a j-j coupling scheme with f-electron hopping integrals through (ff) bonding. From the δ model, we derive an effective model for each lattice structure by using the second-order perturbation theory with respect to (ff). By further applying mean-field theory to the effective model, we find a ${}_{3g}$ antiferro-quadrupole transition for the sc lattice, a ${}_{2u}$ antiferro-octupole transition for the bcc lattice, and a longitudinal triple-q ${}_{5u}$ octupole transition for the fcc lattice.

Key words: multipole ordering, j-j coupling scheme, δ crystalline electric field ground state
PACS: 75.30.Et; 71.10.Fd; 75.40.Cx

In recent decades, various kinds of magnetic and orbital ordering have been found in f-electron systems. In particular, it has been recognized that cubic systems with δ crystalline electric field ground states frequently exhibit higher-order multipole ordering due to their high symmetry. Indeed, octupole ordering has been proposed to reconcile experimental observations for $\text{Ce}_x\text{La}_{1-x}\text{B}_6$ [1,2,3,4,5] and NpO_2 [6,7,8,9]. To understand the origin of such multipole ordering from a unified viewpoint, it is important to analyze a simple microscopic model with correct f-electron symmetry.

In this paper, we study the so-called δ models on three kinds of lattices, simple cubic (sc), bcc, and fcc, based on a j-j coupling scheme. For the description of the model, we define annihilation operators in the second-quantized form for δ electrons with δ orbitals as $f_{r\alpha} = \frac{1}{\sqrt{5}}a_{r5=2} + \frac{1}{\sqrt{15}}a_{r3=2}$, $f_{r\beta} = \frac{1}{\sqrt{5}}a_{r5=2} + \frac{1}{\sqrt{15}}a_{r3=2}$, $f_{r\gamma} = a_{r1=2}$, and $f_{r\delta} = a_{r1=2}$, where a_{rj_z} is the annihilation operator for an electron with the z-component j_z of the total angular momentum $j=5/2$ at site r .

In the tight-binding approximation, the model Hamiltonian is given by [10]

$$H = \sum_{r,r'} t_{r,r'} f_{r\alpha}^\dagger f_{r'\alpha} + U \sum_r n_{r\alpha} n_{r\beta} + U^0 \sum_r n_{r\alpha} n_{r\beta} + J \sum_{r,r'} f_{r\alpha}^\dagger f_{r'\alpha}^\dagger f_{r'\beta} f_{r\beta} + J^0 \sum_{r,r'} f_{r\alpha}^\dagger f_{r'\alpha}^\dagger f_{r'\beta} f_{r\beta} \quad (1)$$

where \mathbf{r} is a vector connecting nearest-neighbor sites, $t_{r,r'}$ is the hopping integral of an electron with $(\delta; \delta)$ at site r to the $(\delta; \delta)$ state at r' through (ff) bonding [11], $n_{r\alpha} = f_{r\alpha}^\dagger f_{r\alpha}$, and $n_{r\beta} = f_{r\beta}^\dagger f_{r\beta}$. The coupling constants U , U^0 , J , and J^0 denote the intra-orbital, inter-orbital, exchange, and pair-hopping interactions, respectively. Note that the form of $t_{r,r'}$ characterizes the lattice structure.

By using the second-order perturbation theory with respect to (ff) including only the lowest energy triplet among the intermediate states, we obtain effective multipole interactions for each lattice structure. The detail of the effective models will be reported elsewhere, and here we report the ordered state obtained in the mean-field theory.

For sc lattice, a ${}_{3g}$ antiferro quadrupole (AFQ) transition occurs at a finite temperature, and as lowering temperature further, we find another transition to

* Corresponding author.




Fig. 1. Ordered states in the sc lattice under a magnetic field along [001]. (a) The $3g$ AFQ state. (b) The FM state with the $3g$ AFQ moment.




Fig. 2. Ordered states in the bcc lattice under a magnetic field along [001]. (a) The $2u$ AFO state. (b) The FM state with the $2u$ AFO moment.

a ferromagnetic (FM) state. Figures 1(a) and (b) show schematic views for the ordered states. In these figures, we depict the surface defined by $r = \sqrt{\sum_j (r_j(\theta; \phi))^2}$ in the polar coordinates, when the $5f$ wave-function is decomposed into radial and angular parts as $(r; \theta; \phi) = R(r) Y(\theta; \phi)$, where Y denotes real spin. Note that white-shift in the surface indicates the increase of the weight of up-spin state. Figure 1(a) shows the AFQ state, in which the spin distribution is isotropic and the weights of up- and down-spin states are the same. In the FM state state as shown in Fig. 1(b), the charge distribution is the same as in Fig. 1(a), while the weight of up-spin state is larger than that of down-spin. Note that the spin distribution is not isotropic due to the spin-orbit interaction.

For the bcc lattice, a $2u$ antiferro octupole (AFO) ordering occurs first, and then, a FM phase transition follows it. Figure 2(a) shows the $2u$ AFO state. Since this state does not accompany any other moment, the charge distribution retains cubic symmetry and the weights of up- and down-spin states are the same in total. However, the spin distribution is anisotropic, and this state has finite octupole moment. In the FM state as shown in Fig. 2(b), the weight of up-spin state increases, and charge distribution is also changed since this state has the $5g$ quadrupole moment.

For the fcc lattice, we have analyzed the effective model in a previous paper [12], in which we have evaluated multipole correlation functions by applying numerical diagonalization to the effective model on a small cluster. Then, we have determined that relevant




Fig. 3. The triple- q $5u$ octupole state in the fcc lattice.

interactions to the ground state are those between $4u$ and $5u$ moments. We have further applied mean-field theory to the simplified model including only these relevant interactions, and reached the conclusion that the ground state is the longitudinal triple- q $5u$ octupole state, which has been proposed for NpO_2 phenomenologically [8,9]. Figure 3 shows the $5u$ octupole state. This state accompany triple- q $5g$ quadrupole moment. Note that this state does not have frustration even in the fcc lattice.

In summary, we have derived the multipole interaction model from the microscopic 8 Hamiltonian. By analyzing the effective model, we find a $3g$ AFQ state for the sc lattice, a $2u$ AFO state for the bcc lattice, and the longitudinal triple- q $5u$ octupole state for the fcc lattice.

We thank S. Kambe, N. Metoki, H. Onishi, Y. Tokunaga, K. Ueda, R. E. Walstedt, and H. Yasuoka for discussions. K. K. is supported by the REIMEI Research Resources of Japan Atomic Energy Research Institute. T. H. is supported from Japan Society for the Promotion of Science and from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

References

- [1] Y. Kuramoto, H. Kusunose, J. Phys. Soc. Jpn. 69 (2000) 671.
- [2] H. Kusunose, Y. Kuramoto, J. Phys. Soc. Jpn. 70 (2001) 1751.
- [3] T. Sakakibara, et al., J. Phys. Chem. Solids 63 (2002) 1147; T. Sakakibara, et al., Physica B 312-313 (2002) 194.
- [4] K. Kubo, Y. Kuramoto, J. Phys. Soc. Jpn. 72 (2003) 1859.
- [5] K. Kubo, Y. Kuramoto, J. Phys. Soc. Jpn. 73 (2004) 216.
- [6] P. Santini, G. Amoretti, Phys. Rev. Lett. 85 (2000) 2188.
- [7] P. Santini, G. Amoretti, J. Phys. Soc. Jpn. 71 (Suppl.) (2002) 11.
- [8] J. A. Paixao, et al., Phys. Rev. Lett. 89 (2002) 187202.
- [9] R. Caciuro, et al., J. Phys.: Condens. Matter 15 (2003) S2287.
- [10] T. Hotta, K. Ueda, Phys. Rev. B 67 (2003) 104518.

[1] K . Takegahara, et al., J. Phys. C : Solid St. Phys. 13 (1980) 583.

[12] K . Kubo, T . Hotta, Phys. Rev. B 71 (2005) 140404(R) .