A large m agnetic storage ring for Bose-E instein condensates

A.S.Amold, C.S.Garvie, and E.Riis

Dept. of Physics, University of Strathclyde, Glasgow G4 ONG, UK

www.photonics.phys.strath.ac.uk

(D ated: February 1, 2022)

C old atom ic clouds and B ose-E instein condensates have been stored in a 10 cm diam eter verticallyoriented m agnetic ring. An azim uthal m agnetic eld enables low -loss propagation of atom ic clouds over a total distance of 2 m, with a heating rate of less than 50 nK /s. The vertical geom etry was used to split an atom ic cloud into two counter-rotating clouds which were recomb ined after one revolution. The system will be ideal for studying condensate collisions and ultim ately Sagnac interferom etry.

The eld of atom optics [1] has seen a plethora of signi cant advances since the advent of laser cooling [2] and gaseous B ose-E instein condensation (BEC) [3]. A nalogs of m irrors, lenses, and beam splitters for m anipulating ultra-cold atom s now exist and are continually being im – proved.

A relatively recent addition to the atom -optical toolbox is the cold atom storage ring, where atom s are guided around a closed path. This is an interesting con guration for performing, for instance, ultra-sensitive Sagnac [4] atom interferom etry. An electrostatic storage ring was rst reported form olecules [5] in 2001. However, for many interferom etry experiments atom s are more suitable candidates, as they can be easily laser cooled and/or prepared in the same quantum state.

The rst atom ic storage ring was formed with the magnetic quadrupole eld created by two concentric current carrying bops [6]. Our storage ring [7] utilises the more symmetric quadrupole eld of a four-bop geometry (Fig. 1). A third group has recently developed a storage Stadium ' [8] based on a magnetic waveguide. Our ring containsmore than 5 10° atom s, has a lifetim e of 50 s and an area of 72 cm²; each corresponding to m ore than an order of m agnitude in provem ent with respect to R efs. [6, 8]. The Sagnac e ect [4] is linearly proportional to the area of an interferom eter, and our ring's area com pares favourably with that of a state-of-the-art therm al beam atom interferom eter gyro (A = 0.22 cm^2 [9]). The atom s can com plete multiple revolutions in the ring, further increasing our e ective area and thus the rotation sensitivity. In addition, our storage ring has the unique feature that we are able to form a BEC in a section of the ring [7], and observe its propagation around the ring.

W e begin by considering the storage ring theoretically, before discussing our experim ental setup. C om parisons will then be drawn between our experim ent and theory.

All magnetic atom -optical elements make use of the Stem-Gerlach potential U = $_B g_F m_F B$ experienced by an atom moving adiabatically in a magnetic eld of magnitude B, where $_B$ is the Bohr magneton, m_F the atom 's hyper ne magnetic quantum number, and g_F is the Lande g-factor. To make a storage ring one must use atom s in weak-eld-seeking magnetic states ($g_F m_F > 0$); which are attracted to minim a of the magnetic eld magnetic eld magnetic eld magnetic states (states eld magnetic states descented to minime a state state).

nitude. U sing the B iot-Savart law, one can express the cylindrically symmetric magnetic eld from a single coil of radius R; b_R (r;z); in terms of elliptic integrals [10]. O ur storage ring comprises four concentric circular coils (Fig. 1), with a toroidal quadrupole total magnetic eld:

$$B (r;z) = b_{R_{R}} (r;z_{z}) b_{R_{R}} (r;z+_{z})$$
$$b_{R+_{R}} (r;z_{z}) + b_{R+_{R}} (r;z+_{z}): (1)$$

In our experiment R = 5.0 cm; $_R = 1.25 \text{ cm}$; and $_z = 1.35 \text{ cm}$; leading to a ring of zero magnetic eld at a radius $R_0 = 4.3 \text{ cm}$ slightly smaller than the mean coil radius R: The axialw ire (Fig. 1) adds a 1=r azim uthalm agnetic eld to Eq. 1, which has little e ect on the ring radius, but yields a storage ring with non-zero magnetic eld.



FIG.1: (Color online) Four circular coils with average diam – eter 10 cm make up the storage ring (thin circle) to which the axial wire adds an adjustable azim uthal magnetic eld. The square coils localise the atom s (sm all cigar) at the top of the ring in a magneto-optical (magnetic) trap if the left and right square coil pairs have the same (opposite) currents.

The theory of the storage ring for cold atom s is relatively simple, as the spatio-tem poral evolution of each atom can be accurately described using the equations of motion arising from the Stem-Gerlach potential. A fter choosing suitable G aussian initial position and velocity distributions it is possible to perform a M onte C arlo simulation to build up a spatio-tem poral atom ic density m ap.

By comparison with a 3D m odel, we have found that a 1D m odel is su cient, i.e. the rigid pendulum equation:

⁽⁰⁾ (t) =
$$(g=R_0) \sin ;$$
 (2)

where and $! = \frac{d}{dt}$ are the angular position and velocity of an atom, g is the acceleration due to gravity and R₀ is the radius of the storage ring. Note that Eq.2 can be integrated analytically (using $\frac{d^2}{dt^2} = !\frac{d!}{d}$) given the initial angle (₀) and angular velocity (!₀) of an atom, to determ ine the tim e-averaged relative probability of nding an atom at a given angle:

P()/1=! =
$$1 = \frac{p}{!_0^2 + 2g(\cos_0 \cos_0) = R_0}$$
: (3)

In our vertically-oriented storage ring there is therefore a high time-averaged probability of nding atoms in the region where they travel the slowest { the top of the ring. The integral of Eq. 3, leads to an expression for t() in terms of an elliptic integral, which can be inverted to nd (t). There are two kinds of trajectories in the ring: if $\cos(\max x) = \frac{\pi_0}{J_{2g}}!_0^2 + \cos_0 j > 1$ an atom will always rotate in the same direction, but if $\cos(\max x) = 1$ (cf. Fig. 2 (b)) the atom will reverse its direction around the ring at the turning points = $\max x$.

The theoretical time-dependent angular distribution of atom s can be seen in Fig.2. An atom ic cloud centered in a parabolic potential can expand or shrink, but willm aintain the same cloud shape. A circular atom ic trajectory has a potential which is only parabolic to second order, and it is the higher order e ects which cause an atom ic cloud released from the top of the ring (=0) to break into two halves (Fig. 2 (a)), leaving a near-zero probability at the top of the ring after around 300 m s. These two halves return to the top of the ring after a further 300 m s and have a non-zero probability of remaining at the top of the ring for all subsequent times. The time for the two halves of the atom ic cloud to 'recombine' has only a weak dependence on the initial atom ic temperature. G ravity is the dom inant e ect for cold atom s, leading to a hot' velocity of 1:4 m s¹ at the bottom of the ring { a good location for future studies of high-energy (20mK) collisions between BECs [11].

A lthough the spread of initial thermal velocities quickly ensures that atom s reach the top of the ring at di erent times, in Fig. 2(a) we still see slow ly decaying bechos' of the originally localised spatial atom ic cloud in the form of time-varying bin odal perturbations of the

distribution. The angular width of these echos is approximately proportional to the initial cloud width, so the echos and their bim odal nature become more pronounced at higher tem peratures.

In m any of our ring experiments the atom s/BEC swere not launched from 0 = 0 (Fig.2(a)). A lthough the total number of atom s observed is similar to the $0 \notin 0$ case, the size of the returning $0 \notin 0$ cloud is much smaller and clearer due to focusing at the turning points of its motion (Fig.2(b)).



FIG.2: (Color online) A M onte-Carlo simulation of atom ic dynam ics in the vertically-oriented storage ring for a 2 K initial cloud of atom s. Im ages (a) and (b) represent lossless atom dynam ics for a cloud released at $_{0} = 0$ and $_{0} = 60$ m rad respectively. The white lines m ark the extent of our 110 m rad experim ental view ing region.

We now turn to the experimental setup, for which m any of the details are in Refs. [7, 12]. Our double m agneto-optical trap (MOT) [13] collects 10^{9} ⁸⁷Rb atom s in both our high and low pressure MOT chambers. The storage ring can be loaded directly from the low pressure MOT or IO e-Pritchard (IP [14]) trap without the need for any additional magnetic guiding or transfer, i.e. the MOT and IP trap are at the top of the ring. Releasing atom s at the top of a vertical ring ensures insensitivity to m agnetic ring corrugations and a fast rotation frequency.

The advantage of our hybrid magnetic trap (Fig. 1) is that, using the same coils, it can be (i) a MOT, (ii) an IP magnetic trap (with trap frequencies $_r = 230 \text{ Hz}$; $_z = 10 \text{ Hz}$) or iii) a toroidal magnetic storage ring with radial gradient 230 G/cm. In modes (i) and (ii) the magnetic eld has an adjustable aspect ratio in the azim uthal direction. The four 2 turn 500 A circular coils in Fig.1 create a toroidal magnetic quadrupole eld, con ning the atom s to a ring. The four 3 turn 500 A square binch' coils are wired in pairs, and con ne the atom s to a localised section of the ring for Io e-P ritchard trapping of the atom s. A n azim uthal eld can be added to the storage ring via an axial wire. The large coil sizes ensure good optical access, and low magnetic eld noise. The absorption in aging (gravity) direction is the x (y) axis.

A fler loading the low pressure MOT, the atom ic cloud is optically pumped, loaded into the IP trap, m agnetically compressed, and then evaporatively cooled to an adjustable temperature. The atom s can then be smoothly loaded into the storage ring in 20m s. We are able to form Bose-Einstein condensates containing N₀ = 2 10° f = 2;m_F = 2i atom s at the top of the storage ring [7] at a typical nalRF evaporation frequency of 750 kHz.

In order to com pare our experim ental ring data (Fig. 3) with our theoretical model (Fig. 2), we have found it convenient to model the total number of atoms in the 'viewing window' of our absorption in aging system as a function of time. Our CCD camera has an area of 4:8 $6:4 \text{ mm}^2$; with a magni cation of 1:20(1):W e have released cold atom ic clouds with a variety of nalevaporation temperatures into a storage ring with no azimuthal eld. In accordance with theory, the atom s dis-400 m s, reappear shortly afterwards, appear after 300 and are present at all subsequent times. However, there is a marked variation in the storage ring dynamics with atom ic cloud release tem perature. This can be explained in term s of non-adiabatic M a jorana spin- ip transitions [15]. The storage ring has a ring of zero magnetic eld, and colder atom s pass closer to the m agnetic eld zero, and are selectively removed from the storage ring.

If we apply a constant magnetic eld across the entire storage ring, perpendicular to the ring axis, it is possible to have only two places in the ring with zero magnetic eld, however there will be a strong angular variation in the trapping potential. A novel feature of our storage ring is that the hole' in our quartz vacuum chamber (Fig. 1) allow s us to use a wire along the axis of the storage ring. This means that we can generate an azim uthalm agnetic bias eld B = 0 10G around the ring which transform s the radialm agnetic potential from a cone to a hyperbola, rem oving the ring of zero magnetic eld and drastically reducing atom ic loss.

This di erence can be seen in the experiments of Fig.3(a) and (b), where 2 K atom ic clouds are released into a ring without and with an azim uthal eld, respectively. A quantitative contrast of the number of visible atom s in each case is shown in Fig.4, as well as a comparison to the M onte C arlo theory of Fig.2. The di erence the azim uthal eld m akes is even greater with condensates: without it BECs vanish before completing one revolution, but with an azim uthal eld m ultiple revolutions of a condensate are possible with low loss (Fig.3(c)).



FIG.3: (Color online) Experimental storage ring dynamics for 2 K atom swith (a) B = 0G and (b) B = 10G. In (c) multiple revolutions of a BEC in a ring with B = 10G are seen. Numbers denote time in ms, absorption in ages are 0.4 5.4 mm².

Note that when we perform ed experiments with an azim uthal eld, the atom s/BECs were not launched from $_0 = 0$: If atoms are launched from $_0 = 0$ (Fig. 2(a)), then although the total number of atom s observed (Fig. 4) is similar to the $_0 \notin 0$ case, the size of

the returning $_{0} \in 0$ cloud is much smaller and clearer due to focusing in the ring (Figs. 2(b), 3(b,c)).

W e do not expect to see phase- uctuations [16] in our condensate before or after propagation in the ring. These e ects have been studied in highly elongated conden-



FIG.4: The relative atom ic population, P (t); viewed by the CCD camera as a function of time for 2 K atom s. Experimental data with B = 0 G and B = 10 G are shown with circles and triangles respectively. The relative M onte C arlo atom ic population in the viewing region of Fig.2(a) (dashed curve) and Fig.2(b) (solid curve) are shown for comparison. G ood quantitative agreement can be seen between the B = 10 G data and the M onte C arlo simulation of an atom ic cloud released from $_0 = 60$ m rad. The B = 0 G data only ts the expected dashed theoretical curve initially due to large M a prana losses.

sates in which the BEC coherence length is less that the length of the condensate. We form the condensate in only a moderately elongated trap, and the azim uthal expansion process is rapid enough that the (density-dependent) phase uctuations do not have time to develop.

Sagnac interferom etry will be performed by locating the condensate at the exact top of the ring, and incoherently splitting the sample by simply releasing it (Fig. 5). We are currently looking for interference fringes after a single revolution. Coherent splitting of the BEC will be achieved using Bragg scattering [17] to send BEC wavepackets in both directions around the ring. Note that ideally the phase sensitivity of a BEC interferom eter scales like $_{\rm BEC}$ / N 1 where N is the number of atom s (cf. $_{\rm T}$ / N $^{1=2}$ for therm al atom s), however the increased sensitivity is only possible if the BEC s are prepared in number states [18].



FIG. 5: A BEC at the top of the ring was split into two counter-rotating clouds by the faster-than-harm onic circular potential. These split atom ic clouds can be seen recombining here after one revolution in the ring. T in es are in m s, each im age is 150 3400 m^2 and is taken after a 3m s ballistic expansion.

In conclusion, we have demonstrated a storage ring for cold 87 Rb atom ic clouds and BECs with an area of 7200 mm². An azim uthal bias eld around the ring enables low -loss BEC propagation with heating of less than 50 nK /s. Our goal is a highly sensitive Sagnac atom interferom eter, in which we are aided by our unprecedented ring area. Rotation sensitivity for a single revolution of $= h = (8m R_0^{2^{11}} \overline{N}) = 310^{11} rad=s$ is feasible.

If one cannot create an azim uthal magnetic eld, it is still possible to prevent spin- ip losses by using an adjustable-radius time-orbiting ring trap (TORT) [19]. We recently learnt that a team in Berkeley has created the rst TORT for BECs [20], with ring area 5 mm^2 . Large BEC swere also form ed in an optically-plugged ring trap with ring area 1 mm^2 [21].

W e are grateful for helpful discussions with K.Burnett and E.H inds. This work was supported by the UK EPSRC and the University of Strathclyde.

- [1] C. S. Adam s and E. Riis, Prog. Quant. Electr. 21, 1 (1997); E. A. Hinds and I. G. Hughes, J. Phys. D 32, R119 (1999).
- [2] S. Chu, L. Hollberg, J. E. B prkholm, A. Cable, and A. Ashkin, Phys. Rev. Lett. 55, 48 (1985); E. L. Raab, M. Prentiss, A. Cable, S. Chu, and D. E. Pritchard, Phys. Rev. Lett. 59, 2631 (1987).
- [3] M. H. Anderson et al., Science 269, 198 (1995); K. B.
 Davis et al., Phys. Rev. Lett. 75, 3969 (1995); C. C.
 Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, Phys. Rev. Lett. 79, 1170 (E) (1997).
- [4] M G.Sagnac, C R.Acad.Sci.157, 708 (1913).
- [5] F.M.H.Crom proets et al., Nature 411, 174 (2001).
- [6] J.A. Sauer, M.D. Barrett, and M.S. Chapman, Phys. Rev.Lett. 87, 270401 (2001).
- [7] C.S.Garvie, E.R iis and A.S.A mold, Laser Spectroscopy XVI, P.H annaford et al.eds., p178 (W orld Scienti c, Singapore, 2004); see also www photonicsphys.strath.ac.uk.
- [8] S.W u, W .Rooijakkers, P.Striehl, and M .Prentiss, Phys. Rev. A 70, 013409 (2004).
- [9] T.L.Gustavson, P.Bouyer and M.A.Kasevich, Phys. Rev.Lett. 78, 2046 (1997).
- [10] R.H.Good, Eur.J.Phys.22, 119 (2001).
- [11] N.R.Thomas, N.Kj rgaard, P.S.Julienne and A.C. W ilson, Phys.Rev.Lett. 93, 173201 (2004); Ch.Buggle, J.Leonard, W .von K litzing and J.T.M.W alraven, Phys.Rev.Lett. 93, 173202 (2004).
- [12] A. S. Amold and E. Riis, J. Mod. Opt. 49, 959-964 (2002).
- [13] K.G ibble, S.Chang and R.Legere, Phys. Rev.Lett. 75, 2666 (1995); C.J.M yatt et al., Opt.Lett.21, 290 (1996).
- [14] D E. Pritchard, Phys. Rev. Lett. 51, 1336 (1983).
- [15] W . Petrich, M. H. Anderson, J. R. Ensher, and E. A. Comell, Phys. Rev. Lett. 74, 3352 (1995).
- [16] D.S. Petrov, G.V. Shlyapnikov and J.T.M. Walraven, Phys. Rev. Lett. 87, 050404 (2001); S. Dettm er et al., ibid.87,160406 (2001); D. Hellweg et al., ibid.91,010406 (2003); S.Richard et al., ibid.91,010405 (2003).
- [17] J. Stenger et al, Phys. Rev. Lett. 82, 4569 (1999); M.
 Kozum a et al, Phys. Rev. Lett. 82, 871 (1999).
- [18] J. Jacobson, G. B jork and Y. Yam am oto, Appl. Phys. B 60, 187 (1995).
- [19] A.S.Amold, J.Phys.B 37, L29 (2004).
- [20] S.Gupta et al, Phys.Rev.Lett.95, 143201 (2005).
- [21] D. S. Naik, S. R. M uniz and C. Raman, Phys. Rev. A 72,051606 (R) (2005).