

Non-equilibrium Differential Conductance through a Quantum Dot in a Magnetic Field

A.C.Hewson and J. Bauer

Department of Mathematics, Imperial College, London SW7 2AZ, UK

A.Oguri

Department of Material Science, Osaka City University, Sumiyoshi-ku, Osaka 558-8585 Japan

(Dated: March 23, 2022)

We derive an exact expression for the differential conductance for a quantum dot in an arbitrary magnetic field for small bias voltage. The derivation is based on the symmetric Anderson model using renormalized perturbation theory and is valid for all values of the on-site interaction U including the Kondo regime. We calculate the critical magnetic field for the splitting of the Kondo resonance to be seen in the differential conductivity as function of bias voltage. Our calculations for small field show that the peak position of the component resonances in the differential conductance are reduced substantially from estimates using the equilibrium Green's function. We conclude that it is important to take the voltage dependence of the local retarded Green's function into account in interpreting experimental results.

PACS numbers:

I. INTRODUCTION

Measurements of the temperature dependence of the differential conductance dI/dV_{ds} through a quantum dot in the linear response regime have been shown to be in good agreement with theoretical predictions based on magnetic impurity models [1, 2]. Measurements of dI/dV_{ds} through a quantum dot in the presence of an applied magnetic field, however, have revealed an apparent discrepancy between the magnetic splitting in the Kondo peaks deduced experimentally and theoretical predictions based on the same models [3, 4, 5]. The differential conductance measurements are made in the presence of a finite applied bias voltage V_{ds} , which is a situation for which there are few theoretical predictions. There are calculations based on the non-crossing approximation (NCA) [6]. This method, however, is known to break down in the Fermi liquid regime in the equilibrium case, so there is some uncertainty about the validity of the predictions in this case. There are also some recent perturbation calculations of Fuji and Uedo [7] using the Keldysh formalism, taking diagrams up to fourth order in the on-site interaction U into account. This approach should reliably describe the trends in the approach to the strong correlation limit, as the interaction strength is increased, but cannot fully describe the Kondo regime, where there is an exponential renormalization of the Kondo resonance width as a function of U . Rosch et al. have calculated the differential conductance using a perturbative renormalization group calculation for the exchange coupling J to leading order in $1/\ln(V_{ds}/T_K)$, which is appropriate for the very high voltage regime ($V_{ds} \gg T_K$) [12] (see also [13]). This approach is, however, not suitable for investigating the Kondo regime, where V_{ds} and the magnetic field are of order of the Kondo temperature T_K .

More accurate methods [8, 9, 10, 11] for the Kondo regime exist for calculations in an applied magnetic field, which have been used to interpret the experimental results [3, 4, 5], but they are restricted to the linear response (equilibrium) regime. It is not clear that using the equilibrium response for dI/dV_{ds} in the presence of an applied bias voltage will lead to reliable predictions. In this paper we examine the effect of a finite applied bias voltage in the small bias regime. In the limit of a small magnetic field B , we calculate asymptotically exactly the shifts of the peaks of the differential conductance of particular spin component for small, but finite V_{ds} . The results are based on renormalised perturbation theory and numerical renormalization group (NRG) calculations. We also deduce the value of the critical field B_c for two distinct peaks to be seen in dI/dV_{ds} . The knowledge of dI/dV_{ds} to order V_{ds}^2 for arbitrary values of B is sufficient to determine B_c exactly.

In our treatment the quantum dot will be described by a single impurity Anderson model with particle-hole symmetry coupled via left and right leads to two reservoirs of free electrons. The Anderson model has the form,

$$H_{AM} = \sum_{\sigma} \epsilon_{d\sigma} d_{\sigma}^{\dagger} d_{\sigma} + U n_{d,\uparrow} n_{d,\downarrow} + \sum_{k,\sigma} (V_k d_{\sigma}^{\dagger} c_{k,\sigma} + V_k^* c_{k,\sigma}^{\dagger} d_{\sigma}) + \sum_{k,\sigma} \epsilon_k c_{k,\sigma}^{\dagger} c_{k,\sigma}, \quad (1)$$

where $\epsilon_{d\sigma} = \epsilon_d - \sigma g \mu_B B/2$ is the energy of the impurity levels with spin $\sigma = \pm 1$ in a magnetic field B , U is the interaction at the impurity site, and V_k the hybridization matrix element to a band of conduction electrons with energy ϵ_k . In the wide band limit the hybridization weighted density of states, $\Delta(\omega) = \pi \sum_k |V_k|^2 \delta(\omega - \epsilon_k)$, can be taken as a constant Δ . Taking this limit justifies the neglect of any magnetic field term acting on the conduction

electrons in (1) (Clogston-Anderson compensation theorem [14]). Any polarization of the conduction electrons only affects the impurity via the hybridization function $\Delta(\omega)$ and any change to the conduction band density of states due to an applied magnetic field is only at the band edges ($\pm D$), and hence negligible if $g\mu_B B \ll D$. Therefore, if $\Delta(\omega)$ is independent of ω the magnetic field does not need to be included in the electron band.

The single impurity model can be applied to a quantum dot which is equally coupled via leads to left and right reservoirs. In this situation the charge on the quantum dot is coupled only to an even combination of states from the left and right channels by an effective hybridization V_k , so it can be mapped into a single channel model. If there is a potential difference eV_{ds} due to a bias voltage V_{ds} (e is the electronic charge), then the chemical potentials in the left and right reservoirs differ from the average chemical potential μ by $eV_{ds}/2$ and $-eV_{ds}/2$, respectively.

A general expression for the current I due to the applied voltage through a quantum dot has been given by Hershfield et al. [15] and Meir and Wingreen [16, 17], and specialising it to this symmetric case it takes the form,

$$I = \frac{e\Delta}{h} \sum_{\sigma} \int_{-\infty}^{\infty} d\omega [f_L(\omega) - f_R(\omega)] [-\text{Im}G_{d\sigma}^r(\omega, V_{ds})], \quad (2)$$

where $G_d^r(\omega, V_{ds})$ is the steady state retarded Green's function on the dot site, and $f_L(\omega)$, $f_R(\omega)$ are Fermi distribution functions for the electrons in the left and right reservoirs, respectively, and h is Planck's constant.

The retarded Green's function $G_{d\sigma}^r(\omega, V_{ds})$ can be written in terms of a self-energy $\Sigma_{\sigma}(\omega, V_{ds})$,

$$G_{d\sigma}^r(\omega, V_{ds}) = \frac{1}{\omega - \epsilon_d + i\Delta - \Sigma_{\sigma}(\omega, V_{ds})} \quad (3)$$

Oguri [18, 19] has shown that this Green's function for the symmetric model ($\epsilon_d = -U/2$) in the Fermi liquid regime in the absence of a magnetic field to order ω^2 and $(eV_{ds})^2$, can be expressed in the form,

$$G_{d\sigma}^r(\omega, V_{ds}) = \frac{z}{\omega + i\tilde{\Delta} - \tilde{\Sigma}(\omega, V_{ds})} \quad (4)$$

where $\tilde{\Sigma}(\omega, V_{ds})$ is a renormalized self-energy [20] given by

$$\tilde{\Sigma}(\omega, V_{ds}) = -ic \left[\omega^2 + \frac{3}{4}(eV_{ds})^2 \right], \quad \text{with } c = \frac{1}{2\tilde{\Delta}} \left(\frac{\tilde{U}}{\pi\tilde{\Delta}} \right)^2 \quad (5)$$

expressed in terms of the renormalized parameters $\tilde{\Delta}$ and \tilde{U} . These parameters [20] are defined in terms of the local impurity self-energy $\Sigma(\omega)$ and local irreducible 4-vertex $\Gamma_{\uparrow\downarrow}(\omega_1, \omega_2, \omega_3, \omega_4)$ [20] for $V_{ds} = 0$,

$$\tilde{\Delta} = z\Delta, \quad \tilde{U} = z^2\Gamma_{\uparrow\downarrow}(0, 0, 0, 0), \quad (6)$$

where z is given by $z = 1/(1 - \Sigma'(0))$. The renormalized parameters $\tilde{\Delta}$ and \tilde{U} , can be calculated directly from NRG calculations, as described in references [22, 23], where they are given in terms of the 'bare' parameters of the model Δ and U . Alternatively they can also be deduced from the Bethe ansatz results [20].

We can generalise this result by including a magnetic field term [23, 24] to lowest order in B , and then $G_{d\sigma}^r(\omega)$ takes the form,

$$G_{d\sigma}^r(\omega) = \frac{z}{\omega + \sigma\tilde{\eta}b + i\tilde{\Delta} - \tilde{\Sigma}_{\sigma}(\omega, V_{ds})}, \quad (7)$$

where $\sigma = \pm 1$, $b = g\mu_B B/2$, and $\tilde{\eta}$ is a renormalization parameter defined as $\tilde{\eta}(b) = z(b)(1 - \Sigma_{\uparrow}(0, b)/b)$. This renormalized parameter can be deduced from NRG calculations as shown in reference [23, 25]. We give an example in the Kondo regime for $U/\pi\Delta = 4$ in figure 1, where the ratio of the parameters to their values in zero field are plotted as a function of $\log(b/T_K)$. As $b \rightarrow 0$, $\tilde{\eta}(b)$ is independent of b such that $\tilde{\eta}(b) \rightarrow R$, where R is the Wilson or ' χ/γ ' ratio for zero magnetic field. In terms of the renormalized parameters for finite field $R(b)$ is given by $R(b) = 1 + \tilde{U}(b)\tilde{\rho}_d^0(0, b)$ [20, 22], where the free quasiparticle density of states $\tilde{\rho}_{d,\sigma}^0(\omega, b)$ is given by

$$\tilde{\rho}_{d,\sigma}^0(\omega, b) = \frac{\tilde{\Delta}(b)/\pi}{(\omega + \sigma b\tilde{\eta}(b))^2 + \tilde{\Delta}^2(b)}, \quad (8)$$




FIG. 1: b dependence of the renormalized parameters for strong coupling ($U/\pi\Delta = 4$), in relation with their $b = 0$ values, $\tilde{U} = \pi\tilde{\Delta} = 4T_K = 0.0817\Delta$. Note the logarithmic scale of the y-axis.

which is independent of the spin index σ for $\omega = 0$. There are in general B^2 and higher order terms in the self-energy but for the moment we work only to first order in B so the renormalized self-energy $\tilde{\Sigma}_\sigma(\omega, V_{ds})$ is still given by equation (5) with

$$c = c(b) = \frac{\pi\tilde{U}^2(b)[\tilde{\rho}_d^0(0, b)]^3}{2}, \quad (9)$$

which reduces to (5) for $b \rightarrow 0$.

II. WEAK FIELD LIMIT

We look at the effects of the different contributions to dI/dV_{ds} in turn. We consider the simplest situation to begin with and ignore the renormalized self-energy term in (7). The differential conductance (2) at $T = 0$ then takes the form,

$$\frac{dI}{dV_{ds}} = \frac{e^2}{h} \sum_{\nu} \frac{\tilde{\Delta}^2}{(eV_{ds}/2 - \nu\tilde{\eta}b)^2 + \tilde{\Delta}^2}, \quad (10)$$

where $\nu = \pm 1$. The total result is expressed as a sum over $\nu = \pm 1$, rather than as a sum over $\sigma = \pm 1$, as both spin up and spin down states contribute to each resonance. Note the result does not simply correspond to the non-interacting case, as it includes the many-body renormalization factor $\tilde{\eta}$. It corresponds to the case in which the interaction between the renormalized quasiparticles (\tilde{U}) is neglected. In the Kondo regime $\tilde{\eta} = 2$, so the effective Zeeman splitting of the resonance in this limit is twice the Zeeman splitting (2b) for non-interacting electrons ($U = 0$). Note that the energy level on the dot is at the average chemical potential μ , which is shifted by $eV_{ds}/2$ with respect to the value at zero bias. The individual terms in (10) for each ν have maxima at $\pm 2\tilde{\eta}b$. The maxima of the sum in general occur at a shifted position $eV_{ds}^{\pm} = \pm 2\tilde{\eta}b f_c(\tilde{\eta}b, \tilde{\Delta})$ with

$$f_c(\tilde{\eta}b, \tilde{\Delta}) = \left[1 - \left(1 - \left[1 + (\tilde{\Delta}/\tilde{\eta}b)^2 \right]^{1/2} \right)^2 \right]^{1/2}, \quad (11)$$

provided that $\tilde{\eta}b > \tilde{\Delta}/\sqrt{3}$. For $\tilde{\eta}b \gg \tilde{\Delta}$ can be approximated by $f_c(\tilde{\eta}b, \tilde{\Delta}) \simeq 1 - \frac{1}{8} \frac{\tilde{\Delta}^4}{(\tilde{\eta}b)^4}$. The peak splitting is therefore $e(V_{ds}^+ - V_{ds}^-) = 4\tilde{\eta}b f_c(\tilde{\eta}b, \tilde{\Delta})$. A maximum of the differential conductance, therefore, occurs when one of the quasiparticle peaks is coincident with left Fermi level at $\mu + eV_{ds}/2$ and at the same time the other peak coincides with the right Fermi level, $\mu - eV_{ds}/2$. This is illustrated in figure 2. In the interpretation of the experimental results




FIG. 2: A schematic plot of the spectrum on the dot and chemical potentials for left/right lead ($\mu \pm eV/2$) and dot (μ) for (a) zero bias and zero magnetic field and (b) finite voltage and finite field.

of dI/dV_{ds} the splitting of the Kondo resonance Δ_{Kondo} was identified with the voltage splitting $e(V_{ds}^+ - V_{ds}^-) = \Delta_{\text{Kondo}}$ [4, 5]. However, for the renormalized quasiparticles the Kondo splitting is $\Delta_{\text{Kondo}} = 2\tilde{\eta}b$, which substituted in the above result gives $e(V_{ds}^+ - V_{ds}^-) = 2\Delta_{\text{Kondo}}$. The factor of 2 difference is due to the many-body shift in the energy level on the dot, which was not taken into account in the interpretation of the experimental results in [4, 5].

We should also take into account the terms in the renormalized self-energy $\tilde{\Sigma}(\omega, V_{ds})$. For the moment we ignore the V_{ds} dependence of $\tilde{\Sigma}(\omega, V_{ds})$, and then

$$\frac{dI}{dV_{ds}} = \frac{e^2 \tilde{\Delta}}{h} \sum_{\nu} \left. \frac{\tilde{\Delta} - \tilde{\Sigma}^I(\omega)}{(\omega - \nu \tilde{\eta}b)^2 + (\tilde{\Delta} - \tilde{\Sigma}^I(\omega))^2} \right|_{\omega=eV_{ds}/2}, \quad (12)$$

where $\tilde{\Sigma}^I(\omega, V_{ds})$ is the imaginary part of $\tilde{\Sigma}(\omega, V_{ds})$ as given in equation (5). If the component spectral densities ($\nu \pm 1$) have maxima at ω_{\pm} then this expression has maxima as a function of V_{ds} for $2\omega_{\pm}$. Working to lowest order in b , we find from the exact result for the self-energy to order ω^2 , that for a maximum,

$$\omega = \pm \tilde{\eta}b + \frac{\tilde{\Delta}}{2} \frac{\partial \tilde{\Sigma}^I}{\partial \omega} = \pm \tilde{\eta}b - \omega \tilde{\Delta} c(b) \quad (13)$$

The peak position in the resonances in the spectral density to first order in b is given by

$$\omega_{\pm} = \frac{\pm \tilde{\eta}b}{1 + \tilde{\Delta} c(b)} \quad (14)$$

For $b \rightarrow 0$ find $\tilde{\Delta} c(b) \rightarrow \frac{1}{2} \left(\frac{\tilde{U}}{\pi \tilde{\Delta}} \right)^2 = \frac{1}{2} (R - 1)^2$ and so this result corresponds to that of Logan and Dickens [9], and as derived from the renormalized perturbation calculations [21]. It shows that the effective Zeeman splitting of the free quasiparticles, $2\tilde{\eta}b$, is reduced in the spectral densities by a factor arising from the ω^2 term in the imaginary part of the self-energy. In the Kondo limit, $\tilde{U} = \pi \tilde{\Delta} = 4T_K$, so $\tilde{U}/\pi \tilde{\Delta} = 1$ and $\tilde{\eta} = R = 2$; hence the peak position of $2b$ is reduced by a factor of $2/3$. This effect will be reflected in the voltage difference between the differential conductance peaks,

$$e(V_{ds}^+ - V_{ds}^-) = \frac{4\tilde{\eta}b}{1 + \tilde{\Delta} c(b)} f_c(\tilde{\eta}b, \tilde{\Delta}), \quad (15)$$

where f_c is a correction factor due to the overlap of resonances similar as above, which can be computed numerically. However, in calculating this voltage difference we cannot neglect the voltage dependence of the self-energy.

When we take the V_{ds} dependence of the retarded Green's function into account,

$$\frac{dI}{dV_{ds}} = -\frac{e^2 \Delta}{h} \sum_{\sigma} \text{Im} G_{d\sigma}^r(eV_{ds}/2, V_{ds}) + \frac{2e\Delta}{h} \sum_{\sigma} \int_0^{eV_{ds}/2} d\omega \left[-\text{Im} \frac{\partial G_{d,\sigma}(\omega, V_{ds})}{\partial V_{ds}} \right]. \quad (16)$$




FIG. 3: The shift of the component resonance ($\nu = 1$) in the differential conductance (units of e^2/h) in a magnetic field for $b/\tilde{\Delta} = 0.05$ as a function of the bias voltage $eV_{ds}/\tilde{\Delta}$, according to the inclusion of different contributions as described in the text. The arrows indicate the respective maxima.

In considering the contribution from the first term on the right hand side we can replace the eV_{ds} in the self-energy by 2ω , and then see where it has a maximum as a function of ω . This will correspond to the calculation we have just done except that the ω^2 term in $\tilde{\Sigma}_I(\omega)$ will have an extra factor of 4, as we can see from equation (5). We get the modified result,

$$e(V_{ds}^+ - V_{ds}^-) = \frac{4\tilde{\eta}b}{1 + 4\tilde{\Delta}c(b)} f_c(\tilde{\eta}b, \tilde{\Delta}). \quad (17)$$

In the Kondo regime and for $b \rightarrow 0$ the reduction factor arising from the self-energy including the voltage dependence is now $1/3$, so the splitting on including this term is reduced by an extra factor of 2 when the voltage dependence in the retarded Green's function is taken into account. We do need to take account, however, of the second term on the right hand side of equation (16) involving an integral over ω . This term can be written as

$$\frac{3c(b)e^2}{h} V_{ds} \tilde{\Delta} \sum_{\nu} \int_0^{eV_{ds}/2} d\omega \frac{(\omega + \nu\tilde{\eta}b)^2 - (\tilde{\Delta} + c(b)(\omega^2 + 3(eV_{ds})^2/4))^2}{[(\omega + \nu\tilde{\eta}b)^2 + (\tilde{\Delta} + c(b)(\omega^2 + 3(eV_{ds})^2/4))^2]^2}. \quad (18)$$

If we plot all the contributions to dI/dV_{ds} in the very weak field regime then, due to overlap, no magnetic field splitting can be observed. We can calculate, however, the shifts in the component resonance for $\nu = \pm 1$. In figure 3 we plot the terms in the differential conductance (in units of e^2/h) given by equation (16) as a function of $eV_{ds}/\tilde{\Delta}$, where we use $\nu = 1$ in (12) for the first term in (16) and (18) for the second term in (16). We take values corresponding to the Kondo regime, with $R = \tilde{\eta} = 2$, $b/\tilde{\Delta} = 0.05$ ($\pi\tilde{\Delta} = 4T_K$). We have distinguished between the different contributions, (a) is the case for the non-interacting quasiparticles, (b) includes the ω^2 term in the renormalized self-energy, (c) includes the first term in equation (16), and (d) takes into account the full expression including the integral term in (18). We see that the integral term arising from the voltage dependence of $G_{d,\sigma}(\omega, V_{ds})$ causes a significant further reduction of the magnetic shift beyond that estimated from the first term in equation (16). In an experimental conductance measurement, this is not observable, however, due to the overlap of the two components.

III. CRITICAL FIELD

Our derivation is restricted to the regime where eV_{ds} is small compared to $\tilde{\Delta}$. These results are sufficient for us to deduce the critical value of the magnetic field b_c at which two distinct peaks begin to appear in the total differential




FIG. 4: The total differential conductance (in units of e^2/h) in the Kondo regime for larger magnetic field values, calculated using equation (16) taking into account the full selfenergy expansion from (21). These results are asymptotically exact for $eV_{ds}/\tilde{\Delta} \ll 1$ and approximate, based on a second order expansion in eV_{ds} for larger values.

response. For values of $b < b_c$ the differential conductance will have a maximum at $V_{ds} = 0$, and for $b > b_c$ this will become a minimum. From the coefficient of dI/dV_{ds} to order V_{ds}^2 we can determine the point at which it changes sign as a function of b , and hence determine b_c . The contribution from the integral term in equation (18) to order V_{ds}^2 can be evaluated trivially, as it is sufficient to this order to put $\omega = V_{ds} = 0$ in the integrand. As a first estimate using the above results the value of b_c can be calculated analytically, and the result expressed entirely in terms of $\tilde{\Delta}$ and the Wilson ratio $R = \eta(0) = 1 + \tilde{U}/\pi\tilde{\Delta}$,

$$\frac{b_c^2}{\tilde{\Delta}^2} = \frac{\sqrt{9 + 20(R-1)^2(1 + 5(R-1)^2)} - 3}{10R^2(R-1)^2}. \quad (19)$$

In the non-interacting case, $R = 1$ and $b_c/\Delta = 1/\sqrt{3} = 0.577$, which corresponds to $f_c(\tilde{\eta}b, \tilde{\Delta}) = 0$ from equation (11), and in the Kondo regime, $R = 2$, $\tilde{\Delta} = 4T_K/\pi$, and $b_c/T_K = 0.582$, with T_K given by

$$T_K = \Delta \left(\frac{U}{2\Delta} \right)^{1/2} e^{-\pi U/8\Delta + \pi\Delta/2U}. \quad (20)$$

If the V_{ds} dependence of the Green's function is neglected the result in the Kondo regime is $b_c/T_K = 0.491$; significantly smaller than if this term is included.

The estimated critical magnetic field is comparable with $\tilde{\Delta}$, and for $U \neq 0$ it may not be sufficient to work to linear order in B . It is possible to work with an arbitrary magnetic field, but in this case the renormalized parameters become field dependent. The renormalized self-energy to order ω^2 and V_{ds}^2 can be expressed in the form,

$$\tilde{\Sigma}_\sigma(\omega, V_{ds}) = -c(b) \left[i \left(\omega^2 + 3 \left(\frac{eV_{ds}}{2} \right)^2 \right) + \frac{\sigma b \tilde{\eta}(b)}{\tilde{\Delta}(b)} \left(\alpha_\omega(b) \omega^2 + \alpha_V(b) \left(\frac{eV_{ds}}{2} \right)^2 \right) \right], \quad (21)$$

where $c(b)$ is given in equation (9). The coefficients for the expansion of the real part of $\tilde{\Sigma}_\sigma(\omega, V_{ds})$ are

$$\alpha_\omega(b) = 2 + \frac{2I(b)\tilde{\Delta}(b)}{\tilde{\xi}(b)[\tilde{\rho}_d^0(0, b)]^2}, \quad (22)$$

where $\tilde{\xi}(b) = \pi\tilde{\rho}_d^0(0, b)\tilde{\eta}(b)b$, $I(b)$ is the integral

$$I(b) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} G_{\downarrow}^0(\omega'') G_{\downarrow}^0(\omega'' + \omega') [G_{\uparrow}^0(\omega')]^3 \frac{d\omega''}{2\pi} \frac{d\omega'}{2\pi}, \quad (23)$$

where $[G_{\sigma}^0(\omega)]^{-1} = \omega + \sigma\tilde{\eta}b + i\tilde{\Delta}\text{sgn}(\omega)$ is the renormalised free propagator for $T = 0$. $\alpha_{\omega}(b)$ is obtained from the second order derivative of the renormalized self-energy evaluated at $\omega = 0$ using the renormalized perturbation theory to order \tilde{U}^2 . This reduces to the exact result of Yamada [27] for the symmetric model for $b = 0$. The corresponding coefficient $\alpha_V(b)$ for finite voltage is calculated from the \tilde{U}^2 contribution to the retarded self-energy in the Keldysh formalism [26], where the propagator $G_{0,\sigma}^{-}(\omega, V_{ds})$ taken to order V_{ds}^2 for $T = 0$ is

$$G_{0,\sigma}^{-}(\omega, V_{ds}) = G_{\sigma}^0(\omega) - \frac{i\tilde{\Delta}\delta'(\omega)(eV_{ds})^2/4}{(\omega - \sigma\tilde{\eta}b)^2 + \tilde{\Delta}^2}, \quad (24)$$

where $\delta'(\omega)$ is the derivative of the delta-function, together with a similar equation for $G_{0,\sigma}^{+-}(\omega, V_{ds})$. The result for $\tilde{\alpha}(b)$ is

$$\tilde{\alpha}_V(b) = 1 + \frac{\tilde{\Delta}(b)}{2\tilde{\xi}(h)\tilde{\eta}(b)b} \left[1 - \frac{\tilde{\eta}(b)b}{\tilde{\Delta}(b)} \tan^{-1} \left(\frac{\tilde{\eta}(b)b}{\tilde{\Delta}(b)} \right) \left(4 + \frac{\tilde{\Delta}(b)}{\tilde{\xi}(h)\tilde{\eta}(b)b} \right) \right]. \quad (25)$$

The contribution to the real part of the self-energy to order ω^2 and V_{ds}^2 is proportional to $\sigma\tilde{\eta}(b)b$. It might be thought that such a term should cancel out in taking the sum over the two spin components. However, there is a σ -independent contribution from a cross term with the effective Zeeman term $\sigma b\eta(b)$, which has to be included. In the limit $b \rightarrow 0$ equation (21) reduces to (5).

The equation for the critical field b_c becomes

$$\frac{b^2}{\tilde{\Delta}^2(b)} = \frac{\sqrt{(3 - \alpha(b)\gamma(b))^2 + 4\gamma(b)(5 - \alpha(b))(1 + 5\gamma(b))} - 3 + \alpha(b)\gamma(b)}{2\gamma(b)(5 - \alpha(b))\tilde{\eta}(b)^2} \quad (26)$$

where $\alpha(b) = \alpha_{\omega}(b) + \alpha_V(b)$, and

$$\gamma(b) = \pi\tilde{\Delta}(b)\tilde{U}^2(b)[\tilde{\rho}_d^0(0, b)]^3 = \pi\tilde{\Delta}(b)\tilde{\rho}_d^0(0, b)(R(b) - 1)^2. \quad (27)$$

Equation (26) is an implicit equation for b_c which can be solved by iteration starting from the much simpler result (19), obtained within the linear approximation.

For a strong coupling situation ($U/\pi\tilde{\Delta} = 4$) the result for the critical field obtained by iterating equation (26) and using the b -dependent renormalized parameters taken from figure 1 is $b_c \simeq 0.459\tilde{\Delta} = 0.584T_K$. This differs only by 0.3% from the value obtained from (19). The small difference is due to the fact that the various correction terms due to the b dependence of the parameters in the more general formula (26) tend to cancel giving only a small resultant change.

Plots of the total differential conductance for various fields above and below the critical field are displayed in figure 4. We have taken the full self-energy as given in (21) into account, including the field dependence of the renormalized parameters. Our results are asymptotically exact only for small V_{ds} and a more complete theory is required to calculate the splitting at larger bias voltages. The major problem to be solved is the dependence of the self-energy on the voltage bias term, when eV_{ds} is comparable with and greater than the Kondo temperature T_K , so that a detailed comparison with experiment can be made with the experimental results in this regime.

IV. CONCLUSIONS

Measurements of the differential conductance in a magnetic field have been used to infer the magnitude of the magnetic splitting of a Kondo resonance [4, 5]. The results reveal an apparent disagreement with theoretical predictions, but the comparisons with theory have been based on calculations of the equilibrium Green's function for the dot. It is not clear that this will constitute a reasonable approximation at the finite bias voltages used in the experiment. To examine this question we have derived an expression for the differential conductivity through a quantum dot, described by a symmetric Anderson model, for small bias voltages and arbitrary magnetic field. This has enabled us to estimate the effect of the voltage dependent terms from the non-equilibrium dot Green's function. Our estimates

of the shifts of the component resonances for small magnetic field values differ significantly from the values obtained using the equilibrium Green's function. We conclude that it is important to take this voltage dependence into account and use the non-equilibrium Green's function for a meaningful comparison with experiment.

Though our calculations are restricted to the small voltage regime we have also been able to estimate the value of the critical field B_c for the emergence of two distinct peaks in the total differential conductance. Our approach is applicable in all ranges of the interaction U from the weak coupling to the strong coupling Kondo regime for $T = 0$. They complement the perturbation calculations in U of Fuji and Ueda [7] for the same model. It is difficult to make a comparison with their results, however, because the interaction simultaneously modifies the renormalized parameters, such as $\tilde{\eta}(b)$, and introduces a dependence on V_{ds} , both directly and through the ω dependence in the self-energy. These three effects of the interaction can be clearly distinguished within our formulation and taken separately into account. It should be possible to extend our calculations to larger values of V_{ds} using a renormalized version of the perturbation theory, and this approach is currently being investigated.

Acknowledgement

ACH wishes to thank the EPSRC for support through the Grant GR/S18571/01). JB thanks the Gottlieb Daimler- and Karl Benz-Foundation and EPSRC for financial support, and AO acknowledges the support by the Grant-in-Aid for Scientific Research for JSPS. We also wish to thank W. Koller for helpful discussions.

-
- [1] D. Goldhaber-Gordon, H. Shtrikman D. Mahalu J. Göres, M.A. Kastner and U. Meirav, Phys. Rev. Lett. **81**, 5225 (1998).
 - [2] T. A. Costi, A. C. Hewson and V. Zlatić, J. Phys.:Cond. Mat., **6**, 2519 (1994).
 - [3] D. Goldhaber-Gordon, H. Shtrikman, D. Mahalu, D. Abusch-Magder, U. Meirav and M.A. Kastner, Nature (London) **391**, 156 (1998).
 - [4] A. Kogan, S. Amasha, D. Goldhaber-Gordon, G. Granger, M.A. Kastner and H. Shtrikman, Phys. Rev. Lett. **93**, 166602 (2004).
 - [5] S. Amasha, I.J. Gelfand, M.A. Kastner and A. Kogan, cond-mat/0411485 (2004).
 - [6] Y. Meir, N.S. Wingreen and P.A. Lee, Phys. Rev. Lett. **70** 2601 (1993).
 - [7] T. Fuji and K. Ueda, J. Phys. Soc. Japan, **74**, 127 (2005).
 - [8] T.A. Costi, Phys. Rev. Lett. **85**, 1504 (2000).
 - [9] D.E. Logan and N.L. Dickens, J. Phys.:Cond. Mat., **13**, 9713 (2001).
 - [10] J. Moore and X.-G. Wen, Phys. Rev. Lett. **85**, 1722 (2000).
 - [11] W. Hofstetter, Phys. Rev. Lett. **85**, 1508 (2000).
 - [12] A. Rosch and T. A. Costi and J. Paaske and P. Wölfle, Phys. Rev. Lett., **90**, 076804-1 (2003).
 - [13] L.I. Glazman and M. Pustilnik, cond-mat/0501007 (2005)
 - [14] A. M. Clogston and P. W. Anderson, Bull. Am. Phys. Soc. **6**, 124 (1961).
 - [15] S. Hershfield and J. Davies and J. Wilkins, Phys. Rev. Lett., **67**, 3720 (1991).
 - [16] Y. Meir and N.S. Wingreen, Phys. Rev. Lett. **68**, 2512 (1992).
 - [17] N.S. Wingreen and Y. Meir, Phys. Rev. B **49**, 11040 (1994).
 - [18] A. Oguri, J. Phys. Soc. Japan, **74**, 110 (2005).
 - [19] A. Oguri, Phys. Rev. B, **64**, 153305 (2001).
 - [20] A.C. Hewson, Phys. Rev. Lett. **70**, 4007 (1993).
 - [21] A.C. Hewson, J. Phys.: Cond. Mat, **13**, 10011 (2001).
 - [22] A.C. Hewson, A. Oguri and D. Meyer, Euro. Phys. J. B **40**, 177 (2004).
 - [23] A.C. Hewson, J. Phys. Soc. Japan, **74**, 8 (2005).
 - [24] A.C. Hewson, J. Bauer and W. Koller, cond-mat/0508465 (2005).
 - [25] $\tilde{\eta}$ corresponds to the ratio $\tilde{\epsilon}_d(h)/h$ in reference [23].
 - [26] L.V. Keldysh, Sov. Phys. JETP **20**, 1018 (1965).
 - [27] K. Yamada, Prog. Theor. Phys. **53**, 970 (1975); Prog. Theor. Phys. **54**, 316 (1975).