

Exact results for the 1D interacting Fermi gas with arbitrary polarization

M .T .B atchelor^{1,2}, M .B orts¹, X .W .G uan^{1,2} and N .O elkers^{1,2}

¹ D epartm ent of T heoretical P hysics, R esearch S chool of P hysical S ciences and E ngineering

² D epartm ent of M athem atics, M athem atical S ciences I nstitute
T he A ustralian N ational U niversity, C anberra A C T 0200, A ustralia

A bstract. W e investigate the 1D interacting two-component Fermi gas with arbitrary polarization. Exact results for the ground state energy, quasim omentum distribution functions, spin velocity and charge velocity reveal subtle polarization dependent quantum effects.

P A C S num bers: 03.75.Ss, 05.30.Fk, 71.10.Pm

Recent achievements in trapping quantum gases of ultra-cold atoms have opened up many exciting possibilities for the experimental investigation of quantum effects in low-dimensional many-body systems. Among these, progress in realizing interacting Fermi gases promises new opportunities for studying the pronounced crossover from a Bardeen-Cooper-Schrieffer (BCS) superfluid to a Bose-Einstein condensate (BEC) [1]. A crossover from weakly attractive Cooper pairs to strongly repulsive molecular dimers can also take place in a confined quasi 1D Fermi gas of atoms by using magnetic field induced Feshbach resonances [2, 3]. In this scenario, a number of key 1D integrable models with a variable interaction parameter have been used to describe the crossover [3, 4, 5, 6, 7, 8, 9]. These include the integrable interacting Fermi gas [10].

The 1D Fermi gas can be experimentally realized [2] by tightly confining the atomic cloud in the radial directions and weakly confining it along the axial direction [3, 4, 5, 6, 11]. To date, the theoretical study of the 1D continuum Fermi gas has been mainly for the zero polarization case in the thermodynamic limit (TL). Our aim here is to present the results of a comprehensive study of the 1D exactly solved two-component Fermi gas with arbitrary polarization, throughout the whole interaction range. This includes finite systems in the weak and strong coupling regions, and in the TL.

The model. The Hamiltonian [10]

$$H = \sum_{i=1}^N \frac{\hbar^2 k_i^2}{2m} + g_{1D} \sum_{1 \leq i < j \leq N} \delta(x_i - x_j) \quad (1)$$

describes a δ -function interacting gas of N fermions each of mass m constrained by periodic boundary conditions to a line of length L . The coupling constant g_{1D} can be written in terms of the scattering strength $c = 2/a_{1D}$ as $g_{1D} = \hbar^2 c/m$. An effective 1D scattering length a_{1D} can be expressed through the 3D scattering length for fermions confined in a 1D geometry. We use the dimensionless coupling constant $\gamma = m g_{1D} L / (\hbar^2 n)$ for physical analysis. Here $n = N/L$ is the linear density. The coupling g_{1D} can vary from $-\infty$ to ∞ by detuning the 3D scattering length. The interaction is attractive for $g_{1D} < 0$ and repulsive for $g_{1D} > 0$.

The energy is given by $E = \frac{\hbar^2}{2m} \sum_{j=1}^N k_j^2$, where the quasimomenta k_j satisfy the Bethe equations (BE) [10]

$$\exp(ik_j L) = \prod_{i=1}^M \frac{k_j - k_i + ic}{k_j - k_i - ic};$$

$$\prod_{i=1}^N \frac{k_i + ic}{k_i - ic} = \prod_{i=1}^M \frac{k_i + ic}{ic}; \quad (2)$$

Here $j = 1; \dots; N$ and $i = 1; \dots; M$, with M the number of spin-down fermions. For convenience we take N even. Of particular interest are the bound states for attractive interaction. In this regime, the model (1) has been extensively studied with $N = 2M$ in the context of the BCS-BEC crossover [3, 4, 5, 6, 7]. For weak attraction, the system describes weakly bound Cooper pairs. In the strongly attractive regime the bound states behave like bosonic tightly bound molecular dimers. On the other hand, for repulsive

interaction, there are no bound states for the ground state. The spin-up and spin-down fermions behave like indistinguishable free fermions as $c \rightarrow 1$. The ground state energy per particle in the TL, i.e., $N/M; L \rightarrow 1$, $N=L; M=L$ constant, can be derived from the Gaudin integral equations [10]. Here we first analyze the BE directly, i.e., for a finite system.

The ground state. We observe that for weak attractive interaction a pair of fermions with spin-up and spin-down states forms a bound state. On the other hand, for weak repulsive interaction the fermion momenta separate in quasimomentum space. For $L/c \gg 1$, it is convenient to distinguish between unpaired $k_j^{(u)}$ ($j = 1; \dots; N-2M$) and paired $k_j^{(p)}$ ($j = 1; \dots; 2M$) momenta. On expanding the BE in powers including $O(c)$ [12], one obtains $k_j^{(u)} = (M-N-1+2j)/L + \frac{(u)}{j}$; $j = 1; \dots; N-2M$, $k_j^{(u)} = (3M-N-1+2j)/L + \frac{(u)}{j}$; $j = N-2M+1; \dots; N-2M$, and $k_j^{(p)} = (1-M+2j)/L + \frac{(p)}{j_+}$ $\frac{1}{c=L}$, ($j = 1; \dots; M$), where $j_+ = j$, if j odd and $j_+ = j-1$ if j even. Therefrom it follows that the binding energy in the weakly attractive case is $c=L$ for finite systems. The deviations from $k_j^{(u,p)}$ are linear in c , with

$$k_j^{(u)} = \frac{c}{L} \sum_{\substack{\text{prime} \\ \text{sum}}} \frac{1}{k_{j;0}^{(u)} k_{j;0}^{(p)}}; \quad (3)$$

$$k_j^{(p)} = \frac{c}{L} \sum_{\substack{\text{prime} \\ \text{sum}}} \frac{1}{k_{j;0}^{(p)} k_{j;0}^{(u)}} + \frac{1}{2} \sum_{\substack{\text{prime} \\ \text{sum}}} \frac{1}{k_{j;0}^{(p)} k_{j;0}^{(u)}}; \quad (4)$$

where the primed sum counts each $k_{j;0}^{(p)}$ only once and excludes $k_{j;0}^{(p)}$. Here $k_{j;0}^{(u,p)}$ are the quasimomenta of non-interacting particles as given above. Note that the unpaired momenta do not interact with each other, which is consistent with the Pauli principle. On the other hand, for strong attractive interaction, i.e., $L/c \rightarrow 1$, tightly bound states with quasimomentum $k_i^{(p)} \rightarrow i \frac{1}{2} \pi$ separate from unbound states with quasimomentum $k_j^{(u)} = \frac{n_j}{L} - 1 + \frac{4M}{Lc} - 1$ with integers $n_j = 1; 3; \dots; (N-2M-1)$. In the above, $i = \frac{n_i}{L} - 1 + \frac{M}{Lc} + \frac{2(N-2M)}{Lc} - 1$ for $n_i = 1; \dots; \frac{1}{2}(M-1)$. In this scenario the bound states behave like hard-core bosons due to Fermi pressure.

We now define the fraction $a = 2M/N$ which characterizes the polarization of the system. The ground state energy per particle in the weakly interacting regime follows from the above asymptotic roots of the BE as (in units of $\hbar^2 n^2/(2m)$), with $a_0 = (1-a)(1 - \frac{1}{2}a)^2 + \frac{1}{2}a$

$$\frac{E}{N} = \frac{a^2}{2} + (1-a)a + \frac{a^3}{12} + \frac{a^2}{3}a_0 + O(c^{-2}); \quad (5)$$

The first two terms represent the binding and the interaction energy, respectively, which are both linear in a . This is different from what is obtained in the TL from the Gaudin integral equations [4, 6, 7], where the binding energy is $\propto a^2$. The other terms in (5) are the kinetic energies for paired and unpaired fermions. In the special case $a = 0$,

The polarization dependence of atom-hole condensation in optical lattices was discussed in Ref. [13].

the model reduces to fully polarized free fermions without s-wave scattering. Using the given asymptotic expressions for the Bethe roots in the strongly attractive regime, we obtain (in units of $\hbar^2 n^2 / (2m)$)

$$\frac{E}{N} = \frac{a}{4} + \frac{a^3}{3} (1 - 2b) + a_1 (1 - 2b) + O(a^2); \quad (6)$$

where $a_1 = (1 - a)^3$, $b_1 = a(2 - a) + 2(1 - a)a$ and $b = 2a$. Whereas the first term represents the binding energy, the remaining interaction energies stem from the interaction between pair-pair and pair-unpaired fermions. The ground state energy is lowest for $a = 1$, compared to the others for $0 < a < 1$. This is in general agreement with the Lieb-Mattis theorem [14].

Now consider the crossover to the TL. First we observe that in the TL, the results (5), (6) reduce to those derived from the Gaudin integrodifferential equations [4, 5, 6, 7]. Furthermore, from (3), (4), we obtain the quasimomentum distribution functions $n_{u,p}(k)$ per unit length in the TL,

$$n_u(k) = \frac{1}{2} + \frac{c}{2} \frac{A}{k^2} \frac{1}{A^2}; \quad A < |k| < B; \quad (7)$$

$$n_p(k) = \frac{1}{2} - \frac{c}{2} \frac{A}{k^2} \frac{1}{A^2} + \frac{B}{B^2 - k^2}; \quad |k| < A; \quad (8)$$

with $A = M/L$ and $B = (N + M)/L$. Especially, for $M = N = 2$, $n_u = 0$. In Fig. 1, $n_{u,p}(k)$ as calculated from the finite BE are compared to Eqs. (7), (8).

We now use the Gaudin integrodifferential equations to numerically compute the dependence of E/N on a for different a . The result is shown in Fig. 2. The numerical data agree with (5) and (6) in the weakly interacting and strongly attractive regimes. A straightforward analysis of the integrodifferential equations in the strongly repulsive regime leads to

$$\frac{E}{N} < \frac{\hbar^2 n^2}{2m} \frac{3}{4} (1 - \frac{4a}{3}) + O(a^2); \quad a < 1$$

$$: \frac{\hbar^2 n^2}{2m} \frac{3}{4} (1 - \frac{4 \ln 2}{3}) + O(a^2); \quad a = 1 \quad (9)$$

which provides a further check on the numerics. As can be seen from (9), for strongly repulsive interaction the mixture behaves like a single component Fermi gas of indistinguishable particles, rather than hard-core bosons [4, 5, 7].

Spin-charge separation. In the TL, the low-energy behaviour of interacting Bose and Fermi gases is universal in the dispersion relation and correlation behaviour [15]. Spin-charge separation is one of the typical Luttinger liquid behaviours, with both the spin and charge degrees of freedom having different propagation velocities at Fermi surfaces. The charge-velocity can be calculated exactly from $v_c^2 = \frac{1}{m} \frac{\partial^2 E}{\partial L^2}$ [16] for arbitrary a , which yields

$$\frac{v_c}{v_F} = \frac{q}{q} \frac{\frac{a^3}{16} (1 - 4b) + a_1 (1 - 4b)}{a(1 - \frac{a-2}{2}) + \frac{a^3}{4} + a_0}; \quad |j| < 1; \quad (10)$$

$$: \frac{a(1 - \frac{a-2}{2}) + \frac{a^3}{4} + a_0}{a(1 - \frac{a-2}{2}) + \frac{a^3}{4} + a_0}; \quad |j| > 1;$$

where the Fermi velocity $v_F = \hbar^2 n / m$.




Figure 1. The quasimomentum distribution function (crosses) for a finite system ($N = 50; M = 16$) for the a) weakly repulsive and b) weakly attractive regime. The full lines connecting the crosses are guides to the eye. The insets are zooms of the distribution near the noninteracting cases (dotted lines), $1/2$ for unpaired fermions and $1/4$ for paired fermions. Here the full lines are the results obtained in the TL (Eqs. (7), (8)).

To illustrate spin-charge separation for non-zero interaction, we show the spin and charge velocities in Fig. 3 as a function of j for $a = 1$. These curves are obtained numerically from the dressed-energy formalism [17, 18]. For weak interaction [6],

$$\frac{v_{c/s}}{v_F} = \frac{1}{2} \left(1 \pm \frac{1}{2} \right); \quad j \gg 1; \quad (11)$$

where the upper sign refers to v_c . Furthermore, $\frac{v_c}{v_F} = 1 - \frac{4 \ln 2}{3}$, $\frac{v_s}{v_F} = \frac{2}{3}$ for $j \rightarrow 1$ and $\frac{v_c}{v_F} = \frac{1}{4} (1 \pm 1)$ for $j \rightarrow 1/6$. In the attractive regime, the charge excitations are still gapless; however, a gap opens in the spin channel [19]. This gap increases with increasing j , leading to a divergent spin velocity in the strongly attractive limit, with $\frac{v_s}{v_F} = \frac{1}{2} \frac{1}{1 - 2j}$ as $j \rightarrow 1/6$. Some possible ways to experimentally observe the




Figure 2. Ground state energy in units of $\tilde{\mu}^2 n^2 = (2m)$ vs interaction strength for different polarizations. Dashed lines compare the analytic approximations (5), (6), (9) with the numerical solution of the integralequations (solid lines) in the different coupling regimes. The curves shown are for $a = 0, 0.2, 0.4, 0.6, 0.8, 1$ (top to bottom).

spin-charge separation in harmonically trapped atoms have been suggested [11].

In conclusion, we have examined in detail the exactly solved two-component 1D interacting Fermi gas for arbitrary polarization a in the whole interaction range.




Figure 3. Spin and charge velocities vs interaction strength for $a = 1$. The straight dotted lines show the lowest order in γ from (11), which is also accessible by field theory [11].

Analytic results have followed for the quasimomentum density profile, the ground state energy and the spin-charge velocities. These quantities reveal significant insight into the nature of quantum effects in the Fermi gas with arbitrary polarization.

Acknowledgments

This work has been supported by the Australian Research Council and the German Science Foundation (grant number BO/2538). The authors also thank the organizers of the Counting Complexity meeting in honour of Tony Guttmann's 60th birthday, where this work was presented.

References

- [1] Jochim S et al 2003 *Science* 302 2101
- Chin C et al 2004 *Science* 305 1128
- Zwierlein M W et al 2004 *Phys. Rev. Lett.* 92 120403
- Regal C A, Greiner M and Jin D S 2004 *Phys. Rev. Lett.* 92 040403
- Bartenstein M et al 2004 *Phys. Rev. Lett.* 92 120401
- Kinast J et al 2005 *Science* 307 1296
- [2] Moritz H et al 2005 *Phys. Rev. Lett.* 94 210401
- [3] Recati A, Fuchs J N and Zwirger W 2005 *Phys. Rev. A* 71 033630
- [4] Astrakharchik G E, Blume D, Giorgini S and Pitaevskii L P 2004 *Phys. Rev. Lett.* 93 050402
- [5] Tokatly I V 2004 *Phys. Rev. Lett.* 93 090405
- [6] Fuchs J N, Recati A and Zwirger W 2004 *Phys. Rev. Lett.* 93 090408
- [7] Iida T Wadati M 2005 *J. Phys. Soc. Jpn.* 74 1724
- [8] Liu X-J, Drummond P D and Hu H 2005 *Phys. Rev. Lett.* 94 136406
- [9] Ortiz G and Dukelsky J 2005 *Phys. Rev. A* 72 043611
- [10] Gaudin M 1967 *Phys. Lett. A* 24 55
- Yang C N 1967 *Phys. Rev. Lett.* 19 1312
- [11] Recati A, Fedichev P O, Zwirger W and Zoller P 2003 *J. Opt. B* 5 S55
- [12] Batchelor M T, Guan X W and McGuire J B 2004 *J. Phys. A: Math. Gen.* 37 L497
- Batchelor M T, Guan X W, Dunning C and Links J 2005 *J. Phys. Soc. Jpn.* 74 S 53
- [13] Lee C 2004 *Phys. Rev. Lett.* 93 120406
- [14] Lieb E and Mattis D 1962 *Phys. Rev.* 125 164
- [15] Haldane F D M 1981 *Phys. Rev. Lett.* 47 1840
- [16] London F 1964 *Superfluids Vol. II* (New York, Dover)
- [17] Takahashi M 1999 *Thermodynamics of One-Dimensional Solvable Models* (Cambridge, Cambridge University Press).
- [18] Schlottmann P 1994 *J. Phys.: Cond. Mat.* 6 1359
- [19] Krivnov V Ya and Ovchinnikov A A 1975 *Sov. Phys. JETP* 40 78
- Musurkin I A and Ovchinnikov A A 1971 *Sov. Phys. Sol. St.* 12 2031