

Dependence of the superfluidity criterion on the resonance between one-particle (Bogoliubov) and two-particle series

V. P. Maslov

Abstract

We study how the superfluidity depends on the resonance between one- and two-particle series. The frequency of the spectrum of two-particle solutions in an interval is calculated.

We consider a system of N identical bosons of mass m that lie on a three-dimensional torus \mathbf{T} with the side lengths L_1 , L_2 , and L_3 . We assume that the bosons mutually interact and that the interaction potential has the form

$$V(N^{1/3}(x-y)), \quad (1)$$

where $V(\xi)$ is an even smooth function rapidly decreasing at infinity and x, y are the boson coordinates on \mathbf{T} . The interaction potential (1) depends on N such that the interaction radius decreases as the particle number N increases, and the average number of particles with which a single particle interacts remains constant in this case.

In the ultrasecond quantization over pairs, an ultrasecond-quantized operator $\widehat{\overline{H}}$, whose explicit form was previously presented, corresponds to the boson system in question. As stated, this ultrasecond-quantized operator satisfies identity

$$\widehat{\overline{H}} = \widehat{\overline{E}}\widehat{A}, \quad (2)$$

where $\widehat{\overline{E}}$ is the ultrasecond-quantized unit operator and \widehat{A} is an operator nonuniquely chosen in the ultrasecond quantization space. Further, we assume that the operator \widehat{A} has the form

$$\begin{aligned} \widehat{A} = & \iint dxdy \widehat{B}^+(x, y) \left(-\frac{\hbar^2}{2m}(\Delta_x + \Delta_y) + V(N^{1/3}(x-y)) \right) \widehat{B}^-(x, y) + \\ & + 2 \int dxdydx'dy' V(N^{1/3}(x-y)) \widehat{B}^+(x, y) \widehat{B}^+(x', y') \widehat{B}^-(x, x') \widehat{B}^-(y, y'), \end{aligned} \quad (3)$$

where $\widehat{B}^+(x, y)$ and $\widehat{B}^-(x, y)$ are the corresponding bosonic creation and annihilation operators for a particle pair in the Fock ultrasecond quantization space. By identity (2), an asymptotic expression for operator (3) can be found for establishing an asymptotic representation for the spectrum of the boson system under consideration in the limit as $N \rightarrow \infty$.

Because the function (1) times N in a sense converges to the Dirac function in the limit as $N \rightarrow \infty$, the operator (3) involves the small parameter $1/N$ in the second term for

this limiting case. This means that to find the asymptotic expressions for eigenfunctions and eigenvalues of the operator \hat{A} , we can use the semiclassical methods developed in [8]. The asymptotic expressions for eigenfunctions and eigenvalues are determined by the symbol of the operator (3), and this symbol is called the true symbol for the ultrasecond-quantized problem. The true symbol corresponding to the operator (3) is the following functional defined for a pair of functions $\Phi^+(x, y)$ and $\Phi(x, y)$:

$$\begin{aligned} \mathcal{H}[\Phi^+(\cdot), \Phi(\cdot)] = & \iint dx dy \Phi^+(x, y) \left(-\frac{\hbar^2}{2m}(\Delta_x + \Delta_y) \right) \Phi(x, y) + \\ & + 2 \int dx dy dx' dy' (NV (N^{1/3}(x - y))) \Phi^+(x, y) \Phi^+(x', y') \Phi(x, x') \Phi(y, y'). \end{aligned} \quad (4)$$

The conservation of the number of particles in the system for the functions $\Phi^+(x, y)$ and $\Phi(x, y)$ implies the condition

$$\iint dx dy \Phi^+(x, y) \Phi(x, y) = \frac{1}{2}. \quad (5)$$

According to the asymptotic methods in [8], in the limit as $N \rightarrow \infty$, an asymptotic series of eigenfunctions and eigenvalues of the operator (3) corresponds to each solution of the equations

$$\Omega \Phi(x, y) = \frac{\delta \mathcal{H}}{\delta \Phi^+(x, y)}, \quad \Omega \Phi^+(x, y) = \frac{\delta \mathcal{H}}{\delta \Phi(x, y)} \quad (6)$$

that also satisfy condition (5). It follows from the explicit form of the true symbol (4) that system (6) can be written as

$$\begin{aligned} \Omega \Phi(x, y) = & -\frac{\hbar^2}{2m}(\Delta_x + \Delta_y) \Phi(x, y) + \\ & + \iint dx' dy' (NV (N^{1/3}(x - y)) + NV (N^{1/3}(x' - y'))) \Phi^+(x', y') \Phi(x, x') \Phi(y, y'), \\ \Omega \Phi^+(x, y) = & -\frac{\hbar^2}{2m}(\Delta_x + \Delta_y) \Phi^+(x, y) + \\ & + \iint dx' dy' (NV (N^{1/3}(x - x')) + NV (N^{1/3}(y - y'))) \Phi(x', y') \Phi^+(x, x') \Phi^+(y, y') \end{aligned} \quad (7)$$

In the limit as $N \rightarrow \infty$, system (7) supplemented with the condition (5) has the family of solutions

$$\begin{aligned} \Phi_k^+(x, y) = & \frac{1}{L_1 L_2^2} \cos(k(x - y)), \\ \Phi_k(x, y) = & \frac{1}{L_1 L_2^2} \sum_l \varphi_{k,l} \exp(il(x - y)), \end{aligned} \quad (8)$$

where k and l are three-dimensional vectors of the form

$$2\pi \left(\frac{n_1}{L_1}, \frac{n_2}{L_2}, \frac{n_3}{L_2} \right),$$

the numbers n_1 , n_2 , and n_3 are integers, the functions $\varphi_{k,l}$ in formula (2) have the form

$$\varphi_{k,l} = \frac{1}{2V_0} \left(\frac{\hbar^2}{2m}(k^2 - l^2) + V_0 \pm \sqrt{\left(\frac{\hbar^2}{2m}(k^2 - l^2) + V_0 \right)^2 - V_0^2} \right), \quad (9)$$

with the plus sign for $l^2 > k^2$ and the minus sign for $l^2 < k^2$, and V_0 here and hereafter denotes the expression

$$V_0 = \frac{1}{L_1 L_2^2} \int dx V(x), \quad (10)$$

in which the integral is taken over the space \mathbf{R}^3 . The vector k in (8) plays the role of a parameter indexing different solutions of system (7) with conditions (5). The solutions (8) are standing waves associated with the series without fluidity.

The leading asymptotic term for the eigenvalues in the series corresponding to solution (8) is equal to the value of the product of the symbol (4) on the functions (8) times N ,

$$E_k = N \left(\frac{\hbar^2 k^2}{2m} + \frac{V_0}{2} \right). \quad (11)$$

The asymptotic expressions for the eigenvalues and eigenfunctions, in particular, the asymptotic terms following (11) are determined by the solutions of system (7) and also by the solutions of the variational system of equations corresponding to (7). The detailed study of solutions of the variational system is here omitted. In what follows, we present the results of this study obtained, in particular, by using the Maple program.

If $k = 0$, then the asymptotic series related to this solution (8) of system (7), (5) is the Bogoliubov series corresponding to the ground state without fluidity. The quasiparticle spectrum for this series is expressed by the well-known formula

$$\lambda_l = \sqrt{\left(\frac{\hbar^2 l^2}{2m} + V_0 \right)^2 - V_0^2}. \quad (12)$$

If $k \neq 0$, then the variational system equations corresponding to (7) gives the quasiparticle spectrum

$$\begin{aligned} \lambda_{1,k,l} &= \\ &= \pm \frac{\hbar^2}{2m} \sqrt{k^4 + \frac{l^2}{2} + \frac{l_1^2}{2} - k^2 l^2 - k^2 l_1^2 + \frac{1}{2}(l^2 + l_1^2 - 2k^2) \sqrt{(l^2 - l_1^2)^2 + \left(\frac{4mV_0}{\hbar^2} \right)^2}}, \\ \lambda_{2,k,l} &= \\ &= \pm \frac{\hbar^2}{2m} \sqrt{k^4 + \frac{l^2}{2} + \frac{l_1^2}{2} - k^2 l^2 - k^2 l_1^2 - \frac{1}{2}(l^2 + l_1^2 - 2k^2) \sqrt{(l^2 - l_1^2)^2 + \left(\frac{4mV_0}{\hbar^2} \right)^2}}, \end{aligned} \quad (13)$$

where $l_1 = l + 2k$ and $l \neq k$. In the formula for $\lambda_{1,k,l}$ the plus sign is chosen for $l^2 > k^2$ and the minus sign for $l^2 < k^2$; in the formula for $\lambda_{2,k,l}$, the plus sign is chosen for $l_1^2 > k^2$ and the minus sign for $l_1^2 < k^2$.

The explicit form (13) implies that this quasiparticle spectrum contains some negative values, which means that the series corresponding to the solution (8) with $k \neq 0$ is non-metastable. In this connection, we suggest the following explanation for the dependence of the critical velocity on capillary width. In what follows, we assume that $L_1 \gg L_2$. We consider the Bogoliubov series corresponding to the flow with the velocity $\hbar k_0/m$ along the capillary, where $k_0 = 2\pi(n_1/L_1, 0, 0)$. For the boson system in question, the leading asymptotic term for the eigenvalues in this series is

$$N \left(\frac{\hbar^2 k_0^2}{2m} + \frac{V_0}{2} \right). \quad (14)$$

We now assume that there is a relation between L_1 and L_2 such that there exists a vector $k = 2\pi(0, n_2/L_2, n_3/L_2)$ for which the corresponding value of the term in (11) is exactly equal to (14). This means that there can be resonance between the current states of the Bogoliubov series and the states of the nonmetastable series corresponding to (8). If the value of L_1 is very large, then resonance is also possible for the case where the value of the term in (11) is close to expression (14) but need not coincide with it. The existence of such a resonance indicates the possibility of a transition from the current state to a nonmetastable state from which the system falls to the lowest energy level, which indicates the loss of superfluidity.

The following versions of series of two-particle solutions are determined by the relation:

$$Q(k_1, k_2, m_1, m_2, m_3, l_1, l_2, l_3) = \frac{l_3^2 - m_3^2}{m_1^2 + m_2^2 + k_1^2 + k_2^2 - l_1^2 - l_2^2}, \quad (15)$$

where $k_1, k_2, l_1, l_2, l_3, m_1, m_2, m_3$ are integers. We impose the following conditions on the numbers $k_1, k_2, l_1, l_2, l_3, m_1, m_2, m_3$:

$$Q > 0, \quad k_1^2 + k_2^2 > 0, \quad (16)$$

where Q is an irreducible fraction, i.e., its numerator and denominator do not have common divisors.

Professor A. A. Karatsuba kindly calculated for me the frequency of the spectrum of two-particle solutions in the interval $[a, a + h]$. The asymptotic expression in the limit as $L \rightarrow \infty$ has the following form.

We are given the functions (15) and the function

$$V = \sqrt{l_1^2 + l_2^2 + \frac{l_3^2}{Q}}, \quad (17)$$

where $k_1, k_2, m_1, m_2, m_3, l_1, l_2, l_3$ are nonnegative integers that do not exceed L ; moreover, $Q > 0$ and $k_1^2 + k_2^2 > 0$.

We are also given the numbers $a \geq 1, 0 < h \leq 1$.

Let K be the number of sets $k_1, k_2, m_1, m_2, m_3, l_1, l_2, l_3$ satisfying the inequality

$$a < V \leq a + h. \quad (18)$$

Then K is determined by the approximate formula

$$K = \left(\frac{1}{5} + \frac{1}{9}\right) \frac{\pi^3}{128} L^2 ((a + h)^6 - a^6) = \frac{7\pi^3}{2880} L^2 ((a + h)^6 - a^6). \quad (19)$$

If the distance between the spectrum points is of the order of $\frac{1}{N}$, i.e., if it is determined with the desired accuracy, then the Bogoliubov series is in resonance.

This readily shows that $a \sim 4$, and hence resonance occurs approximately at the points $8\pi h/mL_2$. This resonance is braking the fluid, and the superfluidity disappears.

We here present A. A. Karatsuba's proof of formula (19).

We introduce the new notation

$$r = m_1^2 + m_2^2 + k_1^2 + k_2^2, \quad l = l_1^2 + l_2^2, \quad \xi = \frac{m_3^2}{l_3^2} \geq 0.$$

In this notation, we have

$$Q = l_3^2 \frac{1 - \xi^2}{r - l} > 0.$$

Two cases, (I) and (II), are possible:

$$(I) \quad 0 \leq \xi < 1, \quad l < r; \quad (II) \quad \xi > 1, \quad l > r.$$

The function V can be rewritten as

$$V = \sqrt{l + \frac{r-l}{1-\xi^2}} = \sqrt{\frac{r-l\xi^2}{1-\xi^2}}$$

and hence (19) becomes

$$a^2 < V^2 = \frac{r-l\xi^2}{1-\xi^2} \leq (a+h)^2. \quad (20)$$

Case (I). From (20) we obtain

$$\begin{aligned} a^2(1-\xi^2) + l\xi^2 &< r \leq (a+h)^2(1-\xi^2) + l\xi^2, \\ l &< r, \quad 0 \leq \xi < 1. \end{aligned} \quad (21)$$

If in (21) we have the relation

$$a^2(1-\xi^2) + l\xi^2 \leq l, \quad \text{i.e.,} \quad a^2 \leq l,$$

then system (21) is equivalent to

$$\begin{aligned} l &< r \leq (a+h)^2(1-\xi^2) + l\xi^2, \\ a^2 &\leq l < (a+h)^2, \\ 0 &\leq \xi < 1. \end{aligned} \quad (22)$$

But if in (21) we have

$$a^2(1-\xi^2) + l\xi^2 > l, \quad \text{i.e.,} \quad l < a^2,$$

then system (21) is equivalent to

$$\begin{aligned} a^2(1-\xi^2) + l\xi^2 &< r \leq (a+h)(1-\xi^2) + l\xi^2, \\ l &< a^2, \\ 0 &\leq \xi < 1. \end{aligned} \quad (23)$$

Let K_1 be the number of sets satisfying either (22) or (23).

It follows from systems (22) and (23) that

$$l < a^2 \quad \text{or} \quad l < (a+h)^2,$$

i.e., we always have $l_1, l_2 < a+h \leq a+1$, since $h \leq 1$.

Precisely in the same way we obtain

$$r < (a+h)^2, \quad \text{i.e.,} \quad m_1, m_2, k_1, k_2 < a+h \leq a+1.$$

Therefore, the six variables $l_1, l_2, m_1, m_2, k_1, k_2$ are bounded by the value $a+1$, and they play an unimportant role for small a . The main parameter is L , $L \rightarrow +\infty$, and this parameter is related only to the variable ξ ,

$$\xi = \frac{m_3}{l_3} < 1, \quad 0 \leq m_3 < l_3 \leq L.$$

The ranges of these variables are determined automatically by the first two inequalities in systems (22) and (23). In what follows, we present two asymptotic formulas (in the limit as $x \rightarrow \infty$).

The number of sets m_1, m_2, k_1, k_2 for which $r = m_1^2 + m_2^2 + k_1^2 + k_2^2 \leq x$ is equal to

$$\frac{1}{24}v_4(\sqrt{x}),$$

where $v_4(\sqrt{x})$ is the volume of a four-dimensional sphere of radius \sqrt{x} , i.e.,

$$v_4(\sqrt{x}) = (\sqrt{x})^4 \frac{\pi^{4/2}}{\Gamma(4/2 + 1)} = x^2 \frac{\pi^2}{2}; \quad (24)$$

the number of sets l_1, l_2 for which $l_1^2 + l_2^2 \leq x$ is equal to

$$\frac{1}{4}\pi x. \quad (25)$$

These are approximate formulas; the larger x , the more precise they are.

From (24) and (22), (23) we obtain

$$\begin{aligned} K_1 = & \sum_{0 \leq \xi < 1} \sum'_{a^2 \leq l < (a+h)^2} \frac{\pi^2}{32} \{((a+h)^2(1-\xi^2) + l\xi^2)^2 - (l)^2\} \\ & + \sum_{0 \leq \xi < 1} \sum'_{l < a^2} \frac{\pi^2}{32} \{((a+h)^2(1-\xi^2) + l\xi^2)^2 - (a^2(1-\xi^2) + l\xi^2)^2\}. \end{aligned}$$

The prime on the sum over l means that the numbers l are regarded with multiplicity taken into account (the number of solutions of the equation is $l = l_1^2 + l_2^2$, $l_1 \geq 0$, $l_2 \geq 0$); the multiplicity is equal to $\pi/4$ (averaged over l), i.e.,

$$\begin{aligned} K_1 = & \sum_{0 \leq \xi < 1} \sum_{a^2 \leq l < (a+h)^2} \frac{\pi^2}{128} \{((a+h)^2(1-\xi^2) + l\xi^2)^2 - l^2\} \\ & + \sum_{0 \leq \xi < 1} \sum_{l < a^2} \frac{\pi^2}{128} \{((a+h)^2(1-\xi^2) + l\xi^2)^2 - (a^2(1-\xi^2) + l\xi^2)^2\} \end{aligned}$$

The sum over l (with a good accuracy) is equal to the integral

$$\begin{aligned} K_1 = & \frac{\pi^3}{128} \sum_{0 \leq \xi < 1} \int_{a^2}^{(a+h)^2} \{((a+h)^2(1-\xi^2) + l\xi^2)^2 - l^2\} dl \\ & + \frac{\pi^3}{128} \sum_{0 \leq \xi < 1} \int_0^{a^2} \{((a+h)^2(1-\xi^2) + l\xi^2)^2 - (a^2(1-\xi^2) + l\xi^2)^2\} dl \\ = & \frac{\pi^3}{128} ((a+h)^6 - a^6) \sum_{0 \leq \xi < 1} \left(\frac{2}{3} - \xi^2 + \frac{1}{3}\xi^4 \right). \end{aligned}$$

Since

$$\xi = \frac{m_3}{l_3}, \quad 0 \leq m_3 < l_3 \leq L,$$

the sum over ξ (with a good accuracy) is equal to the integral

$$\int_0^L \int_0^l \left(\frac{2}{3} - \left(\frac{m}{l} \right)^2 + \frac{1}{3} \left(\frac{m}{l} \right)^4 \right) dm dl = \frac{L^2}{2} \left(\frac{2}{3} - \frac{1}{3} + \frac{1}{15} \right) = \frac{L^2}{5}.$$

Hence we obtain

$$K_1 = \frac{\pi^3}{128} ((a+h)^6 - a^6) \frac{L^2}{5}.$$

Case (II).

$$\xi > 1, \quad l > r, \quad \text{i.e.,} \quad l_3 < m_2, \quad r < l.$$

Let

$$0 < \xi_1 = \frac{1}{\xi} = \frac{l_3}{m_3} < 1, \quad r < l.$$

Then we have the equality

$$V = \sqrt{\frac{r - l\xi^2}{1 - \xi^2}} = \sqrt{\frac{l - r\xi_1^2}{1 - \xi_1^2}}$$

and the condition $a < V \leq (a+h)$ implies

$$a^2 < \frac{l - r\xi_1^2}{1 - \xi_1^2} \leq (a+h)^2$$

or

$$\begin{aligned} a^2(1 - \xi_1^2) + r\xi_1^2 &< l \leq (a+h)^2(1 - \xi_1^2) + r\xi_1^2, \\ r < l, \quad 0 < \xi_1 < 1. \end{aligned} \tag{26}$$

It follows from (26) that $r < (a+h)^2$. If $r > a^2(1 - \xi_1^2) + r\xi_1^2$, i.e., if $r > a^2$, then (26) is replaced by the system

$$\begin{aligned} r - l < l &\leq (a+h)^2(1 - \xi_1^2) + r\xi_1^2, \\ a^2 < r < (a+h)^2, \quad 0 < \xi_1 < 1. \end{aligned} \tag{27}$$

If $r \leq a^2(1 - \xi_1^2) + r\xi_1^2$, i.e., if $r \leq a^2$, then (26) is replaced by (28):

$$\begin{aligned} a^2(1 - \xi_1^2) + r\xi_1^2 &< l \leq (a+h)^2(1 - \xi_1^2) + r\xi_1^2, \\ r &\leq a^2, \quad 0 < \xi_1 < 1. \end{aligned} \tag{28}$$

It follows from Case (I) and systems (22) and (23) that ξ is replaced by ξ_1 , while r and l interchange.

Therefore, we use formulas (25), (27), and (28) to find K_2 , which is the number of sets satisfying (27) and (28):

$$\begin{aligned} K_2 &= \frac{1}{4}\pi \sum_{0 < \xi_1 < 1} \sum'_{a^2 < r < (a+h)^2} ((a+h)^2(1 - \xi_1^2) + r\xi_1^2 - r) \\ &\quad + \frac{1}{4}\pi \sum_{0 < \xi_1 < 1} \sum'_{r \leq a^2} ((a+h)^2 - a^2)(1 - \xi_1^2). \end{aligned}$$

The prime on the sum over r means that the number of solutions r is regarded with multiplicity taken into account (the number of solutions of the equation is $r = m_1^2 + m_2^2 + k_1^2 + k_2^2$).

The number of solutions of the inequality

$$m_1^2 + m_2^2 + k_1^2 + k_2^2 \leq x, \quad m_1, m_2, k_1, k_2 \geq 0, .$$

is approximately equal to

$$\frac{\pi^2}{32}x^2.$$

We obtain

$$\begin{aligned} \sum'_{a^2 < r < (a+h)^2} r &= \sum'_{a^2 < r < (a+h)^2} \sum_{0 \leq t < r} 1 = \sum_{0 \leq t < (a+h)^2} \sum'_{\max(a^2, t) < r < (a+h)^2} 1 \\ &= \sum_{0 \leq t \leq a^2} \sum'_{a^2 < r < (a+h)^2} 1 + \sum_{a^2 < t < (a+h)^2} \sum'_{t < r < (a+h)^2} 1 \\ &= \frac{\pi^2}{32} \left(\sum_{0 \leq t \leq (a+h)^2} ((a+h)^4 - a^4) + \sum_{a^2 < t < (a+h)^2} ((a+h)^4 - t^2) \right) \\ &= \frac{\pi^2}{32} \left((a+h)^4 a^2 - a^6 + (a+h)^4 ((a+h)^2 - a^2) - \frac{1}{3} ((a+h)^6 - a^6) \right) \\ &= \frac{\pi^2}{3 \cdot 16} ((a+h)^6 - a^6); \end{aligned}$$

We derive the following formula for K_2 :

$$\begin{aligned} K_2 &= \frac{\pi}{4} \sum_{0 < \xi_1 < 1} \left\{ (a+h)^2 (1 - \xi_1^2) \frac{\pi^2}{32} ((a+h)^4 - a^4) \right. \\ &\quad \left. - (1 - \xi_1^2) \frac{\pi^2}{3 \cdot 16} ((a+h)^6 - a^6) + ((a+h)^2 - a^2) (1 - \xi_1^2) \frac{\pi^2}{32} a^4 \right\} \\ &= \frac{\pi^3}{12 \cdot 32} ((a+h)^6 - a^6) \sum_{0 < \xi_1 < 1} (1 - \xi_1^2). \end{aligned}$$

Since $\xi_1 = \frac{l}{m}$, $0 < l < m \leq L$, we have

$$\sum_{0 < \xi_1 < 1} (1 - \xi_1^2) = \int_0^L \int_0^m \left(1 - \left(\frac{l}{m} \right)^2 \right) dl dm = \frac{L^2}{3}$$

and

$$K_2 = \frac{\pi^3}{32 \cdot 36} L^2 ((a+h)^6 - a^6).$$

The fact that $K = K_1 + K_2$ implies the relation

$$K = \left(\frac{1}{5} + \frac{1}{9} \right) \frac{1}{128} \pi^3 L^2 ((a+h)^6 - a^6) = \frac{7\pi^3}{2880} L^2 ((a+h)^6 - a^6).$$

Remarks.

1. The formula for K is an approximate formula.
2. This formula is sufficiently exact if L is large and a is not too small but significantly less than L ; for example, $a = \sqrt{L}$, $L \rightarrow +\infty$.
3. If a is chosen to be not large, for example, $a = 4$, $h = \frac{1}{2}$, or $a = 3$, $h = \frac{1}{2}$, or $a = 5$, $h = \frac{1}{2}$, then we can obtain sufficiently exact formulas for $L \geq 8$, since, in this case, we can exactly calculate the multiplicities of l and the l themselves (as sums of two squared numbers). But, in this case, we must perform specific calculations.

Example of calculations for small a .

Let $a = 4$, $h = 1$; $L > 5$.

In system (22), l satisfy the condition

$$16 \leq l = l_1^2 + l_2^2 < 25.$$

The pairs (l_1, l_2) satisfying this condition are:

$$(0, 4), (1, 4), (2, 4), (3, 3), (4, 1), (4, 1), (4, 0),$$

i.e., from (22) we obtain two systems for $l = 16$, $l = 17$, and $l = 20$ and one system for $l = 18$:

$$\begin{aligned} 16 < r &\leq 25 - 9\xi^2, \\ 0 &\leq \xi < 1, \end{aligned} \tag{29}$$

$$\begin{aligned} 17 < r &\leq 25 - 8\xi^2, \\ 0 &\leq \xi < 1, \end{aligned} \tag{30}$$

$$\begin{aligned} 20 < r &\leq 25 - 5\xi^2, \\ 0 &\leq \xi < 1, \end{aligned} \tag{31}$$

$$\begin{aligned} 18 < r &\leq 25 - 7\xi^2, \\ 0 &\leq \xi < 1. \end{aligned} \tag{32}$$

To calculate the number of sets (m_1, m_2, k_1, k_2) for which $r = m_1^2 + m_2^2 + k_1^2$ satisfies systems (29)–(32), we can either use the computer software (in this case, we must divide ξ into intervals, where $9\xi^2$ lies between two neighboring integers from 0 to 9, the same for $8\xi^2$, $5\xi^2$, and $7\xi^2$, and calculate the number of sets directly) or use the formula for the number of integer points in the four-dimensional ball and thus find the points with positive coordinates. This is an approximate formula and if

$$m_1^2 + m_2^2 + k_1^2 \leq x, \quad m_i, k_i \geq 0,$$

then this number is equal to

$$\frac{\pi^2}{32} x^2. \tag{33}$$

Therefore, for system (22), we obtain the number of sets

$$\begin{aligned} \frac{\pi^2}{32} \sum_{0 \leq \xi < 1} \left\{ ((25 - 9\xi^2)^2 - 16^2) \cdot 2 + ((25 - 8\xi^2)^2 - 17^2) \cdot 2 \right. \\ \left. + ((25 - 5\xi^2)^2 - 20^2) \cdot 2 + ((25 - 7\xi^2)^2 - 18^2) \right\}. \end{aligned} \tag{34}$$

Since the number of such l is equal to 2, we use the factor 2. In system (23), we have

$$l = l_1^2 + l_2^2 < 16,$$

and the pairs (l_1, l_2) are: $(0, 3)$, $(0, 2)$, $(0, 1)$, $(0, 0)$, $(1, 2)$, $(1, 3)$, $(1, 0)$, $(2, 0)$, $(2, 1)$, $(2, 2)$, $(2, 3)$, $(3, 0)$, $(3, 2)$, $(3, 2)$.

l=	0	1	4	9	5	10	8	13
multiplicity	1	2	2	2	2	2	1	2

System (23) for each l has the form

$$16(1 - \xi^2) + l\xi^2 < r \leq 25(1 - \xi^2) + l\xi^2, 0 \leq \xi < 1.$$

Using (33), we obtain the following formula (with multiplicity of l taken into account) for the number of sets for system (23):

$$\begin{aligned}
& \frac{\pi^2}{32} \sum_{0 \leq \xi < 1} \left\{ (25(1 - \xi^2))^2 - (16(1 - \xi^2))^2 \right. \\
& \quad + ((25(1 - \xi^2) + \xi^2)^2 - (16(1 - \xi^2) + \xi^2)^2) \cdot 2 \\
& \quad + ((25(1 - \xi^2) + 4\xi^2)^2 - (16(1 - \xi^2) + 4\xi^2)^2) \cdot 2 \\
& \quad + ((25(1 - \xi^2) + 9\xi^2)^2 - (16(1 - \xi^2) + 9\xi^2)^2) \cdot 2 \\
& \quad + ((25(1 - \xi^2) + 5\xi^2)^2 - (16(1 - \xi^2) + 5\xi^2)^2) \cdot 2 \\
& \quad + ((25(1 - \xi^2) + 10\xi^2)^2 - (16(1 - \xi^2) + 10\xi^2)^2) \cdot 2 \\
& \quad + ((25(1 - \xi^2) + 8\xi^2)^2 - (16(1 - \xi^2) + 8\xi^2)^2) \cdot 1 \\
& \quad \left. + ((25(1 - \xi^2) + 13\xi^2)^2 - (16(1 - \xi^2) + 13\xi^2)^2) \cdot 2 \right\} \\
& = \frac{\pi^2}{32} \sum_{0 \leq \xi < 1} \left\{ (25(1 - \xi^2))^2 - (16(1 - \xi^2))^2 \right. \\
& \quad + ((25 - 24\xi^2)^2 - (16 - 15\xi^2)^2) \cdot 2 \\
& \quad + ((25 - 21\xi^2)^2 - (16 - 12\xi^2)^2) \cdot 2 \\
& \quad + ((25 - 16\xi^2)^2 - (16 - 7\xi^2)^2) \cdot 2 \\
& \quad + ((25 - 20\xi^2)^2 - (16 - 11\xi^2)^2) \cdot 2 \\
& \quad + ((25 - 15\xi^2)^2 - (16 - 6\xi^2)^2) \cdot 2 \\
& \quad + ((25 - 17\xi^2)^2 - (16 - 8\xi^2)^2) \cdot 1 \\
& \quad \left. + ((25 - 12\xi^2)^2 - (16 - 3\xi^2)^2) \cdot 2 \right\}. \tag{35}
\end{aligned}$$

Since $\xi = \frac{m_3}{l_3}$, $0 \leq m_3 \leq l_3$, $1 \leq l_3 \leq L$, the number of sets corresponding to Case (I) is either calculated directly according to (34)–(35) or is approximately replaced by an integral of the form

$$\int_0^L dl \int_0^l dm \dots$$

equal to

$$\frac{\pi^2}{64} L^2 \mu,$$

where

$$\begin{aligned}
\mu = \int_0^1 & \left\{ 2((25 - 9u^2)^2 - 16^2) + 2((25 - 8u^2)^2 - 17^2) \right. \\
& + 2((25 - 5u^2)^2 - 20^2) + ((25 - 7u^2)^2 - 18^2) \\
& + (25(1 - u^2))^2 - (16(1 - u^2))^2 + 2((25 - 24u^2)^2 - (16 - 15u^2)^2) \\
& + 2((25 - 21u^2)^2 - (16 - 12u^2)^2) + 2((25 - 16u^2)^2 - (16 - 7u^2)^2) \\
& + 2((25 - 20u^2)^2 - (16 - 11u^2)^2) + 2((25 - 15u^2)^2 - (16 - 6u^2)^2) \\
& \left. + 1((25 - 17u^2)^2 - (16 - 8u^2)^2) + 2((25 - 12u^2)^2 - (16 - 3u^2)^2) \right\} du.
\end{aligned}$$

Thus, we have

$$K_1 = \frac{\pi^2}{64} L^2 \mu.$$

Case (II), i.e., the calculation of K_2 is the same; only the multiplicity of r , $16 < r < 25$, $r = m_1^2 + m_2^2 + k_1^2 + k_2^2$, is calculated either by an exhaustive search of the sets (m_1, m_2, k_1, k_2) (also for $r < 16$) or by the formula for the number of integer points in the four-dimensional ball; the number of sets for $l \leq r$ from (27) and (28) is calculated approximately by the formula

$$\frac{\pi}{4} x.$$

Therefore, if $\kappa(r)$ is the multiplicity of r , then

$$\begin{aligned} K_2 &= \sum_{0 < \xi_1 < 1} \left\{ \sum_{16 < r < 25} \kappa(r) \frac{\pi}{4} (25 - r) (1 - \xi_1^2) + \sum_{r < 16} \frac{\pi}{4} \kappa(r) (25 - 16) (1 - \xi_1^2) \right\} = \\ &= \frac{L^2}{2} \frac{2}{3} \frac{\pi}{4} \left(\sum_{16 < r < 25} \kappa(r) (25 - r) + 9 \sum_{r < 16} \kappa(r) \right). \end{aligned}$$

But we have already calculated that

$$\begin{aligned} \sum_{r < 16} \kappa(r) &= \frac{\pi^2}{32} 4^4; \\ \sum_{16 < r < 25} \kappa(r) &= \frac{\pi^2}{32} (5^4 - 4^4); \\ \sum_{16 < r < 25} \kappa(r) r &= \frac{\pi^2}{48} (5^6 - 4^6). \end{aligned}$$

Hence we have

$$\begin{aligned} K_2 &= \frac{\pi}{12} L^2 \left(25 \cdot \frac{\pi^2}{32} (5^4 - 4^4) - \frac{\pi^2}{48} (5^6 - 4^6) + 9 \frac{\pi^2}{32} \cdot 4^4 \right), \\ K &= K_1 + K_2 = c L^2. \end{aligned}$$

All these formulas are approximate.

The number $\mu = \int_0^1 \dots dm$ and the constant c in the formula for K have not been calculated, but these calculations are very simple.

The author wishes to express his deep gratitude to Professor A. A. Karatsuba and to G. V. Koval' for their help in preparing this paper.

References

- [1] V. P. Maslov. *Ultrascond quantization of thermodynamics*, Theor. Math. Phys., 132, 1222–1232 (2002)..
- [2] V. P. Maslov. *Quantization of Thermodynamics and Ultrascond Quantization* [in Russian], Computer Research Inst., Moscow (2001).
- [3] F. A. Berezin. *Methods of Second Quantization* [in Russian], Nauka, Moscow (1965); English transl., Acad. Press, New York (1986).

- [4] V. P. Maslov. *Operator Methods* [in Russian], Nauka, Moscow (1973); English transl., *Operational Methods*, Mir, Moscow (1976).
- [5] V. P. Maslov and O. Yu. Shvedov. *The Complex Germ Method in the Many-Particle Problem and in Quantum Field Theory* [in Russian], Editorial URSS, Moscow (2000).
- [6] N. N. Bogoliubov. *Selected Works* [in Russian], Vol. 2, Naukova Dumka, Kiev (1970).
- [7] M. V. Fedoryuk. *Method of stationary phase and pseudodifferential operators*, Russ. Math. Surveys, 26, no. 1, 65–115 (1971).
- [8] V. P. Maslov. *Complex WKB Method for Nonlinear Equations* [in Russian], Nauka, Moscow (1977); English transl., *The Complex WKB Method for Nonlinear Equations*, Vol. 1, *Linear Theory*, Birkhäuser, Basel (1994).