

# Coherence Network in the Quantum Hall Bilayer

H. A. Fertig<sup>1</sup> and Ganpathy Murthy<sup>2</sup>

1. Department of Physics, Indiana University, Bloomington, IN 47405

2. Department of Physics and Astronomy, University of Kentucky, Lexington, KY 40506-0055

(Dated: March 23, 2024)

Recent experiments on quantum Hall bilayers near total filling factor 1 have demonstrated that they support an "imperfect" two-dimensional superfluidity, in which there is nearly dissipationless transport at non-vanishing temperature observed both in counterflow resistance and interlayer tunneling. We argue that this behavior may be understood in terms of a coherence network induced in the bilayer by disorder, in which an incompressible, coherent state exists in narrow regions separating puddles of dense vortex-antivortex pairs. A renormalization group analysis shows that it is appropriate to describe the system as a vortex liquid. We demonstrate that the dynamics of the nodes of the network leads to a power law temperature dependence of the tunneling resistance, whereas thermally activated hops of vortices across the links control the counterflow resistance.

PACS numbers:

Introduction{ Over the past decade, there has been accumulating evidence that excitonic superfluid-like [1] properties may be present in quantum Hall bilayer systems [2] when the filling factor of each layer is near  $\nu = 1/2$  and the layers are sufficiently close. The relevance of excitons to this system may be understood by using a filled Landau level in a single layer as a starting reference state; equal densities in the layers may then be reached by creating a particle-hole condensate [3, 4]. It has long been suspected that such a state might have superfluid properties [5].

The most dramatic superfluid-like properties of this system are observed when the layers are separately contacted and a current is passed from one layer to the other. For contacts at opposite ends of the sample, the resulting tunneling conductance  $\sigma_T$  has a sharp resonance precisely at zero voltage [6], which is reminiscent of the DC Josephson effect. When the contacts are on the same side of the sample and the layers on the opposite side are short-circuited, so that currents in the two layers flow in opposite directions ("counterflow" (CF)), the voltage drop in either layer along the direction of flow tends to zero in the low temperature limit [7, 8]. This may be understood as nearly dissipationless flow of electron-hole pairs. Interestingly, the temperature dependences of these two behaviors are very different: the tunneling resistance  $1/\sigma_T$  falls much more slowly than the dissipative in-plane resistance.

The "imperfect" superfluidity apparent in these experiments suggests that a good starting point for understanding this system is in terms of a condensed state with interlayer coherence. However, a true two-dimensional superfluid should have a power law  $I \sim V$  in CF below some critical Kosterlitz-Thouless temperature  $T_{KT}$ , whereas an Ohmic response is observed at all available temperatures. Moreover, a true DC Josephson effect would yield an infinite tunneling conductance below  $T_{KT}$  rather than just a sharp resonance. It is also remarkable that in both the tunneling and CF geometries, current appears to flow

over the entire length of the system, whereas for a true superfluid the current would decay within a Josephson length of the edge [9], which is far smaller. The linear response of the quantum Hall bilayer, exhibiting dissipationless transport, if at all, only at zero temperature, is a new transport regime for condensed matter systems. Most theoretical studies of this system suggest that disorder must be crucially involved in the mechanism(s) that lead to this behavior [10, 11, 12, 13].

In this work, we describe a state where the experimentally observed behavior emerges naturally, and which is indeed induced by the disorder environment of the quantum Hall bilayer. We will argue below that the system breaks up into large regions in which the interlayer coherence is small or absent, separated by narrow regions in which the coherence is relatively strong. The resulting structure is a coherence network, with long, quasi-one-dimensional links connected at nodes. We will demonstrate that dissipation in tunneling may be understood in terms of the phase dynamics of the nodes, whereas dissipation in CF occurs due to thermally activated hops of vortices across the links. Further experimental consequences of our model are discussed below.

Nonlinear Screening and the Coherence Network { We start with the Efros model for nonlinear screening of potential fluctuations in a quantum Hall system [14]. This screening generates puddles of quasiholes or quasi-electrons, which are locally dense and mobile enough to destroy the incompressibility of the quantum Hall fluid. The puddles are separated by narrow regions, with  $\ell_0 = \hbar c/eB$  (the magnetic length) incompressible strips which follow the equipotentials of the total effective potential, and within which there is a charge gap. The strips form links which join together at nodes (associated with saddle points of the disorder potential) to form a random network.

For the bilayer system, the incompressible state at total filling factor  $\nu = 1$  has a phase degree of freedom that may be understood as the wavefunction phase

for excitons in the incompressible regions of the system. The conjugate variable to this phase angle is the interlayer charge imbalance, which may be denoted as  $S_z$ . For equal layer populations, on average  $\langle S_z \rangle = 0$ , and the system becomes analogous to an easy-plane Heisenberg ferromagnet [15]. Vortices of the phase field ("merons") carry both an electric charge ( $e=2$  for either vorticity) and an interlayer electric dipole moment [15, 16, 17]. Thus, in this context the links and nodes of the network carry a phase degree of freedom, while the puddles are coded with vortex-antivortex pairs [18].

The presence of meron-antimeron pairs at link edges implies that the phase angle  $\theta(r)$  turns over many times along a given link. When tunneling is included, these overturns bear a close resemblance to kink solitons of the sine-Gordon model [15]. As discussed below, the phase dynamics of the links is conveniently described in terms of these solitons. Our picture of this coherence network is illustrated in Fig. 1.

Renormalization Group Analysis { For counterflow experiments, dissipation can only occur when merons { i.e., vortices { are mobile and unbound [17]. For this to occur, the vortices in the coherence network must be in a liquid phase; i.e., their discreteness should not inhibit their ability to screen "charge" (i.e., enforced vorticity) at long distances. We begin with a classical model on a square lattice network,

$$H_{CN} = \sum_r \left[ \frac{1}{2} K \sum_{\langle r, r' \rangle} \left[ \theta(r) - \theta(r') + A(r) \right]^2 + h \cos \left( \frac{d}{a_0} \frac{r}{K} \right) \right] = e^{2\gamma} \sum_{r=K}^S \left[ \frac{r^2}{1 + \frac{r^2}{K^2}} + \frac{1}{K} + 2 \right]$$

where  $m_{xy}(r)$  are integer degrees of freedom representing the number of solitons on the link extending from the node at  $r$  in the  $x$  or  $y$  direction, and  $A(r)$  is a Gaussian random variable obeying  $\langle A(r)A(r') \rangle = \delta(r-r')$ ;  $r \neq 0$  which models the effect of disorder. The parameter  $h$  is proportional to the interlayer tunneling matrix element, which explicitly breaks the  $U(1)$  symmetry in energetically favoring  $\theta = 0$ .

It is convenient to recast the problem explicitly in terms of vortices, which can be done by standard techniques [19], leading to  $Z = \int \mathcal{D} \theta \mathcal{D} B e^{-H_V}$ , with

$$H_V = 2 \sum_r \left[ \frac{1}{2} K \left[ \nabla \theta(r) + B(r) + A(r) \right]^2 + \gamma_h \cos 2 \theta(r) + E_c \frac{\partial B(r)}{\partial y} \right] : \quad (1)$$

In this representation  $B$  is an integer field that resides on vertical bonds, and the vorticity is  $m(r) = \partial B / \partial y$ . Finally,  $E_c$  represents a core energy for these vortices, and  $\gamma_h = e^{K_B T} h$ . This form of  $H_V$  is appropriate when  $\gamma_h \ll 1$ , which is easily satisfied under experimental conditions. For  $\gamma_h = 0$  and  $A = 0$ ,  $m$  may be integrated out to recover the usual Coulomb gas.

To facilitate the renormalization group (RG) analysis, we replace the integer field  $B(r)$  with a continuous field

$b(r)$  and add a term  $\sum_r \cos[2\pi b(r)]$  that tends to keep  $b$  at its intended integer values. By matching low energy configurations for the integer and continuous field versions of the theory, one may show  $\gamma_h \ll E_c$  for small  $E_c$  and  $\gamma_h$ . The crucial point is that the irrelevance of  $\gamma_h \ll E_c$  will indicate a liquid state of vortices.

In performing the RG,  $b$  is rescaled according to  $b^0(r^0)e^{-\gamma} = b(r)$ , where  $e^{-\gamma}$  is the rescaling factor, in order to keep the Gaussian part of the Hamiltonian of fixed form. As  $b$  shrinks with rescaling, it is natural to expand  $\cos[2\pi b(r)]$  in a power series in  $b$ , and treat the fourth and higher order terms perturbatively. The quadratic term, however, introduces a contribution to the fixed point of the form  $\frac{1}{2} r^2 b(r)^2$ . A key observation is that this term limits large separations for vortices, so that if  $r \neq 0$  at the fixed point, the system is in a bound vortex state [20, 21]. In the clean limit ( $A = 0$ ), it has been demonstrated that this happens if the temperature is too low,  $E_c$  is too large, or if  $h$  is too large. By contrast, for  $r = 0$ , states of arbitrarily large "vortex dipole moment" may be found in the thermodynamic limit, so that the vortices are effectively unbound [21].

To generalize this to the disordered case, we use the replica trick before proceeding with the RG analysis. In the  $n \rightarrow 0$  limit, the flow equations become, to lowest order in  $E_c$  and  $\gamma_h$ ,

$$\frac{d\gamma}{d\ell} = \gamma^2 \quad \frac{d\gamma_h}{d\ell} = \left( 2 - \frac{P}{K r} \right) \gamma_h$$

where  $\gamma$  is the disorder strength,  $\gamma = \gamma_0$  is a wavevector cutoff,  $a_0$  is the distance between nodes,  $\gamma_0$  is a length scale with initial value  $\gamma(0) = \frac{1}{2} E_c = 4 \pi^2 K a_0$ , and the initial value of  $r(0) = E_c$ . The parameter  $P$  is a number of order unity. To this order in perturbation theory, the parameter  $K$  remains constant. For large disorder and/or small  $K$  [13, 22], it is apparent that  $r, \gamma_h \rightarrow 0$ .

Since all signs of the discreteness of the underlying the vortex density has scaled away at this fixed point, it is a liquid. In this phase neither the stiffness  $K$  nor the symmetry-breaking field  $h$  are effective at preventing phase overturns along a column of nodes (say, at the edge of the system) if their phase angles are driven by an arbitrarily small torque [23]. Moreover, the irrelevance of  $\gamma_h$  means the tunneling term  $h \cos$  may be treated perturbatively in the vortex liquid phase. We now exploit this observation to understand the transport properties of the coherence network.

Dissipation in Transport: Tunneling and Counterflow { We can understand transport in the coherence network based on the following considerations. (i) The total current in the sample is the sum of a CF current and the quantum Hall edge current. Dissipation in the latter is

negligible in both the tunneling and CF geometries we consider here, and so is ignored in what follows. (ii) We assume the solitons are unpinned in the links, as is appropriate for slowly varying disorder, and treat them as an elastic system. The displacement field  $u(x)$  is related to the counterflow current via  $I_{CF} = -\frac{e}{h} \frac{\partial u}{\partial x} K \frac{\partial u}{\partial x}$  (with  $K = \hbar / 4\pi e$ ). From the Josephson relation, it follows that the interlayer electric potential is  $V_{int} = \frac{\hbar}{e} \frac{\partial \theta}{\partial t} = \frac{\hbar}{e} \frac{\partial \theta}{\partial t} u = b_0$ , where  $b_0$  is the average spacing of the solitons. A simple effective Gaussian theory may be written for the solitons on the links [24], whose form would control dissipation at sufficiently high frequencies. Such a purely quadratic theory however cannot dissipate energy at zero frequency [25]. (iii) CF current can be degraded by hopping of merons across the link. This is a thermally activated process with energy  $\epsilon_{link}$ , because the merons must cross the barrier of the incompressible link. (iv) The nodes are relatively meron-free regions with a high stiffness, and can be treated as rigid rotors which are subject to torques  $F_{link} / I_{CF}^{link}$  from each attached link, and a force  $\hbar \sin$  due to the tunneling.

Consider first the tunneling geometry. Since the current flows into (say) the top layer on the left and leaves via the bottom layer on the right, the CF current points in opposite directions at the two ends of the sample. From (ii) above this implies the currents cause the underlying phase angles in the links and nodes to rotate in the same direction, and may be modeled by forcing in solitons at one sample end and removing them at the other, as illustrated in Fig. 2. The dynamics of a typical node may be described by a Langevin equation

$$\frac{d^2}{dt^2} = \sum_{links} F_{link} - \gamma \frac{d}{dt} - \hbar \sin + \langle \epsilon(t) \rangle \quad (2)$$

Here  $I$  is the effective moment of inertia of a rotor, proportional to the capacitance of the node,  $\epsilon$  is a random (thermal) force, and  $\gamma$  is the viscosity due to dissipation from the other node rotors in the system. For a small driving force, the node responds viscously, and the resulting rotation rate determines the rate of flow of solitons via  $\dot{\theta} = F_{link} / I$ . From (ii) and (iv) above one sees that the viscosity  $\gamma$  is proportional to the tunneling conductance  $G_T$  of the system. For  $k_B T \ll \hbar$  one may show the viscosity for an individual node to be [26]

$$\gamma = \frac{\hbar}{2} + \frac{\hbar^2}{2(k_B T)^{3/2}} \quad (3)$$

As each node contributes the same amount to the total viscosity, the total response of the system to the injected CF current obeys

$$I_{CF} / N_{nodes} = \frac{e^2 V_{int}}{h} = G_T V_{int} \quad (4)$$

Note that because the nodes respond viscously, the tunneling conductance is proportional to the area of the bi-

layer. This is so far consistent with experiment but difficult to understand without the network geometry posited here [10]. The dependence of the tunneling conductance on temperature, system area, and  $\hbar$  each constitute an experimentally falsifiable prediction of the coherence network model of the quantum Hall bilayer [27, 28].

Now consider the CF geometry. This is created by short-circuiting the layers at one end of the sample, which we will take to be on the right. There the rotor angles should be subject to free boundary conditions, leading to  $F / \hbar = 0$ . The injected CF current on the left leads to a compression of the solitons (or a rarefaction of antisolitons). In a uniform superfluid, such a force would be fully balanced by the effective force due to tunneling, and the CF current would flow without dissipation. In the coherence network, however, at any  $T \neq 0$ , merons are able to thermally hop across the links to create or destroy solitons, and the other (unpinned) solitons move in response, leading to dissipation.

In equilibrium, the rates of creation and annihilation of solitons in a link obey detailed balance. However, the presence of the "force"  $F / I_{CF}$  destroys this balance, and leads to a net meron current

$$I_{meron} = -I_{CF} e^{-\epsilon_{link}/k_B T}; \quad (5)$$

where  $\epsilon$  is a dimensionless number depending on the details of the link disorder. If, for example, there is a net annihilation of solitons with time on a given link, solitons will move in from the left to take their place. The rate of change of  $\theta$  due to such processes at a point inside the sample a distance  $x$  from the left end is

$$\frac{d\theta(x)}{dt} = 2 \int_x^L I_{meron}(\theta^0) d\theta^0 \quad (6)$$

For temperatures satisfying  $\hbar \gg k_B T \gg \epsilon_{link}$ , the degrading of  $I_{CF}$  due to the nodes may be ignored. In this case, Eqs. 5 and 6 together with the Josephson relation yield an activated longitudinal resistance measured between probes  $x$  apart on a single layer of the form

$$R_{xx} = \rho(x) \frac{\hbar}{e^2} e^{-\epsilon_{link}/k_B T} \quad (7)$$

which is consistent with experiment.

In the CF geometry, the meron current  $I_{meron}$  also creates a Hall voltage within individual layers because merons carry an interlayer dipole moment, leading to nonequilibrium charge imbalances at the upper and lower edges of the sample of opposite sign. Assuming a  $T$ -independent resistive relaxation at the edges (where the electrons are likely to be compressible), this creates a Hall voltage of opposite sign on the two layers which is activated, with the same activation energy as the longitudinal resistance. This is consistent with measurements in electron samples [7, 29].

Experimental Consequences { As shown in this work, the in perfect super uidity observed in quantum Hall bilayers may be understood if we assume the electrons have organized into a coherence network. Several experimental tests of this hypothesis are possible. These include the temperature and size dependence of the tunneling conductance discussed above. Another interesting test would be measurement of the finite frequency tunneling conductance, for which the response of the links should become accessible. An additional possibility is the introduction of an artificial weak link across the direction of counter flow current. This would create a favored channel for soliton destruction/creation that should lead to a decrease in the activation energy for the CF resistance and an enhanced Hall voltage at its endpoints.

The authors would like to thank S. Das Sarma, J.P. Eisenstein, M. Shayegani, and E. Tutuc for useful discussions. This work was supported by the NSF via Grant Nos. DMR 0454699 (HAF) and DMR 0311761 (GM).

- 
- [1] L. Keldysh and Y. Kopaev, Sov. Phys. 6, 2219 (1965).
  - [2] L.V. Butov et al., Nature 417, 47 (2002); D. Snoke, Science 298, 1368 (2002).
  - [3] See J.P. Eisenstein and A.H. MacDonald, Nature 432, 691 (2004) and references therein.
  - [4] H.A. Fertig, Phys. Rev. B 40, 1087 (1989).
  - [5] X.G. Wen and A. Zee, Phys. Rev. Lett. 69, 1811 (1992); Phys. Rev. B 47, 2265 (1993); Z.F. Ezawa and A. Iwazaki, Phys. Rev. B 47, 7295 (1993).
  - [6] I.B. Spielman, J.P. Eisenstein, L.N. Pfeiffer, and K.W. West, Phys. Rev. Lett. 84, 5808 (2000).
  - [7] M. Kellogg, J.P. Eisenstein, L.N. Pfeiffer, and K.W. West, Phys. Rev. Lett. 93, 036801 (2004).
  - [8] E. Tutuc, M. Shayegani, and D. Huse, Phys. Rev. Lett. 93, 036802 (2004).
  - [9] R. Ferrell and R. P range, Phys. Rev. Lett. 10, 479 (1963).
  - [10] L. Balents and L. Radzihovsky, Phys. Rev. Lett. 86, 1825 (2001); A. Stern, S.M. Girvin, A.H. MacDonald, and Ning Ma, Phys. Rev. Lett. 86, 1829 (2001).
  - [11] H.A. Fertig and J.P. Straley, Phys. Rev. Lett. 91, 046806 (2003).
  - [12] Z. Wang, Phys. Rev. Lett. 92, 136803 (2004).
  - [13] D. Sheng, L. Balents, and Z. Wang, Phys. Rev. Lett. 91, 116802 (2003).
  - [14] A. Efros, Sol. St. Commun. 65, 1281 (1988); A. Efros, F. Pikus, and V. Bumett, Phys. Rev. B 47, 2233 (1993).
  - [15] K. Moon, H. Mori, K. Yang, S. Girvin, A. MacDonald, L. Zheng, D. Yoshioka, and S. Zhang, Phys. Rev. B 51, 5138 (1995); K. Yang et al., Phys. Rev. B 54, 11644 (1996).
  - [16] D.H. Lee and C.L. Kane, Phys. Rev. Lett. 64, 1313 (1990).
  - [17] D.A. Huse, cond-mat/0407452 (2004).
  - [18] The compressible puddles might be describable in terms of composite fermions; this does not affect the transport properties of the network.
  - [19] J.V. Jose, L.P. Kadano, S. Kirkpatrick, and D.R. Nelson, Phys. Rev. B 16, 1217 (1977).

[20] H.A. Fertig, Phys. Rev. Lett. 89, 035703 (2002).




FIG. 1: Representation of coherence network. Links and nodes separate puddles of fermions (circles). Fermion charge and electric dipole moments indicated inside circles, as are strings of overturned phase connecting fermion-antifermion pairs.




FIG. 2: Counter flow current in tunneling geometry, created by adding solitons at one edge of sample and removing them at the other, shown for a single set of links.

- [21] H.A. Fertig and J.P. Straley, Phys. Rev. B 66, 201402(R) (2002).
- [22] Y. Joglekar and A.H. MacDonald, Phys. Rev. B 64, 155315 (2001); M.V. Milovanovic and I. Stanic, cond-mat/0412536.
- [23] This behavior has been demonstrated in related simulations at high temperature. See W. Zhang and H.A. Fertig, Phys. Rev. B (in press; cond-mat/0502402).
- [24] H.A. Fertig and Ganpathy Murthy, unpublished.
- [25] D.L. Maslov and M. Stone, Phys. Rev. B 52, R5539 (1995); I. Sa and H. Schulz, ibid. 52, R17040 (1995).
- [26] W. Dieterich, I. Peschel, and W.R. Schneider, Z. für Physik B, 27, 177 (1977).
- [27] Experimental measurements of  $T^{-1}$  so far appear to saturate at low refrigerator temperature. However this may be due to the difficulty of isolating the electrons from noise and cooling them to extremely low temperatures.
- [28] A further temperature dependence arises due to a downward renormalization of the superfluid stiffness in the links and nodes by vortex-antivortex pairs. We expect the  $T^{-3/2}$  dependence to apply in the temperature range  $T_K < T < T_h$ .
- [29] In hole samples it is found that  $R_H$  vanishes with temperature much faster than  $R_{xx}$ . This may be due to the greater Landau level mixing which will affect the dipole moment of the fermions, or due to greater inhomogeneity affecting the relaxation of interlayer charge imbalances.