

Quantum dissociation of an edge of a Luttinger liquid

Eugene B. Kolomeisky and Michael Timmins

Department of Physics, University of Virginia, P. O. Box 400714, Charlottesville, Virginia 22904-4714

In a Luttinger liquid phase of one-dimensional molecular matter the strength of zero-point motion can be characterized by dimensionless De Boer's number quantifying the interplay of quantum fluctuations and two-body interactions. Selecting the latter in the Morse form we show that dissociation of the Luttinger liquid is a process initiated at the system edge. The latter becomes unstable against quantum fluctuations at a value of De Boer's number which is smaller than that of the bulk instability which parallels the classical phenomenon of surface melting.

PACS numbers: 68.65.-k, 61.46.+w, 71.10.Pm, 05.30.-d

A classical three-dimensional solid melts through a first-order transition at a temperature when the free energies of the solid and liquid phases coincide. At sufficiently low temperatures quantum effects dominate and a quantum solid can melt due to zero-point motion [1].

The most curious feature of classical melting is the difficulty in overheating the solid while supercooling the liquid is easy. The latter is expected for the first-order transition while the former is explained by the phenomenon of surface melting: often, as the bulk transition is approached, the melting begins at the free surface of a solid. The surface melting is well-documented experimentally, and phenomenologically it can be viewed as a wetting of the solid by its own melt [2].

A well-understood example of surface melting of a quantum solid is that of the edge melting of the two-dimensional Wigner crystal in a strong magnetic field [3].

The goal of this note is to point out that strictly one-dimensional matter with a free edge can also exhibit an analog of surface melting. Fundamentally this happens because the edge represents a zero-dimensional system subject to stronger quantum fluctuations than the one-dimensional bulk. Due to broken translational symmetry, zero-point motion modifies the cohesive properties of the edge differently from those of the bulk.

Experimentally one-dimensional matter can be realized in carbon nanotube bundles [4]. The latter can play a role of one-dimensional hosts for foreign particles that can find themselves bound in the interstitial channels or inside the tubes [5]. Additionally one-dimensional atomic chains can be constructed on selected templates with the help of scanning tunneling microscopy, or via self-assembly of the deposited material of the chain [6].

Consider a many-body system of identical particles of mass m with pairwise interaction $V(h)$ corresponding to the molecular matter [7]: at large interparticle separation h the interaction is dominated by weak rapidly decaying van der Waals attraction, while at short distances there is a strong overlap repulsion. As a result, the pair potential $V(h)$ has an asymmetric minimum at some intermediate h . Assume that the pair potential is of the form

$$V(h) = \epsilon U(h/l - Q_0), \quad (1)$$

where ϵ is the energy scale of the potential, l is the potential range, Q_0 is a dimensionless parameter, and $U(y)$ is a function common to a family of substances. With this choice the quantum theorem of the corresponding states [8] holds stating that every property measured in appropriate dimensionless units is only determined by the function $U(y)$, particle statistics and De Boer's number

$$\lambda_0 = \frac{\hbar}{\pi l (2m\epsilon)^{1/2}}, \quad (2)$$

measuring the intensity of zero-point motion.

The possibility of several bulk phases in the system translates into the corresponding number of the branches of the energy as a function of λ_0 ; the lowest of them singles out the ground state of the system. When two energy curves cross, the ground state changes via a first-order phase transition. For sufficiently large λ_0 and zero pressure the ground state must correspond to individual particles infinitely far apart from each other. This is a monoatomic gas which will be chosen as the zero reference point for the energy.

In what follows we select the pair interaction potential in the Morse form [9]:

$$V(h) = \epsilon(e^{-2(h/l - Q_0)} - 2e^{-(h/l - Q_0)}), \quad (3)$$

where ϵ is the depth of the potential well and Q_0 is the location of the minimum of (3) measured in units of the potential range l . Similar to the applications of the Lennard-Jones potential to laboratory molecular systems [7], the only reason behind this choice is the possibility of analytic progress. Morse parameters for a series of molecular substances and corresponding De Boer's numbers (2) were computed in Ref.[10]. Hereafter the energy and length scales will be measured in units of ϵ and l , respectively. As appropriate for molecular substances, we restrict ourselves to nearest-neighbor interactions.

In the classical limit, $\lambda_0 = 0$, the ground state of the system is a crystal; its quantum counterpart for sufficiently small λ_0 is a Luttinger liquid [11] whose properties have been computed in Ref.[10] as follows:

The length of any bulk bond h as a function of imaginary time τ is viewed as a quantum-mechanical degree

of freedom subject to the external potential $V(h)$. This bond joins together two half-infinite segments representing the rest of the system, the “bath”. After the bath is approximated by a harmonic liquid, the latter can be integrated out away from the anharmonic bond leading to a problem of the Caldeira-Leggett type [12]. The latter has been analyzed by a combination of variational and renormalization-group techniques, and it has been demonstrated that the approximation is a controlled way of dealing with the interplay of zero-point motion and anharmonicity of the two-body interaction [10]. Similar consideration applied to the edge bond of a half-infinite Luttinger liquid leads to the Euclidian action of the form

$$S_{edge} = \frac{\rho c}{8} \int_{|\omega| < \omega_D} \frac{d\omega}{2\pi} |\omega| |h(\omega)|^2 + \int d\tau V(h), \quad (4)$$

where ρ and c are the mass density and sound velocity, respectively, and $h(\omega)$ is the Fourier transform of the bond-length field; the frequency cutoff is given by the Debye frequency ω_D .

The $\rho c|\omega|$ form of the kinetic energy term of the action (4) can be understood by noticing that if the bond length oscillates with frequency ω , then during one oscillation period $2\pi/|\omega|$ this disturbance propagates in the bulk a distance of order $c/|\omega|$. Thus the usual kinetic energy density, proportional to $\rho\omega^2$ should be multiplied by the size of the region $c/|\omega|$ affected by the motion.

The calculation of the properties of the edge of a Luttinger liquid proceeds through the application to the action (4) of Feynman’s variational principle [13] which states that for any trial action S_0 with associated ground-state energy E_0 , the system’s true ground-state energy is bounded above by $E_0 + (T/\hbar) \langle S - S_0 \rangle_0 >_0$ where the zero-temperature limit $T = 0$ is taken at the end and $\langle \dots \rangle_0$ denotes an expectation value computed with S_0 .

Similar to the bulk problem [10] the trial action is selected in the Gaussian form

$$S_0 = \frac{\rho c}{8} \left(\int_{|\omega| \leq \omega_D} \frac{d\omega}{2\pi} |\omega| |h(\omega)|^2 + \gamma \omega_D \int d\tau (h - Ql)^2 \right), \quad (5)$$

where dimensionless variational parameters Q and γ have a meaning of the bond length and its stiffness, respectively. Then the root-mean-square (rms) fluctuation of the bond length can be computed as

$$\langle f^2 \rangle_0^{1/2} = 2\lambda^{1/2} \ln^{1/2}(1 + \gamma^{-1}), \quad (6)$$

where

$$\lambda = \frac{\hbar}{\pi \rho c l^2} \quad (7)$$

quantifies the strength of zero-point motion in the Luttinger liquid. The binding energy of the edge particle

E_{edge} is approximated by $E_0 + (T/\hbar) \langle S - S_0 \rangle_0 >_0$, i. e. by its upper bound:

$$E_{edge}(\gamma, Q) = (\pi \lambda_0^2 / \lambda) \ln(1 + \gamma) - 2e^{Q_0 - Q} (1 + \gamma^{-1})^{2\lambda} + e^{2(Q_0 - Q)} (1 + \gamma^{-1})^{8\lambda} \quad (8)$$

Minimizing E_{edge} with respect to Q we arrive at the expression for the quantum expansion of the edge bond

$$Q_{edge} - Q_0 = 6\lambda \ln(1 + \gamma^{-1}) \quad (9)$$

Substituting this back into (8), E_{edge} can be written as

$$E_{edge}(\gamma) = (\pi \lambda_0^2 / \lambda) \ln(1 + \gamma) - (1 + \gamma^{-1})^{-4\lambda} \quad (10)$$

Minimizing Eq.(10) with respect to γ , and substituting the outcome back into (10) we find

$$\gamma = (4\lambda^2 / \pi \lambda_0^2) (1 + \gamma^{-1})^{-4\lambda} \quad (11)$$

and

$$E_{edge} = (\pi \lambda_0^2 / 4\lambda^2) (4\lambda \ln(1 + \gamma) - \gamma) \quad (12)$$

respectively. The results (6), (9), (11), and (12) should be compared with their bulk counterparts [10]:

$$\langle f^2 \rangle_0^{1/2} = (2\lambda)^{1/2} \ln^{1/2}(1 + \pi/2), \quad (13)$$

$$Q_{bulk} - Q_0 = 3\lambda \ln(1 + \pi/2), \quad (14)$$

$$\lambda_0 = \lambda(1 + \pi/2)^{-\lambda} \quad (15)$$

$$E_{bulk} = (1 + \pi/2)^{-2\lambda} (\pi \lambda \ln(1 + 2/\pi) - 1) \quad (16)$$

Substituting Eq.(15) back in Eqs.(11) and (12) brings them into a form convenient for analysis

$$\gamma = (4/\pi)(1 + \pi/2)^{2\lambda} (1 + \gamma^{-1})^{-4\lambda} \quad (17)$$

$$E_{edge} = (\pi/4)(1 + \pi/2)^{-2\lambda} (4\lambda \ln(1 + \gamma) - \gamma) \quad (18)$$

The properties of the edge as a function of the quantum parameter λ (7) can be computed by finding a solution $\gamma(\lambda)$ to Eq.(17) minimizing the energy (18) and substituting the outcome in the expressions for the rms fluctuation (6) and quantum expansion (9); the dependence on De Boer’s number (2) follows from Eq.(15).

In the classical limit, $\lambda \rightarrow 0$, the only solution to (17) is $\gamma = 4/\pi$ with the energy (18) $E_{edge} = -1$ as expected. As the degree of zero-point motion intensifies (λ increases), the bond stiffness γ decreases and the energy E_{edge} increases. For finite λ Eq.(17) may have more than one solution. One of them is always $\gamma = 0$ corresponding to the delocalized edge particle. For large λ this solution must correspond to the lowest (zero) energy (18).

For $\gamma \ll 1$ the right-hand-side of Eq.(17) behaves as $\gamma^{4\lambda}$ while for $\gamma \rightarrow \infty$ it approaches a γ -independent limit, thus implying that (17) cannot have more than three solutions and that $\lambda = 1/4$ plays a special role.

For $0 < \lambda \leq 1/4$ Eq.(17) has two solutions and the larger of them (whose $\lambda = 0$ limit is $\gamma = 4/\pi$) corresponds to the lowest energy (18). For $\lambda = 1/4$ the explicit solution to (17) is $\gamma = (4/\pi)(1 + \pi/2)^{1/2} - 1 \simeq 1.0415$.

As λ increases beyond $1/4$, Eq.(17) acquires a third root whose $\lambda \rightarrow 1/4 + 0$ limit is $\gamma = (\pi/4(1 + \pi/2)^{1/2})^{1/(4\lambda-1)} \rightarrow 0$. However this solution leads to a larger energy (18) than even the delocalized solution $\gamma = 0$. The lowest energy (bound) state continues to be described by the largest solution to (17).

As λ continues to increase, the finite solutions to (17) approach each other and at some λ they coalesce. This is a critical phenomenon corresponding to the limit of stability of the bound edge. At that point the slopes of the right- and left-hand-sides of Eq.(17) coincide which leads to the limiting values $\gamma \simeq 0.4920$ and $\lambda \simeq 0.3730$ satisfying the relationship $\gamma = 4\lambda - 1$. At larger values of λ Eq.(17) has only one solution $\gamma = 0$ corresponding to an unbound edge. The transition between the bound and unbound states actually happens before the limit of stability is reached, namely when the energy (18) vanishes. Numerical analysis shows that it happens at $\lambda \simeq 0.3412$. This is close to the limit of stability thus implying that the edge delocalization is a weak first-order transition.

The results of the analysis are summarized in Fig. 1 where we show the bond stiffness γ and the edge binding energy E_{edge} as functions of the quantum parameter λ . The metastability develops in the $1/4 \leq \lambda \leq 0.3730$ range: for $\lambda < 0.3412$ the bound edge has lower energy while for $\lambda > 0.3412$ the ground state corresponds to a delocalized edge particle.

These conclusions should be contrasted with the properties of the bulk Luttinger liquid. Its range of existence is given by [10] $0 < \lambda \leq 1.0591$ (or equivalently $0 < \lambda_0 \leq 0.3896$) which is the condition that a solution $\lambda(\lambda_0)$ to Eq.(15) can be found for given De Boer's number λ_0 (2). Therefore in the $0.3730 < \lambda \leq 1.0591$ range the bulk Luttinger liquid is stable against the disordering effect of quantum fluctuations while the edge is not. This is due to the stronger softening effect that zero-point motion has on the free edge as compared to the bulk of the system. The direct evidence of this is presented in Fig. 2 where we show the quantum expansion and rms fluctuation of the bulk and edge bonds as functions of the quantum parameter λ (7) within their corresponding ranges of existence. The quantum expansion in the bulk (14) is a linear function of λ while the edge bond expands faster than linearly because the bond stiffness γ entering the argument of the logarithm in (9) is a decreasing function of λ as shown in Fig. 1. Since the $\gamma(\lambda)$ dependence is not very strong one can say that the edge expansion is roughly twice the bulk value as suggested by the ratio of




Figure 1: Dimensionless stiffness of the edge bond γ and corresponding binding energy of the edge particle E_{edge} of a half-infinite Luttinger liquid as functions of the quantum parameter λ (7). The region of metastability is confined to the $1/4 \leq \lambda \leq 0.3730$ range.




Figure 2: (Color online) Quantum expansion $Q - Q_0$ and rms fluctuation of the bulk and edge bonds as functions of the quantum parameter λ (7). The rms fluctuation is shown both as the vertical extent of shaded regions centered around the quantum expansion curves, and explicitly in the inset.

pre-logarithmic factors in Eqs.(9) and (14). This can be understood by noticing that any bulk bond joins two half-infinite Luttinger liquids thus implying that its dynamics is twice as inertial as that of the edge. In this sense zero-point motion at the edge is about twice as strong as that in the bulk. The same argument explains why the edge rms fluctuation is roughly square-root of two larger than its bulk counterpart (compare Eqs.(6) and (13)).

In describing the dynamics of the edge bond the rest of the system was approximated by a harmonic liquid with the bulk properties which means that the bond adjacent to the edge has the length and rms fluctuation identical to those in the bulk. This is an artifact and in reality, as




Figure 3: The dependences of the energy per particle for various bulk phases of the system on De Boer's number λ_0 (2) together with edge binding energy. The arrow pointing down is the dimer dissociation threshold.

one goes inside the bulk, the bond lengths and their rms fluctuations decrease approaching the bulk values asymptotically. This deficiency would be acceptable provided the calculated length of the edge bond and its rms fluctuation are not very different from their bulk counterparts. Since for molecular matter with pair interaction potential of the Morse form the classical bond length satisfies the condition $Q_0 \gtrsim 5$ [10], inspection of Fig. 2 shows that even at the limit of its stability the length of the edge bond and its rms fluctuation do not exceed their bulk counterparts by more than an acceptable 25%.

Moreover, the relative fluctuation $\langle f^2 \rangle_0^{1/2} / Q_{edge}$ is always significantly smaller than unity which implies that our conclusions are weakly sensitive to the statistics of the underlying particles and that the deficiencies of the Morse potential in mimicking the true pair interaction at largest and shortest distances are ignorable. The latter allows us to argue that the edge dissociation pre-empting the bulk instability is a general property of one-dimensional molecular matter.

In order to gain an insight into the consequences of this effect in Fig. 3 we plot the ground-state energy per particle of the bulk Luttinger liquid (given by Eqs.(15) and (16)) and the binding energy of the edge particle (determined through Eqs.(15), (17), and (18) as functions of De Boer's number λ_0 (2). Additionally we show the ground-state energy per particle for an infinitely diluted gas of Morse dimers, $E_{dimer}(\lambda_0) = -(1/2)(1 - \pi\lambda_0/\sqrt{2})^2$ [9]. The bold parts of the curves describe the ground states of the bulk matter: as De Boer's number increases, at $\lambda_0 \simeq 0.3365$ the Luttinger liquid evaporates via a discontinuous transition into a gas of dimers followed by a continuous dissociation transition at $\lambda_0 = \sqrt{2}/\pi$ into a monoatomic gas [10]. For a system with a free edge the binding energy of the edge particle E_{edge} can become smaller than its dimer counterpart E_{dimer} : for

$\lambda_0 \gtrsim 0.1981$ the whole Luttinger liquid comes unraveled, *two* particles at a time despite the fact that the bulk condensed state is energetically favorable. Since our bulk and edge binding energies are variational upper bounds, in actuality the dimer gas may not come into play; its role then will be played by the monoatomic gas.

If the escape of the edge particles to infinity is impossible due to a distant obstacle, this will generate a vapor pressure and the bulk Luttinger liquid may coexist with a gas of particles. As λ_0 increases toward the point of the bulk transition, dissociation proceeds inside the bulk in a manner similar to that in surface melting [2]. We hasten to mention the speculative character of the statements of this paragraph which we plan to clarify in the future.

The examples of one-dimensional matter with dissociated edge and stable bulk, $0.1981 < \lambda_0 < 0.3365$, include H_2 and D_2 in free space, and more cases can be found in the presence of a medium [10].

This work was supported by the Chemical Sciences, Geosciences and Biosciences Division, Office of Basic Energy Sciences, Office of Science, U. S. Department of Energy.

-
- [1] P. W. Anderson, *Basic Notions of Condensed Matter Physics* (Addison-Wesley Reading, Massachusetts, 1997), Section IIID.
 - [2] For nice reviews see J. G. Dash, *Rev. Mod. Phys.* **71**, 1737 (1999) and U. Tartaglino, T. Zykova-Timan, F. Ercoleassi, and E. Tosatti, *cond-mat/0504680*.
 - [3] H. A. Fertig, R. Côté, A. H. MacDonald, and S. Das Sarma, *Phys. Rev. Lett.* **69**, 816 (1992); R. Côté and H. A. Fertig, *Phys. Rev. B* **48**, 10955 (1993).
 - [4] S. Iijima, *Nature (London)*, **354**, 56 (1991).
 - [5] W. Teizer, R. B. Hallock, E. Dujardin, and T. W. Ebbesen, *Phys. Rev. Lett.* **82**, 5305 (1999), and references therein.
 - [6] J. N. Crain and D. T. Pierce, *Science*, **307**, 703 (2005), and references therein.
 - [7] N. W. Ashcroft and N. D. Mermin, *Solid State Physics* (Saunders College Publishing, 1976).
 - [8] J. De Boer, *Physica* **14**, 139 (1948); J. De Boer and B. S. Blaisse, *Physica* **14**, 149(1948); J. de Boer and R. J. Lunbeck, *Physica* **14**, 520 (1948).
 - [9] P. M. Morse, *Phys. Rev.* **34**, 57 (1929).
 - [10] E. B. Kolomeisky, X. Qi, and M. Timmins, *Phys. Rev. B* **67**, 165407 (2003), and references therein.
 - [11] V. N. Popov, *Teor. Mat. Fiz.* **11**, 354 (1972) [*Theor. Math. Phys.* **11**, 65 (1972)]; K.B. Efetov and A. I. Larkin, *Zh. Eksp. Teor. Fiz.* **69**, 764 (1975) [*Sov. Phys. JETP* **43**, 390 (1976)]; F. D. M. Haldane, *Phys. Rev. Lett.* **45**, 1358 (1980); **47**, 1840 (1981); *J. Phys. C* **14**, 2585 (1981); *Phys. Lett.* **81A**, 153 (1981).
 - [12] A. O. Caldeira and A. J. Leggett, *Phys. Rev. Lett.* **46**, 211 (1981).
 - [13] R. P. Feynman, *Statistical Mechanics* (Benjamin, Boston, 1972), p. 67.