

Minigap in a SN junction with paramagnetic impurities

B. Crouzy,¹ E. Bascones,² and D. A. Ivanov¹

¹*Ecole Polytechnique Fédérale de Lausanne (EPFL),*

Institute for Theoretical Physics, CH-1015 Lausanne, Switzerland

²*Instituto de Ciencia de Materiales de Madrid, CSIC, Cantoblanco, E-28049 Madrid, Spain*

Theoretische Physik, ETH-Hönggerberg, CH-8093 Zürich, Switzerland

(Dated: March 23, 2022)

We study the effect of spin-flip scattering on the density of states in a long diffusive S-N junction with transparent interface. We calculate the critical value of the spin-flip scattering rate at which the minigap closes and give the dependence of the minigap on the spin-flip rate. For the system we consider, the minigap and the critical spin-flip rate have the scale of the Thouless energy, and not of the superconducting gap.

Proximity structures made of superconducting and non-superconducting elements in contact with each other is a topic of current theoretical and experimental interest. Most of the experimental applications concern diffusive materials for which an appropriate formalism has been derived.¹ In the diffusive regime the motion of the electrons is governed by frequent scattering on impurity atoms: the elastic mean free path l_e is much smaller than the relevant length scales of the system, namely the superconducting coherence length ξ and the dimension of the normal (N) and superconducting (S) electrodes. As a consequence of Andreev reflection and its interplay with disorder, a gap is opened in the spectrum of the N-region (see for example^{2,3,4,5} and references therein). This minigap does not depend on the position in the junction and is of the order of magnitude of the Thouless energy $E_{Th} = \frac{D}{L^2}$, where $D = \frac{v_F l_e}{3}$ is the diffusion constant and L is the length of the normal wire. Here and in the following, we use the units with $\hbar = 1$.

Elastic scattering on impurities is not harmful for isotropic superconductivity and is directly connected to the appearance of the minigap. On the contrary, scattering on magnetic impurities leads to the destruction of singlet superconductivity. The exchange interaction between the spin of the electrons and the magnetic impurities breaks the time reversal symmetry between the electrons in Cooper pairs and reduces the superconducting correlations. Scattering on magnetic impurities can be described in terms of a spin-flip scattering rate Γ_{sf} . In the Born approximation^{6,7} Γ_{sf} is proportional to the impurity concentration. The energy gap E_g in the electron density of states (DoS) of a bulk superconductor is lowered by spin-flip scattering

$$E_g = \Delta \left[1 - \left(\frac{2\Gamma_{sf}}{\Delta} \right)^{2/3} \right]^{3/2} \quad (1)$$

and closes for the critical concentration of magnetic impurities⁸ $\Gamma_{sf}^{bulk} = \frac{\Delta}{2}$. Here, the order parameter Δ itself depends on Γ_{sf} .

In the “dirty” limit, the equations of motion for the quasiclassical matrix Green function (averaged over the fast Fermi oscillations and the momentum directions) can

be reduced to the Usadel equations.^{1,9} Usadel equations have been very useful in the analysis of position dependent problems like proximity effect. Spin-flip and inelastic scattering processes can be included in the Usadel equations through the corresponding scattering rates Γ_{sf} , respectively Γ_{in} .¹⁰

The impact of spin-flip scattering on the proximity effect and the minigap can be controlled by selectively doping the normal and/or superconducting regions with magnetic impurities or applying a magnetic field to a thin film or wire.^{7,11} Belzig *et al*¹² have numerically shown that the minigap is reduced and vanishes for large values of Γ_{sf} . For the case of a junction with $L = 1.1 \xi$ (corresponding to the Thouless energy of the order of Δ), they find that the minigap closes when $\Gamma_{sf} \approx 0.4 \Delta$. Other works^{13,14} have analyzed the effect of magnetic impurities in the transport properties of S-N junctions, finding that the Thouless energy is the scale which controls the effect of spin flip scattering on the proximity effect. Due to the self-consistent treatment and the non-linearity of the Usadel equations the complete numerical solution of the problem is difficult. It is important to obtain simple expressions which allow comparison with experiment without solving the full equations for each set of parameters.

In this note we obtain such expressions for the case of a long junction. We show that a finite, but small Γ_{sf} suppresses linearly the minigap which closes at $\Gamma_{sf}^c \approx 0.62 E_{Th}$. We also present an asymptotic expression for the dependence of the minigap on the spin-flip rate near the critical value. Above this value, we calculate the zero energy density of states.

We consider a finite size normal metal N, of length L connected to a semi-infinite superconducting terminal S by a transparent interface. We assume that electronic motion is diffusive in both the normal and superconducting parts. We restrict our discussion to a quasi one-dimensional geometry/wire or to a thin film and neglect any dependence on transversal coordinates. The origin of the coordinate x is fixed at the S-N interface. Introducing the angular parametrization² $g^R = \cos \theta$, $f^R = \sin \theta$ for the normal, respectively the anomalous component of the retarded Green function, the Usadel equations in the

N-region, where the pairing interaction vanishes, become

$$\frac{D\partial_x^2\theta}{2} + (i\epsilon - \Gamma_{\text{in}})\sin\theta - 2\Gamma_{\text{sf}}\cos\theta\sin\theta = 0. \quad (2)$$

The proximity angle θ is a function of the energy ϵ and the position x . The electron DoS, in units of the normal state bulk value $\nu_0 = \frac{m\mu_f}{2\pi^2}$, is given by

$$\frac{\nu(\epsilon, x)}{\nu_0} = \text{Re}[\cos\theta(\epsilon, x)]. \quad (3)$$

The boundary conditions for the quasiclassical equations have been discussed in Refs. 15,16. At the interface with vacuum, the conservation of the quasiparticle current yields

$$\partial_x\theta(x=L) = 0. \quad (4)$$

We will study these equations analytically using simplified boundary conditions at the S-N interface, where we impose the superconducting bulk value (at zero energy) of the proximity angle

$$\theta(x=0) = \frac{\pi}{2}. \quad (5)$$

This boundary condition is justified for energies much smaller than the superconducting order parameter Δ and if the normal part is much more disordered than the superconducting part.⁵

Since the scale of the superconducting order parameter Δ does not appear in the rigid boundary condition (5), we will write in the following the energies and the length in units of the only other relevant scale for our system: the Thouless energy E_{Th} , respectively the width of the N-region L .

The boundary conditions (4,5) and the calculations presented here for the S-N junction can also be applied to describe S-N-S junctions with no phase difference between the superconducting terminals.¹⁷ It is important to pay attention to the choice of the energy scale E_{Th} . A S-N-S junction of length unity is equivalent to a S-N junction of length $\frac{1}{2}$. Therefore all the energies must be multiplied by a factor four if we consider S-N-S junctions.

In the limit of small energies and scattering rates the Usadel equation (2) can be linearized in terms of the deviation $\delta\theta$ from the value of the proximity angle at the S-N interface (5). At this level of approximation the DoS in the N-region is given by

$$\frac{\nu(\epsilon, x)}{\nu_0} \approx \Gamma_{\text{in}}^{\text{eff}} x(2-x), \quad (6)$$

where $\Gamma_{\text{in}}^{\text{eff}} = \Gamma_{\text{in}}(1 + 2\Gamma_{\text{sf}})$ and $x \in [0, 1]$. We see that if $\Gamma_{\text{in}} = 0$ a small spin-flip scattering does not create states at small energies. If, on the contrary, Γ_{in} is finite, Γ_{sf} modifies the density of states (constant at small energies) induced by the inelastic scattering.




FIG. 1: Numerical gap curve (solid line): comparison with asymptotic expressions (dashed lines).

To study the presence of solutions where the electronic DoS (3) vanishes we focus in the rest of the paper on the situation with $\Gamma_{\text{in}} = 0$. We introduce a new notation for the proximity angle θ below the minigap and write $\theta = \frac{\pi}{2} + i\beta$ with β real. The minigap E_g is, by definition, the maximal energy compatible with a real β and can be obtained^{17,18} using a first integral of (2)

$$\partial_x\beta = 2\sqrt{f(\beta^1) - f(\beta)}, \quad (7)$$

where the superscript 1 denotes the value at $x = 1$ and $f(\beta) = \epsilon \sinh\beta + \Gamma_{\text{sf}} \sinh^2\beta$. Integrating equation (7) over the junction, we get

$$\int_0^{\beta^1} \frac{d\beta}{2\sqrt{f(\beta^1) - f(\beta)}} = 1. \quad (8)$$

Without spin-flip, we recover with this relation the well-known value¹⁷ of the minigap $E_g^0 \approx 0.78$. The critical value of the spin-flip at which the minigap in the DoS closes is (see the discussion in Appendix)

$$\Gamma_{\text{sf}}^c = \frac{\pi^2}{16} \approx 0.62. \quad (9)$$

For energies $\epsilon < E_g^0$ and $\Gamma_{\text{sf}} = 0$ equation (8) is solved for two different values of β^1 . One of these solutions leads to a diverging β^1 when the energy goes to zero and we reject it using a continuity argument. This continuity argument is commonly accepted in the quasiclassical approximation, but the diverging branch may play an important role in the discussion of the presence of a non-zero subgap DoS resulting from mesoscopic fluctuations.¹⁹ Considering a finite spin-flip scattering rate Γ_{sf} , we find that the second branch of the solution no longer diverges at zero energy. If we increase the spin-flip rate up to a critical value the two zero energy solutions merge at Γ_{sf}^c . The critical value can therefore be determined taking the limit $\beta^1 \rightarrow 0$ of the integral in the l.h.s. of equation (8) at zero energy.

The complete dependence of the minigap on the spin-flip rate (Fig. 1) can be obtained by a simple numerical

integration of equation (8). But it is possible to derive the asymptotic form of the gap curve for $\Gamma_{\text{sf}} \rightarrow 0$ and for $\Gamma_{\text{sf}} \rightarrow \Gamma_{\text{sf}}^c$.

In the limit of a small spin-flip rate, we can expand the integrand in (8) in the small parameter $\frac{\Gamma_{\text{sf}}}{\epsilon}$. Denoting $\hat{\beta}^1 \approx 1.421$ the value of the proximity angle corresponding to the zero spin-flip value of the minigap E_g^0 , we find the resulting correction

$$E_g \approx (E_g^0 - C_1 \Gamma_{\text{sf}}), \quad (10)$$

where the coefficient C_1 is given by

$$C_1 = \frac{\int_0^{\hat{\beta}^1} \frac{(\sinh \hat{\beta}^1 + \sinh \beta)}{(\sinh \hat{\beta}^1 - \sinh \beta)^{1/2}} d\beta}{\int_0^{\hat{\beta}^1} \frac{d\beta}{(\sinh \hat{\beta}^1 - \sinh \beta)^{1/2}}} \approx 3.09. \quad (11)$$

The minigap decreases linearly with increasing spin-flip scattering rate. From the magnitude of C_1 , we can see that even a small spin-flip rate strongly affects E_g .

To obtain the analytic behavior close to Γ_{sf}^c is more tricky as all $\delta\Gamma_{\text{sf}} = \Gamma_{\text{sf}}^c - \Gamma_{\text{sf}}$, β and $\frac{\epsilon}{\Gamma_{\text{sf}}}$ are small. Following the procedure detailed in the Appendix we obtain the asymptotic dependence of the minigap near the closing point

$$E_g \approx 2 \left[\frac{2\delta\Gamma_{\text{sf}}}{3} \right]^{3/2}. \quad (12)$$

In Fig. 1, we compare the asymptotics (10) and (12) with the numerical gap curve. Dimensions are reintroduced in the graphs for clarity.

For $\Gamma_{\text{sf}} > \Gamma_{\text{sf}}^c$ the DoS in the N-region is finite at any energy. In this domain the Usadel equation (2) at zero energy has a solution with θ real. Applying again the procedure of the first integral (7), but this time for a real θ , we get

$$\int_{\frac{\pi}{2}}^{\theta(x)} \frac{d\theta}{\sqrt{\cos^2 \theta^1 - \cos^2 \theta}} = -2\sqrt{\Gamma_{\text{sf}}} x. \quad (13)$$

Inverting this elliptic integral, we can find a complete zero energy solution for equation (2). This solution involves the function $\text{dn}(u, k)$, which is one of the Jacobi elliptic functions, defined as inversions of the canonical forms of elliptic integrals (we follow the notations used in Refs. 5, 20)

$$\theta(x) = \arcsin \left[\frac{\sin \theta^1}{\text{dn}(2\sqrt{\Gamma_{\text{sf}}}(x-1), \cos \theta^1)} \right], \quad (14)$$

where θ^1 can be determined imposing the rigid boundary condition (5) at $x = 0$

$$\sin \theta^1 = \text{dn} \left[-2\sqrt{\Gamma_{\text{sf}}}, \cos \theta^1 \right]. \quad (15)$$




FIG. 2: Dependence of the zero energy local DoS, at the open boundary, on the spin-flip scattering rate: asymptotic expression near Γ_{sf}^c (solid line), complete elliptic solution (dashed line) saturating to the normal state bulk value ν_0 and its asymptotics at large Γ_{sf} (dots). The inset shows the dependence on position of the DoS for $\Gamma_{\text{sf}} = E_{\text{Th}}$ and $\Gamma_{\text{sf}} = 5 E_{\text{Th}}$.

In the inset of Fig. 2, we use relation (14) to represent the dependence on position of the DoS in the N-region for two different spin-flip rates. As expected the density of states increases when we move away from the interface. For large spin-flip scattering rates, the elliptic solution (14) saturates to the normal state bulk value $\theta(x) = 0$, everywhere except in a thin domain close to the S-N interface (rigid boundary at $x = 0$).

Near the critical spin-flip rate, we found a square root dependence of the zero energy local DoS on Γ_{sf} . Expanding the integrand in (13) in the small $\cos \theta^1$, we obtain that for $\Gamma_{\text{sf}} \rightarrow \Gamma_{\text{sf}}^c$ the density of states at the interface with vacuum is given by

$$\frac{\nu(\epsilon=0, x=1)}{\nu_0} = \sqrt{2} \left[\frac{\delta\Gamma_{\text{sf}}}{\Gamma_{\text{sf}}^c} \right]^{1/2} - \frac{11}{8\sqrt{2}} \left[\frac{\delta\Gamma_{\text{sf}}}{\Gamma_{\text{sf}}^c} \right]^{3/2} + \dots \quad (16)$$

In the limit of large spin-flip rates, when the density of states approaches the normal state one, the expansion²¹ of the elliptic integral in (13) near $\cos \theta^1 = 1$ leads to an asymptotic expression for the DoS at the interface with vacuum

$$\frac{\nu(\epsilon=0, x=1)}{\nu_0} = 1 - 8e^{-4\sqrt{\Gamma_{\text{sf}}}} + \dots \quad (17)$$

In Fig. 2, we compare the expressions (16,17) for the DoS at the open boundary with values obtained using the complete zero energy elliptic solution (14).

In summary, we have studied the effect of spin-flip scattering on the minigap and zero energy density of states of a normal slab connected to a superconducting electrode. We have obtained analytic expressions for the dependence of the minigap on the scattering rate both for small spin-flip scattering rates and close to the critical value at which the gap closes. This critical value is

controlled by the Thouless energy. For values larger than Γ_{sf}^c we find finite DoS at zero energy with a square root dependence on Γ_{sf} . These results are valid for long S-N or S-N-S junctions with transparent interfaces, at small energies. The results on the minigap are not restricted to the case in which there is no spin-flip scattering in the superconducting regions. The only requirement is that the Thouless energy of the normal part is small compared to the gap in the superconducting one. These results allow easy comparison with experimental measurements.

We thank M. Sigrist for useful discussions and acknowledge financial support from NCCR MaNEP of the Swiss Nationalfonds and the Spanish Science and Education Ministry through Ramón y Cajal contract.

APPENDIX: GAP CURVE CLOSE TO Γ_{sf}^c

In this appendix, we derive the asymptotic form of the minigap curve close to Γ_{sf}^c . We have seen previously that the minigap E_g is the largest energy compatible with equation (8). Introducing

$$\begin{aligned} z &= \sinh \beta + \alpha \\ z_1 &= \sinh \beta^1 + \alpha \end{aligned}$$

$$\text{with } \alpha := \frac{\epsilon}{2\Gamma_{\text{sf}}},$$

we can rewrite equation (8) in the form

$$2\sqrt{\Gamma_{\text{sf}}} = \int_{\alpha}^{z_1} \frac{1}{\sqrt{z_1^2 - z^2}} \cdot \frac{1}{\sqrt{1 + (z - \alpha)^2}} dz. \quad (\text{A.1})$$

The integral in the r.h.s. is a function of z_1 and α , which we denote $Y(z_1, \alpha)$. To determine the minigap, we will

find the maximum value of Y over z_1 for a given value of the parameter α .

The critical spin-flip scattering rate (9) can be obtained setting $\alpha = 0$ and maximizing (A.1). It turns out that $Y(z_1, \alpha)|_{\alpha=0}$ is largest for $z_1 = 0$. The next step is to go to finite energies and expand Y in α . We write

$$Y(z_1, \alpha) \approx Y(z_1, \alpha)|_{\alpha=0} + \frac{\partial Y(z_1, \alpha)}{\partial \alpha}|_{\alpha=0} \alpha, \quad (\text{A.2})$$

where

$$\frac{\partial Y(z_1, \alpha)}{\partial \alpha}|_{\alpha=0} = -\frac{1}{z_1} + \frac{z_1}{1 + z_1^2}. \quad (\text{A.3})$$

For a small α , the maximum of Y is expected to be close to the zero energy value $z_1 = 0$ and the second term $\sim O(z_1)$ can be neglected. The first term in the r.h.s. can be expanded

$$Y(z_1, \alpha)|_{\alpha=0} \approx \int_0^1 \frac{1}{\sqrt{1 - s^2}} \left[1 - \frac{(z_1 s)^2}{2} \right] ds. \quad (\text{A.4})$$

where $s = \frac{z}{z_1}$. Taking the derivative of Y over z_1 , using the expansion (A.2), we find that the maximum of Y is obtained for

$$\hat{z}_1 = \left(\frac{\alpha}{\sqrt{\Gamma_{\text{sf}}^c}} \right)^{1/3}. \quad (\text{A.5})$$

substituting back this result in (A.2) and using the definitions of $\alpha = \frac{\epsilon}{2\Gamma_{\text{sf}}}$ and $Y = 2\sqrt{\Gamma_{\text{sf}}}$ we obtain an expression for $E_g(\Gamma_{\text{sf}})$. Finally we write $\Gamma_{\text{sf}} = \Gamma_{\text{sf}}^c - \delta\Gamma_{\text{sf}}$ and expand in the small $\delta\Gamma_{\text{sf}}$ to get the asymptotic dependence (12).

-
- ¹ K. D. Usadel, Phys. Rev. Lett. **25**, 507 (1970).
 - ² A. A. Golubov and M. Y. Kupriyanov, J. Low Temp. Phys. **70**, 83 (1988).
 - ³ E. Scheer, W. Belzig, Y. Naveh, M. H. Devoret, D. Esteve, and C. Urbina, Phys. Rev. Lett. **86**, 284 (2001).
 - ⁴ G. Rubio-Bollinger, C. de las Heras, E. Bascones, N. Agrait, S. Vieira, and F. Guinea, Phys. Rev. B **67**, 121704 (2003).
 - ⁵ A. Altland, B. D. Simons, and D. Taras-Semchuk, Adv. Phys. **49**, 321 (2000).
 - ⁶ A. Abrikosov, L. Gor'kov, and I. Y. Dzyaloshinskii, *Quantum field theoretical methods in statistical physics* (Pergamon Press, 1965).
 - ⁷ K. Maki, *Superconductivity*, vol. II (Marcel Dekker, Inc., 1966).
 - ⁸ A. A. Abrikosov and L. P. Gor'kov, Zh. Eksp. Teor. Fiz. **39**, 1781 (1960), [Sov. Phys. JETP **12**, 1243 (1961)].
 - ⁹ W. Belzig, F. K. Wilhelm, C. Bruder, G. Schön, and A. D. Zaikin, Superlattices and Microstructures **25**, 1251 (1999).
 - ¹⁰ N. Kopnin, *Theory of nonequilibrium superconductivity* (Oxford University Press, 2001).
 - ¹¹ J. G. Rodrigo, H. Suderow, S. Vieira, E. Bascones, and F. Guinea, J. Phys: Condens. Matter **16**, R1151 (2004).
 - ¹² W. Belzig, C. Bruder, and G. Schön, Phys. Rev. B **54**, 9443 (1996).
 - ¹³ S. Yip, Phys. Rev. B **52**, 15504 (1995).
 - ¹⁴ T. Yokoyama, Y. Tanaka, A. A. Golubov, J. Inoue, and Y. Asano, Phys. Rev. B **71**, 094506 (2005).
 - ¹⁵ M. Y. Kupriyanov and V. F. Lukichev, Zh. Eksp. Teor. Fiz. **94**, 139 (1988), [Sov. Phys. JETP **67**, 1163 (1988)].
 - ¹⁶ A. Zaitsev, Zh. Eksp. Teor. Fiz. **59**, 1015 (1984), [Sov. Phys. JETP **59**, 1015 (1984)].
 - ¹⁷ F. Zhou, P. Charlat, B. Spivak, and B. Pannetier, J. Low Temp. Phys. **110**, 841 (1998).
 - ¹⁸ D. A. Ivanov, R. von Roten, and G. Blatter, Phys. Rev. B **66**, 052507 (2002).
 - ¹⁹ P. M. Ostrovsky, M. A. Skvortsov, and M. V. Feigel'man, Phys. Rev. Lett. **87**, 027002 (2001).
 - ²⁰ D. F. Lawden, *Elliptic Functions and Applications* (Springer, 1989).
 - ²¹ B. C. Carlson and J. L. Gustafson, SIAM J. Math. Anal. **25**, 288 (1994).