

Comment on "Sound Modes broadening in Quasicrystals" and its Peer Review

Gerit Coddens

Laboratoire des Solides Irradiés, Ecole Polytechnique,
F-91128-Palaiseau CEDEX, France

(Dated: today)

Recently de Boissieu et al. proposed an explanation for the broadening of acoustic modes observed in quasicrystals (QC). It is the transcription of a well-known model used for glasses. We raise two fundamental objections against applying it to QC. After the text of the Comment we report the methodology that has been used to thwart the publication of this Comment, which is perfectly valid.

I. THE COMMENT

Recently, de Boissieu et al. [1] proposed a mechanism to explain the broadening of phonons starting from a threshold wavevector in QC, as is systematically observed in experimental data. The mechanism (the coupling of sound waves to a heat wave) is quite general and not new, as it has been known for many years in glasses. There is obviously no doubt that the mechanism itself is sound but we must take issue with the way it is being used through a transcription to the field of quasicrystals. Below, we explain our objections on two levels:

(1) On a general level, phonon broadening is an intrinsic property of quasicrystals, even in a completely harmonic model, such that there is a priori no need for the introduction of an assumption of anharmonicity in the form of a coupling.

(2) On a more detailed level, the authors try to blow new life into a cluster scenario proposed by Janot et al. [2], by using the idea of localized modes on clusters as a basic ingredient for the microscopic realisation of the mechanism. The localized modes on the clusters are this time no longer directly responsible for the broadening, but they are proposed to be *at* modes, that couple to the sound waves. To cite them verbatim: "The building bricks of all QC structures are atomic clusters. These clusters are not mere geometrical constructions but real physical entities responsible for specific features in the QC vibrational spectrum (e.g., responsible for localized modes)". [3]

Hence on two levels assumptions are introduced that are not granted or even not needed, while through the presentation the reader might be left with the impression that the experimental data present evidence in support of these assumptions.

(1) There exists an extensive literature on phonons in QC, e.g. on the Fibonacci chain, based on the transfer matrix method. [4] What transpires from such (rigorous) studies is that the phonon eigenmodes are not at all periodic and even not quasiperiodic. (There exist e.g.

so-called recurrent eigenmodes: As a function of its position in space the amplitude of a mode can exhibit humps around the values a^{-n} , where a is the golden mean, n is the set of integers, and a a constant (length). In between the humps, the amplitude of the mode is never zero but it can be very weak). The simple argument of the non-quasiperiodicity of the eigenmodes shows that their Fourier analysis in terms of wave vectors Q will not show dispersion curves of zero line-width as in crystals, but broader features, and that this broadening is not in energy (as de Boissieu et al. think) but in the wave-vector Q . For the recurrent modes, specialists even wonder if it is a mathematically legitimate to take it for granted that they have a Fourier transform in Q -space, and in the case it is not, what kind of information the neutron data contain about these modes.

These features occur even in models that are perfectly harmonic without any coupling between the modes. Of course calculations in the harmonic approximation on successive approximants can never reveal such broadening, because they give rise to zero linewidths by definition. A key intrinsic difficulty of the problem is thus just being missed by such an approach, but it is from the blind spot inherent to such calculations that one might feel the urge to inject additional assumptions of anharmonicity into the problem under the form of localization or couplings.

It would be cheap to push the present objections away, by arguing that real QCs are not one-dimensional, and that surely there will be a loophole of escape from these objections, when we go to higher dimensions. Such vague arguments would (a) reverse the charge of proof, and (b) contain a tacit denial of the horrendous difficulty of eigenvalue problems (with the correct boundary conditions) on quasiperiodic structures. We may add to this that (c) anharmonicity can be experimentally evidenced by the temperature dependence of the Debye-Waller factor. [5]

When one drops full mathematical rigor, approximate eigenmodes that are periodic or quasiperiodic, leading to zero

approximate linewidths, are only expected to exist in the long-wavelength limit. This correlation between absence of broadening and long wavelengths is confirmed by the experimental observations. But the cluster mechanism proposed by de Boissieu et al. only holds in the limit in which acoustic wavelengths are much larger than a typical cluster size, i.e. the very limit where on the basis of the preceding arguments the broadening of the linewidths is rather expected (and observed) to be minor.

(2) In key positions of their paper (the abstract and the final conclusion) the authors stress the important rôle they propose clusters to play in the microscopic realisation of their mechanism. The authors discuss the relevance of isolated clusters, and present the issue in terms of a unique isolated spherical inclusion in a vibrating medium (e.g. a metallic sphere inside a rubber medium). They dismiss such a scenario with great emphasis, but this does not really clarify the assumptions that underly their paper with respect to the issue if there are isolated clusters in quasicrystals: The scenario chosen to assess this issue is too obviously wrong, and the real issue if there are isolated clusters in quasicrystals is more subtle.

We have to address here a difficult situation of possibly ambiguous terminology, because the authors do indeed introduce a concept of "isolated" clusters, different from the one that might be inferred from their presentation. Let us call the type of isolation evoked by the model of a unique spherical inclusion "type 1" and the type of isolated clusters used by the authors "type 2". The inadequacy of the type-2 isolated cluster model does not hinge on the abundance of clusters in the structure as one might infer from the type-1 model with its unique cluster. It rather consists in tacitly denying the importance of boundary conditions in the set of coupled differential equations that describe the phonon problem (and which are in general expressed in terms of a dynamical matrix).

The point is easily understood as follows. Take e.g. a one-dimensional crystal that is based on the periodic repetition of the small motif LSLLS taken from the Fibonacci chain. The crystal is thus ::LSLLSLSLLSLSLLS::... Based on visual clues we could claim that LSLLS is a cluster, and that the crystal is a dense packing of clusters. The eigenmodes and eigenvalues of an isolated cluster LSLLS are completely different from those of the crystal ::LSLLSLSLLSLSLLS::... In fact, the cluster LSLLS has six discrete ω modes with a certain dynamical form factor, $S(Q)$ that extends throughout reciprocal space. The modes are thus ω due to the finite extension of the cluster in space. The modes of the crystal are completely different: They do not correspond to a few isolated discrete energies, but build a whole dispersion curve and for each eigenmode (i.e. each energy) in the acoustic regime the Q -dependence is Dirac-like in reciprocal space (if we limit ourselves to one Brillouin zone).

Of course, it would be completely inappropriate to claim on the basis of the visual clue that we can discern clusters LSLLS in the crystalline structure that there are ω modes in the crystal, and that there would exist a coupling between sound waves and these ω modes in the crystal. The flaw in such a reasoning is uniquely based on a tacit change of the boundary conditions: It replaces periodic boundary conditions (with a rather smooth variation of the force constants across the "cluster" boundaries) by an abrupt discontinuity at the "surface" of the imaginary cluster.

In other words: Type-1 isolation refers to the absence of similar clusters in the surroundings, the cluster is alone. Type-2 isolation refers to a decoupling of the cluster from the surroundings in terms of the force constants. And of course what is relevant for the phonon problem is not if the cluster is alone (type-1 isolation) but how the clusters couple to their surroundings at their supposed boundaries (type-2 isolation). Even in the example of a unique metallic sphere in a rubber medium, it is type-2 isolation that is physically relevant.

QCs are not periodic, and are subject to other boundary conditions than the ones that prevail in a crystal (see below). But this certainly cannot mean that our example would not be appropriate and that the introduction of "clusters" by the authors would have less of a hidden problem with the prevailing boundary conditions, because the crucial point of our objection lies in the postulated abrupt discontinuity, not in the periodicity. When I am raising an objection against the use of (type 2) isolated clusters, it can thus certainly not imply that I would have missed the passage in the paper where the authors acknowledge that there are no (type 1) isolated clusters in quasicrystals. The statement that there are no (type 1) isolated clusters in QCs (because they are a dense packing of clusters) cannot hide the fact that it is the very use of (type 2) isolated clusters which is the basic ingredient for the microscopic interpretation proposed by the authors.

We could never warn the reader enough against the pitfall that would consist in getting one's attention sidetracked towards the issue if there is convincing evidence for the presence of clusters in QCs or otherwise. That would be certainly an interesting topic in its own right, but rather pointless and misleading in the present discussion, as the issue if there are (type-2) isolated clusters in quasicrystals cannot be replaced by an issue if there are clusters in quasicrystals all together, nor by the issue if clusters are physically meaningful in quasicrystals. The verdict on the latter issues will moreover depend on the context of the application: A possible cluster argument in a problem of stability or electronic properties will be different from the one in a phonon problem.

The authors formulate the statement that clusters are not mere geometrical constructions without any proof as though it would be an obvious truth, and the difficulty that they can overlap is passed under silence. The claim that the origin of the ω modes observed in AFDm can be attributed to a localization on clusters is also put forward without any proof.

To introduce the boundary conditions underlying their cluster assumptions, the authors should have given arguments that there is a discontinuity in the force constants at the surface of these clusters. In certain points on the cluster boundaries, the contrary rather seems to be true, viz. when instead of being isolated clusters overlap, which is often the case. In such points it rather looks as though nothing in the whole set of the atomic forces between pairs of atoms in a QC singles out a cluster as an isolated entity, defined by such a discontinuity. The forces between the atoms inside the clusters are not obviously different from those between an atom of the cluster and a neighbouring atom that lies just outside the cluster (but inside the overlapping cluster of the same type). A few phason jumps can create the illusion that a whole cluster has jumped, which also clearly illustrates the relative arbitrariness of assigning an atom to a cluster and of suggesting that a cluster would be an isolated entity whose existence would be obviously defined by a discontinuity in the atomic forces at its surface.

Mathematically spoken, if a cluster is taken large enough it can even be a covering cluster for the whole QC. One can imagine a crystal that could be depicted as a (periodic) arrangement of physically acceptable, overlapping identical clusters of a certain size, and that would not lead to any localization or broadening. Any attempt to escape from this trivial objection must therefore forcedly end up in a discussion of the global, non-periodic arrangement of the clusters and their overlaps.

Discussing QC problems in terms of clusters rather than atoms, is thus just a kind of renormalization procedure, that merely shifts the intrinsic difficulty of non-periodicity to a different length scale, but does not tackle the difficulty itself. It is a blunt denial of the subtlety and difficulty of the eigenvalue problem to overemphasize the rôle of clusters. We can illustrate this with the Fibonacci chain. It starts with LSLLSLSLSLSLSLSLSLSLSLSLS:::, where we have subdivided the sequence in building bricks LSLLS and LSL. Each of the occurrences :LSL: herein is seen to be followed by LS as both building bricks :LSL: and :LSLLS: begin with LS. Hence the whole sequence can be seen as made from the "covering cluster" LSLLS, whereby we have to allow for overlaps LS, which appear exactly at the positions where we have separated out :LSL:. Similarly we could even consider LSL as a covering cluster (the overlap would then be L). Now, the phonon eigenmodes of the isolated sequences LSL and LSLLS can be calculated (from the corresponding 4×4 and 6×6 dynamical matrices). What does this handful of eigenmodes tell us about the phonons of the Fibonacci chain? Hardly anything! As we pointed out above, even the phonons of the periodic sequences based on LSL or LSLLS do not give us the correct picture, despite the fact that in such sequences the clusters are no longer completely isolated (which would be a completely unrealistic boundary condition) and one at least allows for the point that they are embedded in a larger structure (which completely changes the eigenvalue problem).

The idea of clusters LSL and LSLLS certainly has great eye appeal. One might think at first sight that it must yield great insight in the dynamics of the Fibonacci chain. But as we explained above, all this is mere deception. Already the overlap LSLLSLS of two clusters of the type LSLLS will yield completely different solutions for the eigenvalue problem than LSLLS itself. The same basic objections about the boundary conditions remain perfectly valid in the three-dimensional case, such that the fact that we work on the one-dimensional case does not present a loophole from these objections. All the use of the clusters LSL and LSLLS allows us to do is to rewrite the transfer matrix formalism in terms of matrices that correspond to LSL and LSLLS rather than in terms of the more elementary matrices that correspond to S and L. This illustrates how replacing atoms by clusters is just a renormalization procedure, as we stated. It is an underestimation of the complexity of eigenvalue problems and their boundary conditions (which is global) to suggest that they could be approached locally by focusing one's attention to small building bricks. Putting the bricks together just changes everything.

At least in the present context we can thus state that unless a rigorous proof of the contrary is given, it is wise to adopt cautiously the conservative view point that the rigorous application of the idea of clusters, even if they look physically attractive, has remained limited to just a convenient pictorial shorthand to describe parts of the structure, nothing more. We can appreciate from this discussion how both objections (1) and (2) are linked, in the sense that both are based on a tacit modification of highly sensitive details of an eigenvalue problem, that is very hard to spot. The example of how the recurrent modes completely escape the analysis in terms of periodic approximants, shows to what kinds of catastrophies such lack of rigor can lead. Once again, this concern about rigor should not be misrepresented by saying that I would claim that there are no clusters in quasicrystals, or that clusters could not play a role in quasicrystals, etc...

Without any justification, the localized modes invoked are identified with the flat modes that have been reported in APDMN, and a coupling mechanism between these localized modes and sound waves is proposed. We have two objections to this:

(a) Such an explanation for the Γ modes is just one between several other possibilities. One of the alternatives is documented and can therefore not be ignored: By a scrutiny of the displacement patterns in their numerical simulations Hafner and Kráľci [6] were able to associate the Γ modes with a restriction ("confinement") of the vibrations to disclination lines of atoms that are topologically different from average (e.g. the atoms have a 13-fold coordination, rather than a normal 12-fold one). This has nothing to do with the vibration on a cluster.

(b) The issue if the flat modes are due to a localization on clusters is not open-ended within the present state of knowledge. It can be unambiguously settled. It suffices to check if the structure factor of a flat mode is indeed compatible with the dynamical structure factor of a cluster vibration (as Buchenau has done to prove his model for the dynamics of silica). Although the dynamical structure factors of the flat modes have not been published, it must be straightforward to extract this first-rank information from the authors' already existing data, and a numerical calculation of the vibrational spectrum of a Bergman or a Mackay cluster with realistic force constants, involving typically 33 to 55 atoms, is certainly not unfeasible.

Hence, before one can formulate any possible approach of the type proposed by the authors, it is a peremptory prerequisite that one first proves on the basis of existing data, that (1) the observed structure factors of the flat modes are compatible with an interpretation of these modes in terms of cluster phonons, and (2) that there are anharmonicity within the system, e.g. on the basis of Debye-Waller factor anomalies of which one has proved beyond any doubt that they cannot possibly be attributed to an onset phason hopping. These are necessary but not sufficient, minimal conditions that have to be met. They stand completely free from any theoretical considerations, and therefore add up completely independently to the two main objections outlined in the present Comment.

II. THE PEER REVIEW OF THE COMMENT

We would like to give the reader an inkling about the methods that are used to have valid Comments of this type rejected. I would have preferred to quote the referee reports literally rather than paraphrasing them but it has been pointed out to me that it is illegal to reproduce referee reports literally as this constitutes a violation of copyright. I must state that I feel very uncomfortable about this. First of all, it kind of diminishes my credibility and exposes me to cheap and easy criticism that I am distorting the truth because I would not reproduce what has been written literally. Secondly, when one literally reproduces what has been written, one cannot be accused of being responsible for whatever that might be contained in it and look disgraceful, while when one has to paraphrase it, one becomes subjectively accredited with this responsibility. Thirdly, it is apparently not enough that anonymous peer review exposes people almost defenseless to the huge prejudices that can be inflicted by sham peer review, especially when it becomes systematic if some group has managed to completely invade the horizon of an editorial board. Victims are this way also obstructed from denouncing what they have undergone and making the community aware of it. I must ask the reader to consider how this obligation to mention what has been stated only indirectly, can only result in a down-sized and filtered account of the adversity and the personal attacks I have been faced with.

In a first reply de Boissieu stated that the verbatim quotation I made at the beginning of my Comment: "The building bricks of all QC structures are atomic clusters. These clusters are not mere geometrical constructions but real physical entities responsible for specific features in the QC vibrational spectrum (e.g., responsible for localized modes)" would not be in his paper, and that my Comment would distort the point of view of his paper by making quotations out of context.

He also stated that it was not appropriate to take issue with the fact that his analysis is based on isolated clusters, while on p.5 of his paper he had clearly stated that the picture of one isolated cluster is not adequate for QCs, and that one should rather think of a QC in terms of a dense packing of clusters. The reader should not get confused by this would-be catching me in my own contradictions. The contradictions are entirely from the hand of de Boissieu et al. themselves. Yes, de Boissieu et al. state with great emphasis in the beginning of their paper that they do not use isolated clusters, referring to the example of a metallic sphere in a rubber medium. But, no, this cannot hide that the whole argument is exactly based on an assumed isolation of the clusters. It is just that the isolation at stake is very different from the one suggested by the example of a single metallic sphere in a rubber medium. It is not by sorting two entirely different situations under a same descriptive phrasing that they would become equal. In fact, the problem of isolated clusters is not one of numbers ("one" against "a dense packing"), as one could infer from this reply, but one of mutual overlap. In order to overcome the confusion that could result from this reply, I introduced the definitions and the distinction between type-1 and type-2 isolation in my Comment.

de Boissieu further stated that the scientific content of my comment would be very small, that it contained vague and general views, that were in lack of scientific evidence and that were not supported by recent papers. He stated that I overinterpreted the literature.

Finally, he replaced the true issue of my Comment, viz. that the clusters are not isolated, by another issue, viz. if there are clusters in quasicrystals at all, and then gave a detailed reply on this replacement issue, with several citations. Even if this reply had been entirely correct, it did not address the issues I had raised in my Comment.

With respect to my argument that it is not true that clusters are isolated in the sense of having significantly stronger intra-cluster bonding, de Boissieu stated that he certainly agreed that the existence of clusters is still a matter of debate, but that it has not been proved that the forces are of equal strength throughout the QC and that therefore

my argument had no firm ground. He added that the problem of the energetics of QCs is very complicated and one could not expect the final solutions to be given soon.

With respect to my citation of the work of K Raji and Hafner he stated that this was wrong, because the velocities reported by these authors have an insignificant participation ratio, and because there has been no analysis of their vibration patterns, such that their true nature is still not clear, even if it is clear that they are associated with disclination lines.

It was also stated that electron density measurements on a crystalline approximant of AlReSi indicate a larger bonding character within clusters than in between them. (This argument is clearly not general as can immediately be appreciated from the fact that clusters very often overlap, which is the real issue of my Comment).

It was also stated that Gratias approved the cluster approach, while it very clearly transpires from Gratias' papers that he rather very cautiously considers them as a convenient shorthand to describe the structure.

It succeeds to point out that all this does not address at all the issue I raised, which is that the clusters often overlap. The point is thus not if there are clusters at all in quasicrystals, but that these clusters are not type-2 isolated.

After I had replied to this, the correspondence was sent to two referees. One of them appeared biased to me. But eventually, both of them recommended publication with small modifications. However, I learned later on that one of them wrote a separate note to the editors wherein he/she stated that he/she did not wish to review my manuscript any further, because the manuscript would not be presented in the proper manner. (The "he/she" reflect a terminology that was used by the APS).

When I had adapted my version to these comments, the editors of PRB requested that I should remove the sentence from my paper that very explicitly stated that one should be careful in not being sidetracked towards the fake issue if there are clusters in QCs, as the true issue is if these clusters are isolated in a very specific sense. I had made a citation towards de Boissieu's reply to the editors in this respect. It was argued that I would not have the right to cite correspondence from the peer review process. It was also stated that the editorial board was "positively inclined" to accept my Comment for publication. Judging that the way de Boissieu changed the issues was apt to mislead a many reader, I reformulated the sentence such as to keep its contents but to remove the citation. Then PRB put my paper "on hold" for six months, refusing to give any explanation. They had done that already a first time in the review process. Perhaps I should have understood from this that my Comment was not well considered by the APS itself, and that this gentle use of force was meant to be discouraging enough to make me just give up. Finally after nine months, they sent me all at once a report from a fourth referee. There had very obviously not been the slightest reason to ask advice from a fourth referee.

This referee replaced again the issue if the clusters are isolated by the false issue if there are clusters all together. On this replacement issue he stated that an intense debate was going on within the community, and again developed a whole argumentation about this non-issue, citing the model of closed electronic shells by Janot et al., the fracture experiments by Ebert et al., confirmed by numerical simulations by Roesch et al. and the cluster friction model by Feuerbacher et al. It was also stated that the whole development based on the Fibonacci chain was insignificant, and unsuitable to disprove the assumption that there are clusters in QCs, because real three-dimensional clusters are very different from sections of the Fibonacci chain (e.g. in containing many shells). That the looks of one- and three-dimensional clusters are different may very well be true, but is irrelevant for the real issue, which is not if there are clusters in QCs, but that these clusters are not separated. And the latter issue can equally well be explained on the Fibonacci chain as on a three-dimensional model. It was also stated that as long there was no rigorous proof that there are no clusters in QCs, it would be allowed to assume that clusters are present and to build models on this assumption. Once again, the issue is not if there are clusters all together, but that these clusters are not isolated.

The referee also paraphrased me by stating that I would have claimed that the considerations of de Boissieu are redundant, because phonon broadening is an intrinsic property of QCs. He argued that this might be plausible, but that it would be too simplistic, because the broadening cannot be calculated, while de Boissieu et al. would convincingly explain the experimental data.

I responded more or less as follows (I am forced to paraphrase the exact statements from the report in order not to violate copyright):

I clearly wrote in my Comment that "We could never warn the reader enough against the pitfall that would consist in getting one's attention sidetracked towards the issue if there is convincing evidence for the presence of clusters in QCs or otherwise." This sentence summarizes a number of arguments that are clearly developed in my paper: The issue is not if there are clusters or otherwise. The issue is that de Boissieu et al. ignore the consequences of the crucial fact that the clusters often overlap, and that they therefore are not isolated (in the type-2 sense defined in my paper). (And the issue that the clusters are not isolated is a very different one from the one that de Boissieu dismissed after evoking an isolated metallic sphere in a rubber medium, which of the type-1 defined in my paper). The fact that I define two types of isolation already clearly shows that the issue is not if there are clusters all together in QCs. The issue is that these clusters are not isolated in the type-2 sense. I have a whole discussion of this in terms of boundary conditions within the paper. The referee finds it convenient to ignore this all together, and cites work of Janot et al.,

Ebert et al., Roesch et al. and Feuerbacher et al. as proofs for the existence of clusters while this is totally pointless, as clearly explained in my paper and the sentence I quoted from it above. The referee thus goes resolutely for the pitfall I warned against in my paper, which I formulated because de Boissieu had attempted this elusive move already in his first reply.

The referee builds further on this swap towards a false issue, by stating that the example of the Fibonacci chain, which would take a disproportionate large part of my Comment could not be used to prove the cluster assumption wrong. In my paper, the Fibonacci chain is not being used in order to prove that there would be no clusters in a QC. As I already pointed out above, the issue if there are clusters or otherwise is pointless for our discussion. What matters is that the clusters often overlap (i.e. are not isolated in the type-2 sense) and in order to point this out, the Fibonacci chain is as good as a full-edged 3D model. And the referee further insists on focusing the attention to this pointless issue of the existence of clusters or otherwise when he writes that three-dimensional clusters are comparable to sections of the Fibonacci chain, and that it is legitimate to use the cluster assumption as long as there is no striking argument against it.

The latter contains actually a reversal of de Boissieu's charge of proof: One could make as many unproved claims as one likes, it is up to others to prove them wrong. de Boissieu and the referee know very well that this is not viable. This shines through clearly enough when it comes down to attacking my work, rather than defending the work of de Boissieu, and one tries to make prevail totally irrelevant objections against it to make us wonder if perhaps I did not fail to meet my charge of proof on some very tiny loopholes.

One of these irrelevant objections is that the Fibonacci chain would not be pertinent in the present discussion. The arguments developed on the Fibonacci chain are not affected by his irrelevant distinctions between the Fibonacci chain and 3D QCs she would like us to believe crucial. The Fibonacci chain is perfectly suited for discussing the type-2 isolation and other issues at stake. Making the same points on a 2D or 3D QC would require to include into the paper elaborate Figures to show the clusters and how they overlap, etc... While with the Fibonacci chain one can describe the whole situation exhaustively and very clearly by referring to the letters L and S. One issue of my paper is that the possible overlap of the clusters (the lack of type-2 isolation) is pointing out a tacit cheat about the values of the interaction forces: It tacitly implies that the bonds between atoms within the cluster are stronger than the bonds between atoms of a cluster and surrounding atoms. That this is not true is totally obvious when two clusters A and B overlap, as discussed in my paper. And the referee can check it also on a 3D model. It is nothing specific for the Fibonacci chain only. This entails that the clusters are not at all doing what de Boissieu claims they are doing. This point is exactly the same one as voiced by Henley in his paper "Clusters, phason elasticity, and entropic stabilization: a theory perspective" (Phil. Mag. 86, 1123 (2006): Ames conference proceedings) in the first sentence of the section "2. Clusters" on page 1124. It is this issue, and not the mere presence of clusters or otherwise, that is essential and makes de Boissieu's position untenable. To make this untenable position prevail nevertheless, the referee carefully eludes discussing this crucial point of overlap. He diverts the attention away from it by hammering incessantly on the irrelevant issue if clusters exist all together.

I also addressed the paraphrasing of my argument that the use of clusters would be redundant. In fact, this is a completely false presentation of the issues, as anyone who reads the paper carefully can see. The referee operates a very subtle shift when he paraphrases my objection by stating that I would have claimed the considerations of de Boissieu et al. are redundant. As far as I can see I wrote: "There is a priori no need for the introduction of an assumption of anharmonicity in the form of a coupling." If one thinks carefully about it, this does not mean that the considerations are redundant, but that they could be redundant. The snag to this almost subliminal shift is that if I had claimed that the considerations are redundant I would be invested with a charge to prove it. While if the considerations could be redundant, it is the charge of proof of de Boissieu that has not been properly met. Hence this is a hidden reversal of the charge of proof, that is very hard to spot. That phonon broadening is an intrinsic property is not merely plausible as he states, it is a mathematical certainty, because the eigenmodes are not quasiperiodic. Hence here the referee unduly questions (again in an incohesive looking way) an established obvious factual mathematical truth. That nobody is able to solve the horrendously difficult problem of the calculation of the q-dependence of the intrinsic broadening, does not exclude that the broadening observed could be entirely due to this intrinsic broadening. Let one please not jump onto this sentence to put again things in my mouth that I do not say. I do not say that the broadening is entirely intrinsic, I say that cannot be excluded that it could be entirely intrinsic. The mechanism of intrinsic broadening has at least the merit that it is physically sound, while the cluster scenario is conclusively proved wrong by e.g. the type-2 isolation issue, which the referee carefully eludes to discuss.

It is therefore ridiculous to exploit the difficulty of the problem of calculating the intrinsic broadening to compare this scenario unfavourably with the (illusory) success of the cluster model by stating that the result of the q-dependence of the line width presented by Boissieu et al. is convincing. The model is proved wrong and that the data can be fitted with a polynomial of the fourth degree is hardly informative and astounding that could be derived from scores of other models.

I have never stated that the broadening would be uniquely intrinsic. As I explained it already above I have only

evoked this as a possibility. Because, what the argument of the intrinsic broadening was intended to show is that the assumptions de Boissieu makes are in lack of justification, by giving a counter example. E.g. the tacitly implied assumption that there is anharmonicity is gratuitous. In view of the possibility of intrinsic broadening which will occur even in a completely harmonic model, it is a peremptory prerequisite to prove that the forces are anharmonic before one can introduce the assumptions that underly de Boissieu's model. Asking to develop the intrinsic broadening scenario into a full calculation as the referee does is, again, a reversal of the charge of proof. Moreover it tries to saddle me up with the obligation of a *demonstratio diabolica*.

The editorial board of PRB refused to consider my reply to this referee report. They even refused to state this refusal. They just moved on towards a statement that this ended the review of my Comment. They eluded answering by addressing non-scientific issues, maintained my Comment rejected and even eluded responding to an appeal of mine. They had artificially made the whole procedure last for more than two years. They even recommended that I try to have it published in another journal, because other journals have other criteria for approval than the APS.

We may finally point out that we already had attempted to write a Comment on the artificial cluster issue, back in 1993 when it was introduced in reference [2]. Janot et al. dropped an off-hand comment on my work towards the end of that paper that my interpretation of the quasielastic data in terms of phason hopping would be wrong. In reality, their data did not warrant such questioning of my work. In fact, reference [2] reported a failure to observe the quasielastic scattering that I had measured and that corresponded to the decrease of the elastic intensity when the temperature was raised. Such elastic data can never be used to challenge the interpretation of the much more detailed and specific quasielastic data. Nevertheless, Janot et al. did this, denigrating my work. I had to discover this as an accomplished fact in the published literature. To undo the damage, I was forced to write a Comment to reference [2], with reversed rights of reply. Using this reversed situation, Janot answered that my samples were suspicious and that the quasielastic signal was due to preferential segregation of Cu into the grain boundaries. On the editorial board of Physical Review Letters S. Moss stated that he felt much more sympathetic towards my arguments, but that the exchange would be too long to publish it in Physical Review Letters. S. Moss and R. Schuhmann suggested to me that I send it to Physical Review B. But when I did this, I was told that Physical Review B does not handle Comments on papers published in Physical Review Letters. In my Comment I had pointed out that the cluster scenario was analogous to scenarios used in glasses. But in reference [1] it is stated towards the end, that after the work was finished, the authors discovered that a similar approach had been used in glasses.

After the rejection of my Comment on [2], a proposal of mine for beam time at the ILL to measure phason dynamics was rejected on the basis a statement by Dubois in the scientific evaluation committee that the experiment had already been done by Janot. It was just not true. When I protested, and I expressed my fears that my ideas would be stolen, Janot wrote a letter to me with copy to the director of the ILL, wherein he stated that I would be paranoid, and that they did not intend to measure phason dynamics. A few months later he and de Boissieu made the experiment in my place on IN16, but they melted their sample. They had made their attempt to measure phason dynamics with the same type of sample, on the same type of instrument, in the very same Q -range, with the same energy resolution, and in the same energy and temperature range. Nevertheless, they wrote an ILL report about it wherein they stated that this experiment would be different from mine and wherein they reported that they had figured out in the meantime that the interpretation of the Debye-Waller factor in [2] was wrong. The interpretation of the temperature dependence of the Debye-Waller factor had been the only element of justification on which the whole introduction of the cluster issue had been based. It was wrong. And already at that stage, the obvious error in the reasoning had been that the clusters are not isolated but overlap. Nevertheless, these issues were introduced again in reference [1].

[1] M. de Boissieu, R. Currat, S. Franconal and E. Kats, Phys. Rev. B 69, 054205 (2004).

[2] C. Janot, A. Margl, B. Frick and M. de Boissieu, Phys. Rev. Lett. 71, 871 (1993); C. Janot and M. de Boissieu, Physica B 219-220, 328 (1996).

[3] (a) Our verbatim citation can be found on page 5, at the top of the second column of reference [1]. (b) The authors qualify the optic modes observed in QCs as "dispersionless". As acoustic phonons in the long-wavelength limit are also dispersionless, we prefer the terminology "flat", which not only indicates that the slope is constant with the wave vector Q , but also that this constant is zero.

[4] F. Delyon and J. Periere J. Stat. Phys. 64, 363 (1991); F. Delyon and D. Petritis Comm. Math. Phys. 103, 441 (1986); F. Delyon and B. Souillard Comm. Math. Phys. 89, 415 (1983); A. Suto, in Beyond Quasicrystals, p.481-549 edited by F. Axel and D. Gratias, (Editions de la Physique, Les Ulis).

[5] In AlCuFe e.g. the phonons neatly follow the general scheme of broadening, but there is no evidence that would tally with a universal ansatz of anharmonicity. A change of slope occurs in the temperature dependence of the Debye-Waller factor, but this is entirely due to the onset of phason hopping dynamics (see e.g. G. Coddens, S. Lyonnard Y. Calvayrac, Phys. Rev. Lett. 78, 4209 (1997); see also H. de Araj, A. Gomes, and J.B.M. da Cunha, Solid State Comm. 97, 1025 (1996)).

- [6] J. Hafner and M. Krajić, J. Phys.: Cond. Matter 5, 2489 (1993); M. Krajić and J. Hafner, J. Non-Cryst. Solids, 192-193, 338 (1995).