

Spin transverse force on spin current in an electric field

Shun-Qing Shen

Department of Physics, and Center for Theoretical and Computational Physics,
The University of Hong Kong, Pokfulam Road, Hong Kong, China
(dated: Revised on July 27, 2005)

As a relativistic quantum mechanical effect, it is shown that the electric field exerting a transverse force on an electron spin 1/2 only if the electron is moving. The spin force, analogue to the Lorentz force on an electron charge in a magnetic field, is perpendicular to the electric field and the spin current which spin polarization is projected along the electric field. This spin-dependent force can be used to understand the zitterbewegung of electron wave packet with spin-orbit coupling and is relevant to the generation of the charge Hall effect driven by the spin current in semiconductors.

PACS numbers: 85.75.-d, 72.20.My, 71.10.Ca

Recent years spintronics has become an emerging field because of its potential application to semiconductor industry, and more and more attentions are focused on the effect of spin-orbit coupling in metals and semiconductors.[1] The spin-orbit coupling is a relativistic effect describing the interaction of the electron spin, momentum and electric field, and provides a route to manipulate and to control the quantum spin states via external fields.[2, 3] It is desirable to understand the motion of electron spin with spin-orbit coupling in an electromagnetic field. In electrodynamics it is known that a magnetic field would exert a Lorentz force on an electric charge if it is moving. This Lorentz force can generate a lot of fundamental phenomena such as the Hall effect in solid.[4] The interaction of the spin in the electromagnetic field behaves as if the spin is a gauge charge and the interaction is due to the SU(2) gauge field.[5] It is essential that the electron spin is an intrinsic quantum variable, not a just classical tiny magnetic moment. The physical meaning of the interaction is closely associated to the Aharonov-Casher effect and the Berry phase.

In this paper it is found that an electric field exerts a transverse force on an electron spin if it is moving and the spin is projected along the electric field. The force is proportional to the square of electric field, and the spin current projected along the field, and its direction is always perpendicular to the electric field and the speed of the spin. The force stems from the spin-orbital coupling which can be derived from the Dirac equation for an electron in a potential in the non-relativistic limit or the Kane model with the $k \cdot p$ coupling between the conduction band and valence band. From an exact solution of a single electron with the spin-orbit coupling it is heuristic to understand that the zitterbewegung of electronic wave packet is driven by the spin transverse force on a moving spin. The role of spin transverse force is also discussed in the spin Hall effect and its reciprocal effect driven by the pure spin current in semiconductors.

We start with the Dirac equation of an electron in a confining potential V and a vector potential A for a mag-

netic field, $B = \nabla \times A$,

$$i\hbar \frac{\partial}{\partial t} \psi = \frac{\hbar^2}{2m} \nabla^2 \psi + \frac{e}{c} \mathbf{A} \cdot \nabla \psi + m c^2 \psi + V \psi; \quad (1)$$

where α and β are the 4 \times 4 Dirac matrices. m and e are the electron mass and charge, respectively, and c is the speed of light. We let $\psi = e^{-imc^2 t/\hbar} \psi'$ such that the rest mass energy of electron is removed from the energy eigenvalue of the electron. In the nonrelativistic limit, ψ' is a very small component, $\frac{1}{m c^2} \nabla \cdot \mathbf{p} \psi + \frac{e}{c} \mathbf{A} \cdot \nabla \psi'$ where σ are the Pauli matrices. Thus the component ψ' of the wavefunction satisfies the following equation, $i\hbar \frac{\partial}{\partial t} \psi' = H' \psi'$, where

$$H' = \frac{\mathbf{p} + \frac{e}{c} \mathbf{A}}{2m} + V_{\text{eff}} + \frac{\hbar}{4m^2 c^2} \nabla \cdot \mathbf{p} \nabla \cdot \mathbf{p} + \frac{e}{4m^2 c^2} \mathbf{A} \cdot \nabla \nabla \cdot \mathbf{A} \quad (\nabla \cdot \mathbf{V}), \quad (2)$$

where $B = \nabla \times \mathbf{A}$ and $V_{\text{eff}} = V + \frac{\hbar^2}{8m^2 c^2} \nabla^2 V$. In the last step we neglect the higher order terms of expansion. Thus the Dirac equation is reduced to the Schrodinger equation with the spin-orbit coupling. The same form of effective spin-orbit coupling and Zeeman splitting can be also derived from the 8 \times 8 Kane model that takes into account only the $k \cdot p$ coupling between the Γ_6 conduction band and the Γ_8 and Γ_7 valence bands, although the effective mass, the effective Lande g-factor, and the effective coupling coefficients have to be introduced as material-specific parameters such as $m^* = m$, $B = g_B = 2$, and $\tilde{A}^2 = (4m^2 c^2) \nabla^2 r_{41}^{6c6c}$. [6]

In the Heisenberg picture the kinetic velocity is

$$\mathbf{v} = \frac{1}{i\hbar} [\mathbf{r}; H] = \frac{1}{m} \mathbf{p} + \frac{e}{c} (\mathbf{A} + \mathbf{A}') \quad (3)$$

where $\mathbf{A}' = \frac{\hbar}{4m c^2} \nabla \times \mathbf{V}$ and comes from the spin-orbit coupling. It indicates clearly that \mathbf{A}' plays a role of a SU(2) gauge vector potential. The spin dependence of the gauge field can separates the charged carriers with different spins in cyclotron motion experimentally.[7]

Even though we have $[\mathbf{p}, \mathbf{p}] = 0$ for canonical momentum, the analogous commutators do not vanish for the kinetic velocity

$$[\mathbf{v}, \mathbf{v}] = \frac{i\hbar e}{m^2 c} \mathbf{B} + \frac{e^2}{m^2 c^2} [\mathbf{A}, \mathbf{A}] \quad (4)$$

where the total magnetic field $\mathbf{B} = \mathbf{B} + \mathbf{r} \times \nabla A$; and $\mathbf{r} \times \nabla A = \frac{\hbar}{2m} \nabla^2 V - (\mathbf{r} \cdot \nabla) \mathbf{V} = (4m\alpha e) \mathbf{z}$: Notice that $[\mathbf{A}, \mathbf{A}] = 2\hbar^2 i (\mathbf{r} \cdot \nabla) \otimes \nabla V = (4m\alpha e)^2 \mathbf{z}$: We can derive the quantum mechanical version of the force,

$$m \frac{d\mathbf{v}}{dt} = \mathbf{F}_h + \mathbf{F}_g + \mathbf{F}_f \quad (5)$$

with

$$\mathbf{F}_h = \frac{e}{c} \mathbf{v} \times \mathbf{B} + \mathbf{r} \times (\nabla_{\text{eff}} + \nabla_B) \mathbf{B} \quad (6a)$$

$$\mathbf{F}_g = \frac{\hbar}{2m c^2} [(\mathbf{B} \cdot \mathbf{r} \nabla) \mathbf{B} - (\mathbf{r} \nabla) \mathbf{B}]; \quad (6b)$$

$$\mathbf{F}_f = \frac{\hbar}{8m^2 c^4} (\mathbf{r} \nabla) (\mathbf{v} \cdot \mathbf{r} \nabla) \quad (6c)$$

This is the quantum mechanical analogue of Newton's second law. Of course we should notice that this is just an operator equation. The uncertainty relationship tells that the position and momentum cannot be measured simultaneously, and there is no concept of force in quantum mechanics. To see the physical meaning of the equation, we take the expectation values of both sides with respect to a Heisenberg state $|j\rangle$ which does not vary with time. The expectation values of the observable describe the motion of the center of the wave package of electrons. In this sense we have an equation of the force experienced by the moving electron. Actually the first term \mathbf{F}_h in Eq. (5) is the Lorentz force for a charged particle in a magnetic field $\hbar \mathbf{B}$ which contains the contribution from the SU(2) gauge field \mathbf{A} as well as the conventional electromagnetic field. We have recovered the Ehrenfest theorem as one of the examples of the corresponding principle in quantum mechanics. The term, $\hbar (\mathbf{r} \times \nabla_B) \mathbf{B}$; results from the non-uniform magnetic field. Its role was first realized in the Stern-Gerlach experiment, where a shaped magnet is used to generate a non-uniform magnetic field to split the beam of silver atoms. In the classical limit it is written as the interaction between the magnetic momentum $\mu_B = \hbar \mathbf{B}$ and magnetic field. This spin force depends on the spin. Recently it is proposed that the force can generate a pure spin current if we assume \mathbf{B}_z is a constant.[8] In the term \mathbf{F}_g we can also use \mathbf{z} to replace the spin. It is non-zero only when the electric and magnetic fields coexist, as suggested by Anandan and others.[5] This term will play an essential role in generating spin Hall current in two-dimensional Rashba system, which we will discuss it later.

The last term, \mathbf{F}_f , comes from the SU(2) gauge potential or spin-orbital coupling. As the force is related

to the Planck constant it has no counterpart in classical mechanics and is purely quantum mechanical effect. The force is irrelevant of the magnetic field. In the classical limit we cannot simply use the magnetic momentum to replace the spin in the potential A . Otherwise the force vanishes. To see the physical meaning of the force, we write the spin force for a single electron on a quantum state in a compact form,

$$\hbar \mathbf{F}_f = \frac{e^2 \mathbf{J} \cdot \mathbf{j}}{4m^2 c^4} \mathbf{B} \cdot \mathbf{E} \quad (7)$$

where the spin current is defined conventionally, $\mathbf{j}_s = \frac{\hbar}{4} \nabla \psi$; $\mathbf{E} = \mathbf{J} \times \mathbf{j}$; and in the last step the relation $\mathbf{r} \times \nabla A = \mathbf{z}$ has been used. This is the main result in this paper. The force is proportional to the square of electric field \mathbf{E} and the spin current which polarization is projected along the field. It is important to note that an electron in a spin state perpendicular to the electric field will not experience any force. Comparing with a charged particle in a magnetic field, $\mathbf{j}_c = \mathbf{B}$; where \mathbf{j}_c is a charge current density, the spin force is nonlinear to the electric field and depends on the spin state of electron.

We discuss several examples relevant to the spin force. Though the gauge field \mathbf{A} provides a spin-dependent magnetic field $\mathbf{r} \times \nabla A$, the Lorentz force caused by the field on the charge will vanish in a uniform electric field $\mathbf{r} \cdot \nabla V = e\mathbf{E}$: The spin dependent force \mathbf{F}_g also vanishes in the absence of magnetic field. Here we consider the motion of an electron confined in a two-dimensional plane subjected to a perpendicular electric field,

$$H = \frac{p^2}{2m} + (p_x y - p_y x) \quad (8)$$

where $\mathbf{z} = \hbar \mathbf{E} = (4m^2 c^2) \mathbf{z}$ from Eq.(2). This can be regarded as counterpart of a charged particle in a magnetic field. On the other hand it has the same form of the Rashba coupling in a semiconductor heterojunction with the structural inversion asymmetry,[9] where the spin-orbit coupling is induced by the offsets of valence bands at the interfaces and the structure inversion asymmetry.[6] A typical value of this coefficient is of order $10^{-4} c$ (c the speed of light), and can be adjusted by an external field.[10] Because of the spin-orbit coupling the electron spin will precess with time,

$$\frac{d\langle \mathbf{S} \rangle}{dt} = \frac{2}{\hbar} \langle \mathbf{S} \rangle \times (\mathbf{p} \times \mathbf{z}) \quad (9)$$

Since the momentum \mathbf{p} is a good quantum number, without loss of generality, we take $\mathbf{p} = p_x \hat{x}$; just along the x direction. Correspondingly the wave function in the position space $\langle \mathbf{r} | j \rangle = \exp(ip_x x) \psi_s = L$ where ψ_s is the initial spin state. Equivalently the spin-orbit coupling provides an effective magnetic field along the y direction, $\mathbf{B}_{\text{eff}} = p_x \hat{y}$. This problem can be solved analytically, and the electron spin precesses in the spin

x-z plane,[11] $x(t) = x \cos \omega_c t - z \sin \omega_c t$; $z(t) = z \cos \omega_c t + x \sin \omega_c t$; and $y(t) = y$ where the Larmor frequency $\omega_c = 2p_x / \hbar$. The spin $\mathbf{z}(t)$ varies with time and the spin current is always along the x direction, $\langle \mathbf{h}_s^x \rangle = \frac{\hbar}{2m} (\langle \mathbf{z} \rangle \cos \omega_c t + \langle \mathbf{x} \rangle \sin \omega_c t) \hat{x}$ where $\langle \cdot \rangle$ means the expectation value over an initial state $\langle \mathbf{h}_s^x \rangle_i$. As a result the spin transverse force on the spin is always perpendicular to the x direction. Correspondingly the kinetic velocity v_x and v_y at a time t are

$$\hbar v_x i_t = \frac{p_x}{m} + \langle \mathbf{h}_s^y \rangle_i; \quad (10a)$$

$$\hbar v_y i_t = \langle \mathbf{h}_s^x \rangle_i \cos \omega_c t - \langle \mathbf{z} \rangle_i \sin \omega_c t; \quad (10b)$$

respectively. Though $p_y = 0$ the kinetic velocity $\hbar v_y i_t$ oscillates with the frequency ω_c while $\hbar v_x i_t$ remains constant. The y-component of the position is

$$\hbar y i_t = \hbar y i_{t=0} + \frac{\hbar}{p_x} \sin \frac{\omega_c t}{2} \langle \mathbf{h}_s^x \rangle_i \cos \frac{\omega_c t}{2} - \langle \mathbf{z} \rangle_i \sin \frac{\omega_c t}{2}; \quad (11)$$

If the initial state is polarized along y direction the electron spin does not vary with time as it is an energy eigen state of the system, as discussed by Datta and Das.[2] In this case the spin current $\langle \mathbf{h}_s^x \rangle_i$ carried by the electron is always zero and the spin transverse force is zero. Thus $\hbar v_y i_t = 0$: If the initial state is along the spin z direction at $t = 0$, i.e., $\langle \mathbf{z} \rangle_i = s = \pm 1$; it is found that $\hbar v_y i_{t;s} = s \sin \omega_c t$: Different spins will move in opposite directions. It can be understood that the spin precession makes the spin current which polarization is projected along the electric field changes with time such that the spin force along the y direction also oscillates with the frequency ω_c . This force will generate a non-zero velocity of electron oscillating along the y direction. Though $\hbar v_y i_{t;s=1} - \hbar v_y i_{t;s=-1} = 2 \sin \omega_c t$; the velocity does not contribute to the spin current along the y direction, i.e., $\langle \mathbf{h}_s^y \rangle_i = \langle \mathbf{g}_i^y \rangle_i = 0$: The trajectory oscillates with the time. The amplitude of the oscillation is $\sim p_x$ and the frequency is $\omega_c = 2p_x / \hbar$: For a typical two dimensional electron gas the electron density is $n_e = 10^{11} - 10^{12} \text{ cm}^{-2}$ and the wave length near the Fermi surface $\sim p_x / \hbar \approx 10 \text{ nm}$. For a typical Rashba coupling $\alpha = 10^{-4} \text{ eV} \cdot \text{Å}$, $\omega_c = 0.3 - 1.0 \times 10^{14} \text{ s}^{-1}$: The rapid oscillation of the electron wave packet is known in literatures as the zitterbewegung of electron as a relativistic effect, which is usually regarded as a result of admixture of the positron state in electron wave packet as a relativistic effect.[12] In semiconductors the Rashba coupling reflects the admixture of the particle and hole states in the conduction and valence bands. Recently Schliemann et al obtained the solution of the trajectory and proposed that this effect can be observed in a III-V zinc-blende semiconductor quantum wells.[13] In the p-doped semiconductors described by the Luttinger model there exists also a spin force, and will generate the zitterbewegung as calculated by Jiang et al.[14] Though the spin




FIG. 1: The electric field exerts opposite forces on electrons with different spins polarized along the field.

transverse force on a moving spin is very analogous to the Lorentz force on a moving charge, because of spin precession, its effect is completely different with the motion of a charged particle in a magnetic field, where the amplitude of the Lorentz force is constant and the charged particle will move in a circle. The zitterbewegung of the electronic wave package near the boundary will cause some edge effect as shown in recent numerical calculations.[15] The edge effect is determined by the electron momentum. The smaller the momentum, the larger the edge effect. The amplitude and frequency of the zitterbewegung satisfy a relation that $(\sim 2p_x) (2p_x / \hbar) = \omega_c$, which is the amplitude of oscillation of the velocity v_y . In Fig.1 it is illustrated that for two electrons with different spins experience opposite forces in an electric field.

Furthermore we consider a two-dimensional electron gas lacking both the bulk and structural inversion symmetries. A Dresselhaus term $(p_x \times p_y \times y)$ will be included in the total Hamiltonian in Eq. (8).[16] In this model the spin force formula gives[17]

$$\hbar \mathbf{F}_s = \frac{4m^2}{\hbar^2} (\mathbf{p}^2 - p_x^2) (\mathbf{j}_s^z - \hat{z}) \quad (12)$$

for each moving electron. First of all the force disappears at the symmetric point of $\mathbf{p} = 0$: At this point the operator $\mathbf{p} \times \mathbf{y}$ is a good quantum number and there is no spin flip in the system. For $\mathbf{p} \neq 0$; the moving electron will experience a spin-dependent force and the force will change its sign near $\mathbf{p} = 0$. A heuristic picture from this formula is that when a non-zero spin current \mathbf{j}_s^z goes through this system the spin-orbit coupling exerts the spin transverse force on the spin current, and drives electrons to form a charge Hall current perpendicular to the spin current. The injected spin current can be generated in various ways, such as by the spin force $\mathbf{r} \times (\mathbf{B} \times \mathbf{v})$ [8] and circularly or linearly polarized light injection.[18, 19] For instance we assume the spin current \mathbf{j}_s^z be generated by the linear polarized light injection which is proportional to the transition rate of from the valence band to the conduction band with finite momentum and the lifetime of electrons at the excited states. In the relaxation time approximation in a steady state the drift velocity

orthogonal to the spin current is $h\nu_y i = \frac{4m}{\hbar^2} (\frac{\hbar^2}{2} - \frac{\hbar^2}{2}) j_y^z$ if \hbar is not so long, i.e. $2\mu_B \frac{\hbar^2}{2} + \frac{\hbar^2}{2} \ll \hbar$: This non-zero drift velocity will form a Hall current orthogonal to the spin current. This is the charge Hall effect driven by the spin current. In ferromagnetic metal or diluted magnetic semiconductors the charge current is spin polarized it can generate the spin polarized Hall current via the spin transverse force. Thus the spin transverse force can be also regarded a driven force of an anomalous Hall effect [20] and the spin-resolved Hall effect. [17, 21] A detailed calculation for this Hall conductance is given by the Kubo formula as a linear response to the field $B = B_x \hat{x}$. This field will generate a spin force, $r(g_B B_x) = g_B B_x \hat{x}$; which will circulate a spin current along the x direction, and furthermore the spin-orbit coupling provides a driving force to generate a transverse charge current, j_{cy} . In the clear limit the Hall conductance $\kappa_{xy} = j_{cy}/j_{cy} = (g_B B_x) = 0$ for $\hbar = 0$ and $(\hbar = 4\hbar) (\frac{\hbar^2}{2} - \frac{\hbar^2}{2}) = \frac{\hbar^2}{2}$. However following Inoue et al and Mishchenko et al [22] the inclusion of impurities scattering will suppress the Hall conductance completely just like the spin Hall effect. On the other hand numerical calculation in mesoscopic systems shows the existence of the effect. [19] Another example is the two-dimensional p -doped system with the cubic Rashba coupling, [23]

$$H = \frac{p^2}{2m} + i(p_+^3 - p_-^3) \quad (13)$$

where p_{\pm} are spin increasing and decreasing operators, and $p = p_x - ip_y$. The spin force on the moving electron in this system is $F_f = (2m p_{\pm} = \hbar)^2 (\frac{\hbar^2}{2} - \frac{\hbar^2}{2})$: The linear response theory gives the Hall conductance $\kappa_{xy} = (2\hbar)^{-1}$ which is robust against the vertex correction from impurities scattering. Calculations by means of the Green-Keldysh function technique and linear response theory [8, 19] show the existence of charge Hall effect driven by the spin current, and the Onsager relation between the charge Hall effect and its reciprocal. The key features of the numerical results are in good agreement with the picture of spin force qualitatively.

In conclusion, an electric field exerts a transverse force on a moving spin just like a magnetic field exerts a Lorentz force on a moving charge. This force is proportional to the square of electric field and the spin current with spin projected along the field. This is a purely relativistic quantum mechanical effect. As the origin of the force the spin current should be also observable physically. From the solution of the motion of a single electron in an electric field, the zitterbewegung of electronic wave packet in the spin-orbit coupling can be regarded as an explicit consequence of this force. Due to the similarity of this spin transverse force and the Lorentz force, the spin transverse force plays a similar role in the formation

of the charge Hall effect driven by the spin current and the spin Hall effect driven by the charge current as the Lorentz force does in the Hall effect in a magnetic field.

The author thanks F. C. Zhang for helpful discussion. This work was supported by the Research Grant Council of Hong Kong (No.: HKU 7039/05P), and a CRCG grant of The University of Hong Kong.

-
- [1] G. A. Prinz, Science 282, 1660 (1998); S. A. Wolf et al, Science 294, 1488 (2001); D. Awschalom, D. Loss, and N. Samarth (eds), Semiconductor Spintronics and Quantum Computation (Springer, Berlin, 2002).
 - [2] Datta and Das, Appl. Phys. Lett. 56, 665 (1990)
 - [3] M. I. D'yakonov and V. I. Perel, JETP Lett. 13, 467 (1971); J. E. Hirsch, Phys. Rev. Lett. 83, 1834 (1999); J. Sinova et al, ibid. 92, 126603 (2004); S. Murakami et al, Science 301, 1348 (2003); S. Q. Shen et al, Phys. Rev. Lett. 92, 256603 (2004).
 - [4] The Hall Effect and its Applications, edited by C. L. Chien and C. R. Westgate (Plenum, New York, 1980)
 - [5] J. Anandan, Phys. Lett. A 138, 347 (1989); R. C. Casella and S. A. Werner, Phys. Rev. Lett. 69, 1625; A. V. Balatsky and B. L. Altshuler, Phys. Rev. Lett. 70, 1678 (1993); C. M. Ryu, Phys. Rev. Lett. 76, 968 (1996)
 - [6] R. Winkler, Spin-orbit coupling effects in two-dimensional electron and hole systems, (Springer, Berlin, 2003)
 - [7] L. P. Rokhinson et al, Phys. Rev. Lett. 93, 146601 (2004)
 - [8] P. Zhang et al, cond-mat/0503505.
 - [9] E. I. Rashba, Fiz. Tverd. Tela (Leningrad) 2, 1224 (1960) [Sov. Phys. Solid State 2, 1109 (1960)]; J. Luo et al, Phys. Rev. B 38, R10142 (1988).
 - [10] J. Nitta et al, Phys. Rev. Lett. 78, 1335 (1997).
 - [11] S. Q. Shen, Phys. Rev. B 70, 081311(R) (2004).
 - [12] E. Schrodinger, Sitzungsber. Preuss. Akad. Phys. Math. Kl. 24, 418 (1930); H. Feshbach and F. Villars, Rev. Mod. Phys. 30, 24 (1958).
 - [13] J. Schliemann et al, Phys. Rev. Lett. 94, 206801 (2005)
 - [14] Z. F. Jiang et al, cond-mat/0410420
 - [15] A. Reynoso et al, Phys. Rev. B 70, 235344 (2004); B. K. Nikolic et al, cond-mat/0503415
 - [16] D. Dresselhaus, Phys. Rev. 100, 580 (1955)
 - [17] J. Li, L. Hu, and S. Q. Shen, Phys. Rev. B 71, 241305(R) (2005)
 - [18] E. Ya. Sherman et al, Appl. Phys. Lett. 86, 122103 (2005)
 - [19] E. M. Hankiewicz et al, cond-mat/0505597/Phys. Rev. B 72, 155305 (2005).
 - [20] R. Karplus and J. M. Luttinger, Phys. Rev. 95, 1154 (1954); J. Smith, Physica (Amsterdam) 21, 887 (1955); L. Berger, Phys. Rev. B 2, 4559 (1970)
 - [21] E. N. Bulgakov et al, Phys. Rev. Lett. 83, 376 (1999)
 - [22] J. Inoue et al, Phys. Rev. B 70, 041303(R) (2004); E. G. Mishchenko et al, Phys. Rev. Lett. 93, 226602 (2004)
 - [23] R. Winkler, Phys. Rev. B 62, 4245 (2000); R. Winkler et al, ibid. 65, 155303 (2002); J. Schliemann and D. Loss, ibid. 71, 085308 (2005)