

Universal properties of three-dimensional magnetohydrodynamic turbulence: Do Alfvén waves matter?

Abhik Basu¹ [*] and Jayanta K Bhattacharjee²

¹Max-Planck-Institut für Physik Komplexer Systeme,

Nothnitzer Strasse 38, D-01187, Dresden, Germany,

and Poornaprajna Institute of Scientific Research, Bangalore, India

²Department of Theoretical Physics,

Indian Association for the Cultivation of Sciences,

Jadavpur, Calcutta 700032, INDIA.

(Dated: March 23, 2024)

Abstract

We analyse the effects of the propagating Alfvén waves, arising due to non-zero mean magnetic fields, on the nonequilibrium steady states of three-dimensional (3d) homogeneous magnetohydrodynamic (MHD) turbulence. In particular, the effects of Alfvén waves on the universal properties of 3d MHD turbulence are studied in a one-loop self-consistent mode-coupling approach. We calculate the kinetic- and magnetic energy-spectra. We find that even in the presence of a mean magnetic field the energy spectra are Kolmogorov-like, i.e., scale as $k^{-5/3}$ in the inertial range where k is a Fourier wavevector belonging to the inertial range. We also elucidate the multiscaling of the structure functions in a log-normal model by evaluating the relevant intermittency exponents, and our results suggest that the multiscaling deviations from the simple Kolmogorov scaling of the structure functions decrease with increasing strength of the mean magnetic field. Our results compare favourably with many existing numerical and observational results.

PACS numbers: 64.60Ht, 05.70.Ln, 47.65+a

I. INTRODUCTION

The effects of propagating waves on the statistical properties of systems out of equilibrium remain an important topic of discussion. In the context of a coupled spin model in one-dimension ($d = 1$) [1] it has been shown that the presence of such waves leads to weak dynamic scaling in that model. In contrast, in a coupled Burgers-like model in $d = 1$ propagating waves do not affect the scaling properties of the correlation functions at all [2]. So far such issues have been considered only within very simplified one-dimensional nonequilibrium models [2, 3]. Magnetohydrodynamic (3dMHD) turbulence, which is a hydrodynamic description of the coupled evolution of the velocity fields u and the magnetic fields b in a quasi-neutral plasma, stands as a very good candidate for a natural system with propagating waves in three dimensions as most of its natural realizations have propagating Alfvén waves arising due to the presence of mean magnetic fields. Examples of such physical situations include solar wind, neutral plasma in fusion confinement devices etc. The presence of the propagating Alfvén modes, in addition to the usual dissipative modes due to the fluid and magnetic viscosities makes MHD turbulence a good natural example to study the interplay between the propagating and the dissipative modes in a system and their combined effects on the scaling properties of the correlation and structure functions, which are important issues from the point of view of nonequilibrium statistical mechanics.

The scaling of magnetic- $[E_b(k)]$ and kinetic- $[E_u(k)]$ energy spectra in the inertial range (i.e., wavevector k lies in the region $L^{-1} < k < \nu_D^{-1}$, L and ν_D being the integral scale given by the system size and the dissipation scale, respectively) in 3dMHD in the presence of Alfvén waves originating due to a mean magnetic field B_0 remains controversial till the date. Numerical simulations, due to the lack of sufficient resolutions failed to conclusively distinguish between the Kolmogorov's and Kraichnan's predictions (see below). In this communication, within one-loop self-consistent mode-coupling (SCMC) approximations, we obtain the following results for homogeneous but anisotropic (due to the mean magnetic field) 3dMHD :

The bare Alfvén wave speed, proportional to B_0 , renormalizes to acquire a singular k -dependence to become $B_0 k^{-1/3}$ where k is a Fourier wavevector belonging to the inertial (scaling) range. From this result we are able to conclude that even in the presence of a mean magnetic field the kinetic- and the magnetic- energy spectra scale

as $k^{-5/3}$ in the inertial range, identical with the situation without any mean magnetic field.

The dimensionless Kolmogorov's constants for the kinetic- and magnetic-energy spectra depends on a dimensionless parameter which we identify as the ratio of the renormalised Alfvén wave speed and the renormalised viscosities (see below).

The intermittency exponents of the Elsässer fields $z = u, b$ which approximately and qualitatively characterise the multiscaling properties of the structure functions in a log-normal model depend on the parameter and decrease with increasing .

Thus our results show that Alfvén waves in 3dMHD do not affect the scaling properties of the two-point correlation functions. However, the multiscaling properties of the structure functions are shown to be affected by the mean magnetic fields.

The famous Kolmogorov's arguments [4] for fluid turbulence can be easily extended to 3dMHD turbulence. In MHD, in the unit where mass density $\rho = 1$, the kinetic energy dissipation rate per unit mass (ϵ_K) and the magnetic energy dissipation rate per unit mass (ϵ_M) have same physical dimensions. Thus, as in fluid turbulence, by claiming that the structure functions of the velocity and magnetic field differences in the inertial range must be constructed out of the mean energy dissipation rate per unit mass and the local length scale r (belonging to the inertial range) one obtains for the n -th order structure function [5, 6]

$$S_n^a(r) = \langle [a(x+r) - a(x)]^n \rangle \sim (\epsilon r)^{n/3}; \quad \epsilon_D \ll r \ll L \quad (1)$$

where $\epsilon = \epsilon_K$ or ϵ_M and $a = u, b$ for the velocity and the magnetic fields, ϵ_D is the (small) dissipation scale and L is the (large) system size. This yields, for the energy spectra in the inertial range (as a function of wavevector k)

$$E_a(k) \sim k^{-5/3}; \quad a = u, b \quad (2)$$

This is known as the K41 theory in the relevant literature. However, in the presence of a mean magnetic field B_0 , the Alfvén waves are generated (see below) with the propagation speed B_0 . Thus in such a case in addition to the usual Kolmogorov time scale $\tau \sim k^{-2/3}$ [7, 8], there exists another time scale constructed out of the mean magnetic field B_0 , known as the Kraichnan time scale $(B_0 k)^{-1}$ [9]. Kraichnan argued that this time scale would determine

the energy cascade process and hence would enter in the expression of the structure function yielding $E_a(k) \propto k^{-3/2}$; $a = u; b$. There has not been any satisfactory resolution of this issue till this date; due to the particularly difficult vectorial nature of the 3d MHD equations (see below) it is rather difficult to achieve high Reynolds number in Direct Numerical Solutions (DNS) of the 3d MHD equations. Numerical solutions of an MHD shell model in the presence of a small mean magnetic field-like term did not find any dependence of the multiscaling of the structure functions on the mean magnetic field [6]. Analytically, it has been shown within the context of a 1d coupled Burgers-like model [2] that the energy spectra are independent of a mean magnetic field and in case of 1d MHD turbulence it exhibits the Kolmogorov scaling for the energy spectra. Similar conclusions followed from an analogous one-dimensional model [3]. Various phenomenological approaches, including weak turbulence theories and three-wave interaction models, yield, in general, mean magnetic field dependences of the energy spectra [10, 11, 12, 13] when the mean magnetic field is strong. However, these theories, despite their predictions are either not directly derivable from the underlying 3d MHD Eqs. of motion or involve additional assumptions on the flows. Analyses starting from the 3d MHD Eqs. in this regard are still lacking. Most observational results on astrophysical systems seem to favour the K41 results [14]. Simulations of incompressible MHD [15] find results close to the K41 result. Simulations of compressible MHD [16], even though in some cases find energy spectra closer to K41, in general yields a less clear picture. Recent numerical results of Muller et al [17] suggest that in presence of finite magnetic helicity structure functions parallel and perpendicular to the mean magnetic fields are affected differently by the mean magnetic field. In this paper, we address some of these issues by starting from the 3d MHD Eqs. without making any further assumptions on the velocity and the magnetic fields, except for the validity of the perturbative approaches. In particular, we show by applying one-loop mode coupling methods on the 3d MHD equations with a mean magnetic field and in the absence of any magnetic helicity that the one-dimensional energy spectra in 3d MHD turbulence are independent of the mean magnetic field B_0 and scale as $k^{-5/3}$ in the inertial range where k is a Fourier wavevector belonging to the inertial range. We then proceed to calculate the Kolmogorov's constants for the kinetic- and the magnetic-energy spectra and show that they depend on B_0 . Lastly, we calculate the intermittency exponents for the velocity and the magnetic fields and find that they decrease with increasing B_0 . We do not distinguish between the longitudinal and the transverse structure functions. Even

though for analytical convenience we assume a weak mean magnetic field, we are able to obtain new and interesting results concerning scaling and multiscaling in 3dMHD in the presence of a mean magnetic field of small magnitude. In this respect our results can be considered as complementary to some of the existing results [10, 11, 13]. The rest of the paper is organised as follows: In Sec.II we discuss the 3dMHD equations for incompressible fluids. In Sec.V we show that for incompressible fluids the bare Alfvén wave speed B_0 renormalizes to pick a correction $k^{-1=3}$ in the inertial range. This, as we argue, implies that the energy spectra scale as $k^{-5=3}$ in the inertial range even in the presence of a mean magnetic field. In Sec.VI we calculate the Kolmogorov's constants for the kinetic and the magnetic energy spectra. We introduce a parameter β which is the dimensionless ratio of the renormalised mean magnetic field and renormalised viscosities and show that the Kolmogorov's constants depend on β . In Sec.VII we elucidate the multiscaling properties of the structure functions by calculating the intermittency exponents and show that they decrease with increasing β , i.e., with increasing B_0 . This suggests that a mean magnetic field tends to reduce multiscaling corrections to the K41 results for the structure functions. In Sec.VIII we summarize our results.

II. MODEL EQUATIONS

We begin by writing down the 3dMHD equations for the velocity fields u and the magnetic fields b : The velocity field u is governed by the Navier-Stokes equation modified by the inclusion of the Lorentz force [18]

$$\frac{\partial u}{\partial t} + (u \cdot \nabla) u = \frac{r}{\rho} \nabla^2 u + \frac{(r - b)}{\rho} \nabla \cdot b + r^2 u + f; \quad (3)$$

and the dynamics of the magnetic field b is governed by the Induction equation [18] constructed out of the Ohm's law for a moving frame and the Ampère's law:

$$\frac{\partial b}{\partial t} = r \cdot (\nabla \times b) + r^2 b + g; \quad (4)$$

Here, ρ is the mass density, p is the pressure, r is the fluid viscosity and η is the magnetic viscosity (inversely proportional to the electrical conductivity of the fluid medium concerned). Functions f and g are external forces needed to maintain a statistical steady state. In the present approach these are taken to be stochastic forces. We assume them to be zero-mean

and Gaussian distributed with specified variances (see below). In addition to Eqs. (3) and (4) we also have $\nabla \cdot \mathbf{b} = 0$ (Maxwell's equation) and, for incompressible fluids $\nabla \cdot \mathbf{u} = 0$.

If the magnetic fields $\mathbf{b}(\mathbf{x};t)$ are such that $\langle \mathbf{b} \rangle = \mathbf{B}_0$ (a constant vector) then replacing \mathbf{b} by $\mathbf{b} + \mathbf{B}_0$ where now $\langle \mathbf{b} \rangle = 0$, in Eqs. (3) and (4) one obtains additional linear terms proportional to wavevector \mathbf{k} leading to wave-like excitations, known as Alfvén waves. The resulting equations are

$$\frac{\partial \mathbf{u}}{\partial t} + \alpha_1 (\mathbf{u} \cdot \nabla) \mathbf{u} = \frac{\nabla p}{\rho} + \alpha_2 \frac{(\nabla \cdot \mathbf{b})}{\mu} \mathbf{b} + \frac{(\nabla \cdot \mathbf{b})}{\mu} \mathbf{B}_0 + \nu \nabla^2 \mathbf{u} + \mathbf{f}; \quad (5)$$

and

$$\frac{\partial \mathbf{b}}{\partial t} = \alpha_3 \nabla \times (\mathbf{u} \times \mathbf{b}) + \eta \nabla^2 \mathbf{b} + \mathbf{g}; \quad (6)$$

with $\langle \mathbf{b}(\mathbf{x};t) \rangle = 0$. The parameters $\alpha_1; \alpha_2; \alpha_3$ are kept for book keeping purposes and can be set to unity (see Sec.IV). Note that on dropping the nonlinear and the dissipative terms from Eqs. (5) and (6) the resulting linear coupled partial differential equations admit wave-like solutions with dispersion relation linear in wavevector \mathbf{k} . These are known as the Alfvén waves [19] in the literature which propagate with speed proportional to B_0 . In the following sections we would calculate the kinetic and the magnetic energy spectra, the Kolmogorov's constants and the intermittency exponents in the presence of the Alfvén waves, i.e., for $B_0 \neq 0$.

III. CORRELATION AND STRUCTURE FUNCTIONS IN MHD

In the statistical steady state the time dependent correlation functions of \mathbf{u} and \mathbf{b} exhibit scaling which are characterized by the roughness exponents α_u and α_b respectively, and the dynamic exponent z . In terms of the scaling exponents $\alpha_u; \alpha_b$ and z the velocity and the magnetic field correlators have the form (as a function of wavevector \mathbf{k} and frequency ω)

$$\begin{aligned} C_{ij}^u(\mathbf{k}; \omega) &= \langle u_i(\mathbf{k}; \omega) u_j(-\mathbf{k}; -\omega) \rangle = D^u P_{ij} k^{d-2-\alpha_u-z} f_u(k^z = \omega); \\ C_{ij}^b(\mathbf{k}; \omega) &= \langle b_i(\mathbf{k}; \omega) b_j(-\mathbf{k}; -\omega) \rangle = D^b P_{ij} k^{d-2-\alpha_b-z} f_b(k^z = \omega); \end{aligned} \quad (7)$$

where f_u and f_b are scaling functions, P_{ij} is the transverse projection operator: $P_{ij} = \delta_{ij} - \frac{k_i k_j}{k^2}$ to account for the incompressibility of the fields. The kinetic- and magnetic-energy are simply related to the correlation functions: They are just the equal time velocity

and magnetic field auto correlation functions multiplied by appropriate phasefactors. We now set out to find whether the Alfvén waves are relevant perturbations on the system. It should be noted that the correlators (7) are chosen as they would be in the fully isotropic case. In the presence of a mean magnetic field B_0 there would however be additional anisotropic terms in the expression for the correlators above. In the small B_0 limit the lowest order corrections to the expressions (7) are $O(B_0)^2$. Neglecting these correction terms does not affect our scaling analyses below, and in any case, we are interested to see whether anisotropic Alfvén waves are relevant perturbations in the large scale, long time limit on the isotropic 3dMHD as characterised by expressions (7). Therefore, it suffices for us to work with the expressions (7) for our mode-coupling analyses below.

A complete characterisation of the nonequilibrium steady state (NESS) of MHD, however, requires informations about, not just the scaling of the energy-spectra, but also of the n -th order equal time structure functions of the velocity and the magnetic fields in the inertial range. These are defined by

$$S_n^a(r) = \langle [a(x+r) - a(x)]^n \rangle; \quad a = u, b; \quad \delta_D \ll r \ll L; \quad (8)$$

which scale as $r^{-\frac{n}{3}}$ where r belongs to the inertial range ($\delta_D \ll r \ll L$). According to the Kolmogorov theory [K41] the multiscaling exponents $\frac{n}{3} = n/3$; i.e., they are linear in n . Subsequent numerical and observational results for homogeneous and isotropic 3dMHD, i.e., without any mean magnetic fields, [20, 21] suggested deviations from the K41 results which are similar to those of pure fluid turbulence [22]. Much less results for the multiscaling of the structure functions are available when there are mean magnetic fields. Recent numerical simulations of Müller et al [17] suggested that structure functions parallel and perpendicular to the mean magnetic fields are differently affected by it when there is a finite magnetic helicity. Below we investigate the issue of multiscaling in presence of a mean magnetic field B_0 within the context of a log-normal model by evaluating the relevant intermittency exponents which are found to depend explicitly on B_0 . In our analyses we ignore the distinction between the structure functions parallel and perpendicular to the direction of the mean magnetic field; nevertheless, as we discuss below, our results are significantly new.

IV . SYMMETRIES OF THE EQUATIONS OF MOTION

We begin by re-expressing 3dMHD Eqs. (3) and (4) by writing the magnetic elds \mathbf{b} as a sum of a space-time dependent part and a constant vector: $\mathbf{b}(\mathbf{x};t) = \mathbf{b}(\mathbf{x};t) + \mathbf{B}_0$. In terms of these elds and parameters, the 3dMHD Eqs. become

$$\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} = \frac{\mathbf{r} \cdot \mathbf{p}}{\rho} + \frac{(\mathbf{r} \cdot \mathbf{b})}{\rho} \mathbf{b} + \frac{(\mathbf{r} \cdot \mathbf{b})}{\rho} \mathbf{B}_0 + \nu \nabla^2 \mathbf{u} + \mathbf{f}; \quad (9)$$

and

$$\frac{\partial \mathbf{b}}{\partial t} = \nabla \times (\mathbf{u} \times \mathbf{b}) + \eta \nabla^2 \mathbf{b} + \mathbf{g}; \quad (10)$$

It should be noted that in Eqs. (9) and (10) \mathbf{B}_0 is not the mean magnetic eld in general; it would be so only if $\langle \mathbf{b}(\mathbf{x};t) \rangle$ is zero in Eqs. (9) and (10). Note that our above way of splitting the magnetic elds does not change the actual mean magnetic eld in the system: It is still given by $\mathbf{B}_0 + \langle \mathbf{b} \rangle = \mathbf{B}_0$. The Eqs. of motion (9) and (10) are invariant under the following continuous transformations [8, 23]:

The Galilean transformation (TI): $\mathbf{u}(\mathbf{x};t) \rightarrow \mathbf{u}(\mathbf{x} + \mathbf{v}t; t) + \mathbf{u}_0$; $\frac{\partial}{\partial t} \rightarrow \frac{\partial}{\partial t} - \mathbf{v} \cdot \nabla$; and $\mathbf{b} \rightarrow \mathbf{b}$ [2, 8, 24] with $\gamma_1 = \gamma_3 = 1$ in Eqs. (5) and (6). This implies non-renormalization of γ_1 [2, 24, 25].

The transformation (TII) $\mathbf{B}_0 \rightarrow \mathbf{B}_0 + \mathbf{c}$, $\mathbf{b}(\mathbf{x};t) \rightarrow \mathbf{b}(\mathbf{x};t) - \mathbf{c}$, $\mathbf{u} \rightarrow \mathbf{u}$. This allows one to work with the effective magnetic elds defined by $\mathbf{b} - \mathbf{c}$ such that the coefficient of the Lorentz force vertex constructed out of the effective magnetic elds does not renormalize. This, therefore, ensures γ_2 can be set to unity [8, 23] by treating all magnetic elds as effective elds. Here the shift \mathbf{c} is a vector. A transformation similar to TII above exists in a problem of passive scalar turbulence [26].

The transformation TII essentially signifies the freedom to split the total magnetic elds as a sum of a constant part and a space-time dependent part. It should be noted that the transformation TII keeps the mean magnetic eld unchanged. In fact, nonrenormalization of γ_2 can be shown in a simpler way: Let us assume that under mode eliminations and rescaling $\mathbf{b} \rightarrow \mathbf{b}$. This scale factor of \mathbf{b} can now be absorbed by redefining the units of the magnetic elds by $\mathbf{b} \rightarrow \mathbf{b}$. Therefore, γ_2 can be set to unity if all magnetic elds are considered as effective or rescaled magnetic elds. Since the Induction Eq. (4) is linear in

the magnetic field $b(\mathbf{x};t)$, such rescaling leaves every conclusion unchanged. Of course, the external force g is also scaled by a factor l^p — which does not affect our analysis here as the assignment of canonical dimensions to various fields and parameters is done after absorbing l^2 in the definition of b . More specifically under the rescaling $\mathbf{x} \rightarrow l\mathbf{x}; t \rightarrow l^z t; u_i \rightarrow l^u u_i; b_i \rightarrow l^b b_i$ the bare parameters scale as $\nu_{1,3}(l) \rightarrow l^{u+z-1} \nu_{1,3}; \eta_2(l) \rightarrow l^{2-b-u+z-1} \eta_2$ and $B_0 \rightarrow l^{b-u+z-1} B_0$. Since there are no fluctuation corrections to the nonlinearities $\nu_1; \eta_2; \eta_3$, which are the consequences of the invariances under the transformations T I and T II, they can be kept invariant under rescaling of space and time as mentioned above leading to $u = b = 0$ and $u + z = 1$. Thus, under naive rescaling the bare parameter $B_0 \rightarrow B_0 l^{z-1}$. Furthermore, the invariance under the transformation T II and the resulting Ward identity ensure that different ways of implementing one-loop RG by having different values for $h b_i$ and B_0 in Eqs. (9) and (10) subject to the same bare mean magnetic field $B_0 = B_0 + h b_i$. Since any renormalisation scheme must respect a freedom of choice as represented by the transformation T II, respective terms must scale the same way under the rescaling of space and time [23]. Hence, due to fluctuation corrections $B_0(l) \rightarrow 1$. Furthermore, since both B_0 and b have the same scaling dimensions given by the exponent b , the physical mean magnetic field in the system $B_0 = B_0 + h b(\mathbf{x};t)$, if it receives fluctuation corrections under mode eliminations, must scale as $B_0(l) \rightarrow 1$, or if there are no fluctuation corrections to it due to some special symmetries (as in the one-dimensional Burgers-like model for MHD in Ref.[2]) will be irrelevant (in an RG) sense as it will flow to zero as l^{z-1} (since z for fully developed turbulence is less than unity). This clearly suggests the possibility of a renormalisation of the Alfvén wave speed, akin to the renormalisation of the sound speed in compressible fluid turbulence [27]. As we will see below, for 3dMHD there are infra-red singular fluctuation corrections to the bare mean magnetic field leading it to renormalise in a way consistent with our predictions from the Ward identities above. It should be noted that our analyses above is independent of the strength of the bare value of the mean magnetic field. Although, in the discussion above we have rescaled space isotropically (i.e., $x; y$ and z coordinates are rescaled the same way) and analyze the scale dependence of the resulting effective parameters and the fields, it should be noted that such a rescaling does not imply that the effective parameters have isotropic structures.

V. ENERGY SPECTRA FOR INCOMPRESSIBLE FLUIDS

We begin by writing down the 3dMHD equations in the incompressible limit (i.e., $\nabla \cdot \mathbf{u} = 0$). We write down the equations of motion in \mathbf{k} -space in terms of the Elsasser variables $\mathbf{z} = \mathbf{u} \pm \mathbf{b}$. The equations are (we take the mean magnetic field \mathbf{B}_0 to be along the \hat{z} direction)

$$\begin{aligned} \frac{\partial \mathbf{z}_\pm^+ (\mathbf{k}; t)}{\partial t} &= i\mathbf{B}_0 k_z \mathbf{z}_\pm^+ + i\mathbf{P}_\perp \mathbf{k}_s \sum_{\mathbf{q}} \mathbf{z}_s^+ (\mathbf{q}; t) \mathbf{z}_\pm^+ (\mathbf{k} - \mathbf{q}; t) + \text{c.c.} + \text{noise}_\pm^+ (\mathbf{k}; t); \\ \frac{\partial \mathbf{z}_\pm^+ (\mathbf{k}; t)}{\partial t} &+ i\mathbf{B}_0 k_z \mathbf{z}_\pm^+ + i\mathbf{P}_\perp \mathbf{k}_s \sum_{\mathbf{q}} \mathbf{z}_s^+ (\mathbf{q}; t) \mathbf{z}_\pm^+ (\mathbf{k} - \mathbf{q}; t) + \text{c.c.} + \text{noise}_\pm^+ (\mathbf{k}; t) \end{aligned} \quad (11)$$

The stochastic forces noise_\pm^+ are linear combinations of $\mathbf{f}_1, \mathbf{g}_1$, the tensor \mathbf{P}_\perp is the transverse projection operator which appears due to the divergence-free conditions on \mathbf{u} and \mathbf{b} , and $\text{c.c.} = \text{complex conjugate}$.

In the absence of any crosscorrelations between the velocity and the magnetic fields, the simplest choice for the noise variances, consistent with the divergence-free conditions on the velocity and the magnetic fields, are

$$\begin{aligned} \langle \text{noise}_\pm^+ (\mathbf{k}; t) \cdot \text{noise}_\pm^+ (\mathbf{k}; t) \rangle &= 2\mathbf{P}_\perp D_1 k^\gamma (t); \\ \langle \text{noise}_\pm^+ (\mathbf{k}; t) \cdot \text{noise}_\pm^+ (\mathbf{k}; t) \rangle &= 2\mathbf{P}_\perp D_2 k^\gamma (t); \end{aligned} \quad (12)$$

In Eqs.(12) we choose $\gamma > 0$. In particular, the choice of $\gamma = d$ in d -space dimensions ensures that the energy flux in the inertial range is a constant (up to a weak logarithmic dependence on wavevector k) which forms the basis of the K41 theory [4], and as a result, the parameters D_1 and D_2 pick up dimensions of energy dissipation rate per unit mass. Hence, we will use $\gamma = d$ in our calculations below. In experimental realisations of MHD, external forces act on the large scales. In analytical approaches, such forces are replaced by stochastic forces with variances given by Eqs. (12) for calculational convenience. It should however be noted that in numerical solutions of the Navier-Stokes equation driven by stochastic forces with variances similar to Eqs. (12) structure functions of the velocity fields are shown to exhibit multiscaling similar to the experimental results [28]. Our preliminary results from the numerical solutions of the isotropic 3dMHD equations (i.e., no mean magnetic fields) yield multiscaling similar to those obtained from the 3dMHD Eqs. driven by large-scale forces [29]. Although no such numerical results exist for the case with a mean magnetic field, it will presumably be true for such a situation also. In a stochastically driven Langevin description,

correlation functions are proportional to the noise correlations and hence it is necessary to force the Induction Eq. (6) stochastically, as is common in the literature (see, e.g., [2, 8] and references therein). For such noises in the absence of any mean magnetic field (i.e., for the isotropic case) it has been shown by using a one-loop mode coupling theory that the scaling exponents have values $\beta = 1/3$; $z = 2/3$ [8].

The Eqs. of motion (11) are coupled even at the linear level leading to a bare propagator matrix for the Eqs. (11) of the form (a 'o' refers to bare quantities)

$$G_o^{-1} = \begin{pmatrix} 0 & 1 \\ i! & -iB_o k_z + \frac{\omega}{2} k^2 \\ \frac{\omega}{2} k^2 & i! + iB_o k_z + \frac{\omega}{2} k^2 \end{pmatrix} A$$

We use a one-loop mode coupling theory which is conveniently formulated in terms of the self-energy matrix and the correlation functions given by Eqs. (7). The self-energy matrix is defined by $G^{-1} = G_o^{-1} - \Sigma$ where G is the renormalised propagator matrix. In terms of the scaling exponents β and z , and the renormalised parameters the self-energy matrix is given by

$$\Sigma = \begin{pmatrix} 0 & 1 \\ i! & -iB(k)k_z + \frac{\omega}{2} k^z \\ \frac{\omega}{2} k^z & i! + iB(k)k_z + \frac{\omega}{2} k^z \end{pmatrix} A$$

where $B(k)$ is the renormalised or the effective Alfvén wave speed. If there are diagrammatic corrections to the imaginary parts of $\omega(k; !)$ at frequency $! = 0$ which are singular in the infra-red limit, then $B(k)$ is different from B_o , the bare Alfvén wave speed, else they are the same.

There are two one-loop diagrams which contribute to the fluctuation corrections to $\omega(k; ! = 0) = -iB(k)k_z + \frac{\omega}{2} k^z$. These are shown in Fig.(1). These have both real and imaginary parts at frequency $! = 0$. Thus there are fluctuation corrections to the bare Alfvén wave speed which are singular in the small wavenumber limit. We assume, in the spirit of mode-coupling methods, $B(k) = B k^{-s}$ (we assume an isotropic scale dependence for $B(k)$ for simplicity which suffices our purpose of finding the scale dependence of the effective Alfvén wave speed). Clearly, if $s + 1 < z$, the small wavenumber limit of the problem is dominated by underdamped waves, if $s + 1 > z$ the Alfvén waves are damped out in the small wavenumber limit. In contrast if $s + 1 = z$ then in the small wavenumber limit both

the propagating and the dissipative modes are present. Our analyses in Sec.IV clearly suggest that if there are fluctuation corrections to B_0 , then the effective scale dependent Alfvén wave speed $B(l) \propto k^{-z}$ yielding $s = -z$. Furthermore, since $\epsilon + z = 1$ we have $s + 1 = z$ leading to the co-existence of the underdamped Alfvén waves and the dissipative modes in the large-scale, long-time limit. Note that this situation allows us to define a dimensionless parameter $B_+ = B_0 / \nu$ where B_0 and ν are the renormalised Alfvén speed and the viscosity respectively.

In the SCMC approach vertex corrections are neglected. Lack of vertex renormalisations in the zero wavevector limit in 3dMHD allows SCMC to yield exact relations between the scaling exponents ϵ and z as in the noisy Burgers/Kardar-Parisi-Zhang equation [25]. In this model the nonlinearities and the noise variances do not renormalise, thus leading to $z = 2/3$; $\epsilon = 1/3$, satisfying the exponent identity $\epsilon + z = 1$. In the present case, due to the singular nature (in the small wavevector limit) of the bare noise correlations (12), they do not pick up any further singular corrections to their scaling; however the amplitudes get modified (this is similar to the results in Ref.[8]). We denote the renormalized amplitudes by D and D^* , respectively. Furthermore, for the 3dMHD Eqs. (5) and (6) there are infra-red singular fluctuation corrections to B_0 [see the one-loop diagrams in Fig. (1)] leading to $B_0(l) \propto l^{-1}$ consistent with our arguments above. The SCMC approach involves consistency in the scaling and the amplitudes of the mode coupling equations. It should be noted that the exponent values $z = 2/3$; $\epsilon = 1/3$ satisfy the mode coupling integral equations regardless of the strength of the mean magnetic field. For amplitude consistency the one-loop integrals are required to be evaluated. Due their complicated structure, we evaluated them by assuming that the strength of the mean magnetic field is small. Therefore, only our amplitude relations and not the scaling exponents are affected by the approximation of small mean magnetic fields.

Demanding amplitude consistency in the mode coupling equations we obtain,

$$\begin{aligned}
 \frac{D}{B_+} &= \frac{D}{B_+} \left(1 - \frac{2}{d} + \frac{1}{d(d+2)} \right) + \frac{D^*}{d(d+2)} \left(\frac{2D}{B_+} \left(1 - \frac{2}{d} + \frac{1}{d(d+2)} \right) - \frac{2D^*}{B_+ d(d+2)(d+4)} \right); \\
 \frac{B}{B_+} &= B_0 \frac{D}{d(d+2)^3}; \\
 \frac{1}{1+\epsilon} &= \frac{(1+\epsilon^2) \left(1 - \frac{2}{d} + \frac{2}{d(d+2)} \right) + 0.5 \left(1 - \frac{2}{d} \right) + F_1(\epsilon)}{(1+\epsilon^2) \left(1 - \frac{2}{d} + \frac{2}{d(d+2)} \right) + 0.5 \left(1 - \frac{2}{d} \right) + F_2(\epsilon)};
 \end{aligned} \tag{13}$$




FIG . 1: One-loop diagrams contributing to renormalization of G_0^+ . A single line refers to a propagator and a line with a filled circle refers to a correlator.

where $F_1(\epsilon) = 2^2(1 + 3\epsilon^2)(1 - d + \frac{1}{d(d+2)} - \frac{1}{d(d+2)(d+4)})$; $F_2(\epsilon) = 2^2(3 + \epsilon^2)(1 - d + \frac{1}{d(d+2)} - \frac{1}{d(d+2)(d+4)})$ are the ϵ -dependent parts of the amplitude-ratio. While calculating the amplitude consistency relations we worked in the limit of small $B = +$ and the renormalised magnetic Prandtl number ($- = \frac{+}{++}$) close to unity. Away from these limits the amplitudes of the underlying one-loop integrals become much more complicated functions of ϵ and also of the renormalised magnetic Prandtl number, but the qualitative picture remains unchanged. The main physical picture that emerges from the expressions (13) is that in the absence of a bare mean magnetic field B_0 the effective Alfvén wave speed is also zero which is a restatement of the fact that if the original theory is isotropic, so will be the renormalised theory. Moreover the renormalised Alfvén speed increases with increasing B_0 . With these results we are now in a position to calculate the energy spectra in the inertial range. We use the form of the effective (scale dependent) viscosity and the Alfvén wave speed to obtain the equal time correlation function of z_i in the long wavelength

limit. We define (in 3d)

$$\begin{aligned}
C_{ij}^{++}(\mathbf{k};!) &= h z_i^+(\mathbf{k};!) z_j^+(\mathbf{k};!) i = \frac{2D k^3 P_{ij}(\mathbf{k})}{(i^2 + B k^{1=3} k_z)^2 + k^{4=3}}; \\
C_{ij}(\mathbf{k};!) &= h z_i(\mathbf{k};!) z_j(\mathbf{k};!) i = \frac{2D k^3 P_{ij}(\mathbf{k})}{(i^2 + B k^{1=3} k_z)^2 + k^{4=3}}; \\
C_{ij}^+(\mathbf{k};!) &= h z_i^+(\mathbf{k};!) z_j(\mathbf{k};!) i = \frac{2D k^{4=3} P_{ij}(\mathbf{k})}{[i(i^2 + B k^{1=3} k_z) + k^{2=3}][i(i^2 + B k^{1=3} k_z) + k^{2=3}]} \quad (14);
\end{aligned}$$

As discussed before, we have omitted anisotropic corrections of $O(B_o)^2$ or of $O(\epsilon)^2$ to the correlation functions as we are trying to find out the relevance (in an RG sense) of Alfvén waves on the scaling of the isotropic correlation functions in the absence of any mean magnetic fields. Therefore, the equal time correlation functions have the following form (in $d = 3$):

$$C_{ij}^{++}(\mathbf{k};t=0) = C_{ij}(\mathbf{k};t=0) = D k^{3-2=3} P_{ij}(\mathbf{k}); \quad (15)$$

Therefore, the equal time autocorrelation functions of z are independent of any mean magnetic field. This holds true regardless of the scaling of the noise variances. The equal time cross correlation function $C_{ij}^+(\mathbf{k};t=0)$ requires more careful considerations. On integrating $C_{ij}^+(\mathbf{k};!)$ over all frequency $!$ one obtains (in 3d)

$$C_{ij}^+(\mathbf{k};t=0) = \frac{D k^3 P_{ij}}{i B k^{1=3} k_z + k^{2=3}}; \quad (16)$$

a form which is valid in the inertial range. It is clear from the expression (16) that both the real and the imaginary parts of $C_{ij}^+(\mathbf{k};t=0)$ scale as $k^{3-2=3}$ in the inertial range. Thus the one-dimensional kinetic- and the magnetic-energy spectra (which are simply related to the correlators defined in Eqs. (14)) scale as $k^{5=3}$ in the inertial range. The emerging physical picture is as follows: We find from the expression for $C_{ij}^+(\mathbf{k};!)$ above that this, as a function of frequency $!$, has a maximum at $! = B k^{1=3} k_z$ and the width at half maximum $\propto k^{2=3}$. In contrast, the auto correlations of z^+ ; z are maximum at $! = 0$ and their widths scale as $k^{2=3}$. Thus in the long wavelength limit the width and the location of the maximum of $C_{ij}^+(\mathbf{k};!)$ scale in the same way leading to the presence of the underdamped Alfvén waves in the hydrodynamic limit. Therefore, it immediately follows that the kinetic- and the magnetic-energy spectra, being linear combinations of the correlators discussed above times appropriate phase factors, scale as $k^{5=3}$ even in the presence of a non-zero mean magnetic field. It should be noted that we have used a small approximation to arrive at our results for

the self-consistent amplitude relations. For a finite one would require to work with a fully anisotropic form of the correlation functions and obtain self-consistent relations for scaling and anisotropic amplitudes, a task, which is analytically challenging, remains to be done in the future. However, based on our calculations above, the exponent identity $\alpha + z = 1$ and the Ward identity suggesting that the renormalised Alfvén wave speed should scale as $k^{-1=3}$ with wavevector k being in the inertial range, we argue that even for a finite, i.e., a finite mean magnetic field the energy spectra will scale as $k^{-5=3}$ in the inertial range, a result supported by many observational evidences [14]. The self-consistent amplitude equations will then be anisotropic reflecting the presence of underdamped Alfvén waves in the inertial range. The full correlation matrix will be anisotropic in the hydrodynamic limit; its eigenvalues have different amplitudes, but all of them scale the same way. Our confidence on our result that the scaling of the correlation functions along directions parallel and perpendicular to the direction of the mean magnetic field is same is derived from the fact that our exponent values $z = 2=3$; $\alpha = 1=3$ satisfy the one-loop integral equations regardless of the strength of the mean magnetic field and are consistent with the Ward identity discussed above.

VI. KOLMOGOROV'S CONSTANTS

According to the Kolmogorov's hypothesis for fluid turbulence [4], in the inertial range energy spectrum $E(k) = K_0^{-2=3} k^{-5=3}$, where K_0 , a universal constant, is the Kolmogorov's constant and ϵ is the energy dissipation rate per unit mass. Various calculations, based on different techniques by different groups [7, 30, 31, 32] show that $K_0 \approx 1.5$ in three dimensions. Having noted that the energy spectra, even in the presence of a mean magnetic field scale as $k^{-5=3}$ extensions of Kolmogorov's hypothesis for 3dMHD allows one to define Kolmogorov's constants for the Elsasser fields: $E(k) = K_0^{-2=3} k^{-5=3}$. Since $z = u$ b, we have $h z_i^+(k;!) z_j^+(\mathbf{k};!) i = h z_i^-(k;!) z_j^-(\mathbf{k};!) i$ and $z_+ = z_- = z_{MHD}$ in absence of any crosscorrelations between the velocity and the magnetic fields, we have $K_0^+ = K_0^- = K_{MHD}$. The noise strength D and the rate of energy dissipation per unit mass is connected by the Novikov's theorem [8, 33]: $\epsilon = 2D \frac{S_3}{(\epsilon^2)^3}$. Noting that the energy spectra $E(k)$ of z in the inertial range is given by

$$E(k) = D k^{-3} = \epsilon k^{2=3}; \quad (17)$$

where D is the effective or renormalised noise strength, we identify

$$K_{MHD} = 1.6(1 + 0.7\beta^2)^{-6} \approx 105\beta^{-2} : \quad (18)$$

The notable feature of the expression (18) is that the constant K_{MHD} depends on the dimensionless parameter β which we introduced before. For $\beta = 0$, i.e., for no magnetic fields, we find $K_{MHD} = K_0 = 1.6$ for pure uid turbulence which is well within the accepted range of values [31]. Before closing this Sec. we would like to point out that the presence of multiscaling raises questions about Kolmogorov's constant K_{MHD} being universal: A small but finite intermittency correction (i.e., multiscaling) over the simple K41 scaling implies the presence of an arbitrary scale which may spoil the universality of K_{MHD} . We however refrain ourselves from getting into this question and adopt a point of view that regardless of whether or not K_{MHD} remains universal due to multiscaling, the numerical value of this constant is likely to get affected by the presence of Alfvén waves in the system which is reflected by the expression (18).

VII. POSSIBILITIES OF VARIABLE MULTIFRACTALITY

Experiments and numerical simulations [20, 34] find nonlinear multiscaling corrections to the K41 prediction of $\zeta_p^a = p-3$ for the structure functions in the inertial range. Until the date, no controlled perturbative calculation for ζ_p^a is available. To account for multiscaling in uid turbulence, however, Obukhov [35] and Kolmogorov [36] assumed a log-normal distribution for dissipation to arrive at

$$S_p^v(r) = \langle |v_j^v(r)|^p \rangle = C_p^{-p-3} r^{p-3} \frac{L}{r}^{\bar{\zeta} - 2(p-3)} ; \quad (19)$$

where $\bar{\zeta}$ is the mean value of ζ and (a bound for $\bar{\zeta}$ in uid turbulence has been discussed in Ref.[37]):

$$\langle |v(x+r)|^2 \rangle / \langle |v|^2 \rangle = r^2 \langle |v|^{\bar{\zeta}} \rangle : \quad (20)$$

For small $\bar{\zeta}$, $\bar{\zeta} < 9$. A standard calculation on the randomly stirred model yields intermittency exponent $\bar{\zeta} = 0.2$ [32] where $\bar{\zeta} = 9$, whereas the best possible estimate from experiments is 0.23 [32]. This model, despite having well-known limitations and difficulties [38], serves as a qualitative illustration of multiscaling. As in uid turbulence, in MHD the dissipations of the Elsasser variables z fluctuate in space and time, and as a result one may

define two intermittency exponents for them. In the present problem $\beta_+ = \beta_- = \beta_{MHD}$. Below we calculate the exponent β_{MHD} in a one-loop expression which will give us an estimate of the Alfvén wave speed-dependent deviation of the scaling of the structure functions from their simple-scaling values as predicted by the K41 theory. We closely follow Ref.[32]. We work with the self-consistent forms for the self-energies and correlation functions given above along with the consistency relations for the amplitude-ratios and β . Following Ref.[32], we find the dissipation correlation functions in 3d to be

$$\langle \delta u(x+r) \delta u(x) \rangle' = 12.4 \beta_{MHD}^2 K_{MHD}^2 \ln \frac{L}{r}; \quad (21)$$

with β being defined by the relation $\beta_+ = \beta_- = \beta_{MHD}$. From the self-consistent amplitude-relations (13) we find

$$\beta = 0.41 + 0.73 \beta^2 - 6 \beta^3 - 29 \beta^4 = 105 \beta^{1/3} - 1 - 0.5 \beta^4 (0.7 - 0.03 \beta)^{1/3}; \quad (22)$$

Thus, β which is a universal coefficient in ordinary fluid turbulence, varies with the parameter β_{MHD} , or with the Alfvén wave speed in MHD. Substituting the values of K_0 and β we find

$$\beta_+ = \beta_- = \beta_{MHD} = 0.21 + 0.73 \beta^2 - 6 \beta^3 - 29 \beta^4 = 105 \beta^{4/3}; \quad (23)$$

Thus we find that a decrease in the value of β with an increase in β_{MHD} , i.e., with increasing mean magnetic field. Despite the limited applicabilities of log-normal models in characterising multiscaling, we can conclude, from our expression (23) for the intermittency exponent in MHD in the presence of a mean magnetic field, that the intermittency corrections to the simple K41 scaling is likely to get affected (reduced in our calculations) in the presence of Alfvén waves. Further calculations and/or numerical simulations are needed to find the exact extent of the dependence of multiscaling on the mean magnetic field and the possibilities of anisotropic multiscaling for the structure functions parallel and perpendicular to the mean magnetic field as demonstrated recently in Ref.[17]. It should be noted that our conclusions on the multiscaling properties of the structure functions depend on a log-normal model for 3d MHD. Such a description, unfortunately, is unable to distinguish between the structure functions parallel and perpendicular to the direction of the mean magnetic field in the system. Moreover, the intermittency exponents above [expressions (23)] are evaluated in the lowest order in mean magnetic field. Therefore, even though from our results we are not able to make firm comments on the possibility of parallel and perpendicular structure

functions exhibiting different multiscaling, the real importance of our results lie in their elucidation of the multiscaling properties depending on the mean magnetic field.

V III. C O N C L U S I O N

In this paper we have considered the effects of the Alfvén waves on the statistical properties of the correlation functions of the velocity and the magnetic fields or the Elsasser fields. We considered the case when the velocity fields are incompressible. In a one-loop approximation we find that the effective or the renormalised Alfvén wave speed scales as $k^{-1/3}$ where the wavevector k is in the inertial range. This immediately yields that the energy spectra, even in the presence of a mean magnetic field, scale as $k^{-5/3}$ in the inertial range. We identify a dimensionless parameter η which is the ratio of the effective Alfvén wave speed and the renormalised viscosity. We obtain self-consistent relations between the amplitude ratio of the correlation functions and η . These relations allow us to calculate the dimensionless Kolmogorov's constant and we show that it depends explicitly on η or on the mean magnetic field. Finally, we calculate the intermittency exponent which in a log-normal model gives a qualitative account of the multiscaling in terms of the deviation from the K41 scaling for the structure functions. We would like to emphasize that although the one-dimensional Burgers-like model of MHD of Ref.[2] and the 3dMHD Eqs. yield energy spectra independent of the Alfvén waves, the long wavelength physical pictures are different. In the former case, due to the nonrenormalization of the bare Alfvén wave speed, Alfvén waves are overdamped in the hydrodynamic limit; the dominant process in that limit is viscous dissipation. In contrast, in 3dMHD, the Alfvén wave speed picks up a singular correction in the hydrodynamic limit leading to the K41 scaling of the energy spectra and the presence of underdamped Alfvén waves in the hydrodynamic limit. Despite similar mathematical structures these crucial differences arise principally because the one-dimensional model decouples completely when written in terms of the Elsasser variables, allowing to come with the waves of each of them separately. As a result correlators of each of them are independent of the Alfvén waves and the bare Alfvén wave speed remains unrenormalised leading to overdamped Alfvén waves in the hydrodynamic limit. However, the 3dMHD equations do not decouple when written in terms of the Elsasser variables and hence oppositely propagating waves cannot be made to vanish by commuting. Despite the limitations of the one-loop methods [7] and the small

approximation to facilitate easier analytical manipulations for the amplitude relations, we obtain results which are significantly new and open the intriguing possibilities of MHD multiscaling universality classes being parametrised by the mean magnetic field. Some of the quantitative details will change if one retains terms which are higher order in ϵ ; however, we believe that the qualitative picture will essentially remain the same. A SMHD turbulence forms a natural example of a driven nonequilibrium system with Alfvén waves, our results are the first of its kind for a natural system. In our log-normal model approach, we did not distinguish between the longitudinal and the transverse structure functions our results cannot be compared directly with those of Ref.[17], where the longitudinal and the transverse structure functions are found to scale differently with multiscaling exponents which depend on the magnitude of the mean magnetic field in the presence of a finite magnetic helicity. In contrast our results apply to the multiscaling of the usual structure functions in the absence of any magnetic helicity which are combinations of the transverse and the longitudinal structure functions as considered in Ref.[17] which would also then exhibit a mean magnetic field dependent scaling. In particular we find that the usual structure functions multiscale less in presence of an increasingly strong mean magnetic field which is in qualitative agreement with those of Ref.[17]. Although here we have restricted ourselves to the study of Alfvén waves as relevant perturbations on the amplitude and the scaling of the isotropic correlation functions in the limit of small ϵ , our results indicate the possibility of the multiscaling exponents varying with the amplitude of the mean magnetic field. Our analyses, when extended for finite values of ϵ and for full anisotropic structure of the correlation functions, are likely to provide further understanding of and resolve some of the discrepancies between the various phenomenological scenarios available for situations with large mean magnetic fields [10, 11, 13]. We leave this task for the future.

From a broader point of view our results demonstrate the critical role of wave-like excitations in determining the statistical properties of 3dMHD. The presence of propagating waves is not confined to MHD only; they are a generic feature in many other naturally occurring soft-matter systems where such waves can be present in a viscous environment, e.g., active polar gels in cytoskeletal dynamics [39] and in the dynamics of self-propelled particles [40] etc. We believe our results will lead to similar theoretical and experimental studies in relevant nonequilibrium soft-matter systems.

IX . ACKNOWLEDGEMENT

This work was started when one of the authors (AB) was an Alexander von Humboldt Fellow at the Hahn Meitner Institut, Berlin. AB wishes to thank the AvH Stiftung, Germany for partial financial support.

[*] email: abhik@mpipks-dresden.mpg.de

- [1] D. Das, A. Basu, M. Barmann and S. Ramaswamy, *Phys. Rev. E*, **64**, 021402 (2001).
- [2] A. Basu, J. K. Bhattacharjee and S. Ramaswamy, *Eur. Phys. J. B*, **9**, 425 (1999).
- [3] M. K. Verma, *Phys. Plasmas*, **6**, 1455 (1999).
- [4] A. N. Kolmogorov, *C. R. Acad. Sci. USSR* **30**, 301 (1941).
- [5] P. Goldreich and S. Sridhar, *Astrophys. J.*, **438**, 763 (1995).
- [6] A. Basu, PhD thesis, Indian Institute of Science, Bangalore (2000).
- [7] V. Yakhot and S. A. Orszag, *Phys. Rev. Lett.* **57**, 1722 (1986).
- [8] A. Basu, *Europhys. Lett.*, **65**, 505 (2004).
- [9] R. H. Kraichnan, *Phys. Fluids*, **8**, 1385 (1965).
- [10] S. Galtier et al, astro-ph/0008148.
- [11] C. S. Ng and A. Bhattacharjee, *Astrophys. J.*, **465**, 845 (1996).
- [12] J. V. Shebalin et al, *J. Plasma Phys.*, **29**, 525 (1983).
- [13] D. Montgomery and L. Turner, *Phys. Fluids*, **24**, 825 (1981).
- [14] B. J. Rickett and A. G. Lyne, *Mon. Not. R. Astron. Soc.*, **244**, 68 (1990); S. R. Spangler and C. R. Gwinn, *Astrophys. J.*, **353**, L29 (1990); Y. Gupta, B. J. Rickett and W. A. Coles, *Astrophys. J.*, **403**, 183 (1993); P. N. Wilkinson, R. Narayan and R. E. Spencer, *Mon. Not. R. Astron. Soc.*, **269**, 67 (1994).
- [15] J. Cho and E. T. Vishniac, *Astrophys. J.*, **539**, 273 (2000); J. Cho, A. Lazarian and E. T. Vishniac, *Astrophys. J.*, **564**, 291 (2002).
- [16] J. Cho and A. Lazarian, *Mon. Not. R. Astron. Soc.*, **345**, 325 (2003).
- [17] W.-C. Müller, D. Biskamp and R. G. Rappin, *Phys. Rev. E*, **67**, 066302 (2003).
- [18] J. D. Jackson, *Classical Electrodynamics*, 2nd edn., Wiley Eastern, New Delhi, (1975); A. Raichaudhuri, *The Physics of Fluids and Plasmas*, Cambridge University Press, Cambridge

- (1998).
- [19] D.Montgomery, Lecture Notes on Turbulence, eds. J.R.Herring and J.C.McWilliam, World Scientific, Singapore, 1989.
 - [20] A.Basu, A.Sain, S.K.Dhar and R.Pandit, Phys. Rev. Lett., 81, 2687 (1998).
 - [21] D.Biskamp and W.Muller, Phys. Plasmas, 7, 4889 (2000).
 - [22] K.R.Sreenivasan and R.A.Antonia, Ann. Rev. Fluid Mech., 29, 435 (1997); and S.K.Dhar, A.Sain, A.Pande, and R.Pandit, Pramana - J. Phys., 48, 325 (1997).
 - [23] A.Basu, unpublished.
 - [24] D.Forster et al, Phys. Rev A, 16, 732 (1977); J.K.Bhattacharjee, J. Phys. A, 31, L93 (1998).
 - [25] E.Frey and U.Tauber, Phys. Rev. E, 50, 1024 (1994).
 - [26] N.V.Antonov, Physica D, 144, 370 (2000).
 - [27] I.S.Taroselsky, V.Yakhot, S.Kida and S.A.Orszag, Phys. Rev. Lett., 65, 171 (1990); J.K.Bhattacharjee, Mod. Phys. Lett., 7, 881 (1993).
 - [28] A.Sain, Manu and R.Pandit, Phys. Rev. Lett., 81, 4377 (1998).
 - [29] A.Basu and R.Pandit, work in progress.
 - [30] R.H.Kraichnan, J. Fluid Mech., 47, 525 (1971).
 - [31] C.D.Leslie, Developments in the Theory of Turbulence, Clarendon Press, Oxford, 1973.
 - [32] A.Das and J.K.Bhattacharjee, Europhys. Lett., 26, 527 (1994).
 - [33] W.A.Woyczynski, Burgers-KPZ Turbulence-Göttingen Lectures, Springer-Verlag, Berlin, Heidelberg, 1998; W.D.McComb, The Physics of Fluid Turbulence, Clarendon Press, Oxford, 1990.
 - [34] J.Cho, A.Lazarin and E.T.Vishniac, ApJ, 564, 291 (2002); D.Biskamp and W.Muller, Phys. Plasmas, 7, 4889 (2000).
 - [35] A.M.Obukhov, J. Fluid Mech., 13, 77 (1962).
 - [36] A.M.Kolmogorov, J. Fluid Mech., 13, 82 (1962).
 - [37] C.R.Doring and C.Foias, J. Fluid Mech., 467, 289 (2002).
 - [38] U.Frisch, Turbulence: The Legacy of A.N.Kolmogorov (Cambridge University Press, Cambridge, 1995).
 - [39] K.Krusø et al, e-print: physics/0406058.
 - [40] R.A.Sinha and S.Ramaswamy, Phys. Rev. Lett., 89, 058101 (2002).